

Protective effect of Vitamin D on imidacloprid-induced testicular injury in rats

Hikmet Keles¹, Alper Yalcin², Hasan Aydin³

¹Department of Pathology, Faculty of Veterinary Medicine, Afyon Kocatepe University, Afyonkarahisar, Turkey

²Department of Histology, Faculty of Medicine, Adiyaman University, Adiyaman, Turkey

³Department of Pharmaceutical Toxicology, Faculty of Pharmacy, Adiyaman University, Adiyaman, Turkey

Submitted: 22 May 2019

Accepted: 21 June 2019

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms.2019.86776>

Copyright © 2019 Termedia & Banach

Corresponding author:

Assoc Prof. Hikmet Keles PhD

Department

of Pathology

Faculty of Veterinary

Medicine

Afyon Kocatepe

University

03030 Afyonkarahisar

Turkey

E-mail: hikmetkeles@gmail.

com

Abstract

Introduction: The degenerative effects of imidacloprid via oxidative stress are known. Irisin is a recently discovered peptide with energy regulator and antioxidant effects. In addition, the antioxidant potential of Vitamin D has been reported in previous studies. The current study was performed to investigate the effect of Vitamin D on testis morphology and irisin immunoreactivity in imidacloprid-treated rats.

Material and methods: Thirty-two Wistar albino male rats were divided into groups: control ($n = 6$), corn oil ($n = 6$), Vitamin D ($n = 6$), imidacloprid ($n = 7$) and imidacloprid + Vitamin D ($n = 7$). Testis tissues were used to evaluate the histopathological, biochemical and immunohistochemical changes. Oxidative state in testis tissue was determined with total antioxidant and oxidant status markers, total antioxidant status (TAS) and total oxidant status (TOS) respectively.

Results: In microscopic examination, degenerative changes in the seminiferous tubule epithelium, interstitial edema and increased irisin immunoreactivity were observed in animals given imidacloprid. Also increased TOS and decreased TAS levels were measured in these animals. It was observed that Vitamin D improved the testicular damage histopathologically when compared to the imidacloprid group. However, increase in TAS levels and decrease in both TOS levels and irisin immunoreactivity were found insignificant in animals given Vitamin D.

Conclusions: In the present study it was observed that Vitamin D ameliorated testis injury caused by imidacloprid. Furthermore, imidacloprid was found to increase the immunoreactivity of irisin. In the light of our findings, we conclude that the use of Vitamin D could be beneficial against testicular damage caused by imidacloprid.

Key words: vitamin D, imidacloprid, irisin, testis.

Introduction

Oxidative stress generated by insecticides causes the formation of reactive oxygen species (ROS) in living organisms and accordingly induces changes in antioxidant systems [1, 2]. Oxidative stress, which may cause an imbalance in the oxidant-antioxidant status, regulates inflammatory mediators and inhibits neutrophil infiltration [3], is effective in insecticide-induced tissue damage.

Neonicotinoids are known to have an important role as synthetic neuroactive insecticides and many of them are used in crops and livestock

for pest control [4]. Imidacloprid (IMC), one of the neonicotinoid insecticides, is widely used to eliminate pests in agriculture [5]. Residues of the IMC can be encountered in the food chain and therefore it is important to investigate the side effects of this insecticide. Previous studies have reported adverse effects of IMC in mammals [6–8]. Furthermore, side effects of IMC on reproductive ability have been reported in parenteral and offspring rats [9].

Vitamin D (VD) is a steroid hormone that is frequently incepted from dietary sources. In addition, it is also synthesized in the skin following exposure to UVB light [10, 11]. The balance between oxidants and antioxidants causes oxidative stress when disrupted in favor of oxidants. Vitamin D has both a pro- and anti-oxidative effect potential. Vitamin D deficiency is quite common worldwide and affects people of all ages [12]. The association of vitamin D deficiency with coronary artery disease has been demonstrated [13–15], and obese people are at greater risk [16]. The male reproductive system includes widely distributed vitamin D receptors (VDR) [17]. The expression of VDR and VD metabolizing enzymes in the male reproductive system indicates the function of the testis in VD production and regulation [18]. Vitamin D has important functions in the male reproductive system and VD deficiency has been reported to have significant effects on the male reproductive system [19]. Vitamin D deficiency (VD) and intake of high-fat diets may cause morphological and physiological changes related to infertility in the testis [20].

Irisin is a recently discovered peptide hormone and is synthesized in rats during exercise [21]. Moreover, it was determined that the level of irisin increased during oxidative stress status [22]. The peptide is found in skeletal and cardiac muscle tissue as well as brain, skin, spleen, testis, pancreas and stomach tissues [23]. Irisin regulates glucose and lipid homeostasis. Anti-apoptotic, anti-inflammatory and anti-oxidative properties of irisin have attracted the attention of scientists, since these properties are associated with the stages of many diseases such as cancer, myocardial infarction, lung injury, kidney diseases, atherosclerosis, obesity, liver diseases and type 2 diabetes [24]. Hence, it is important to find substances that could reverse the degenerative effects of IMC in mammalian tissues.

The aim of the present study is to examine the protective effects of VD in IMC-induced testis injury and to determine its possible effect on irisin immunoreactivity.

Material and methods

Animals, diets and experimental protocols

A total of 32 male Wistar albino rats aged 8–10 weeks (200–220 g), supplied by Adiyaman Univer-

sity Experimental Animal Production and Research Center, were divided into five groups: the control group ($n = 6$), corn oil (CO) group ($n = 6$), VD group ($n = 6$), IMC group ($n = 7$), and IMC + VD group ($n = 7$). The animals were maintained and fed ad libitum with standard rodent pellet diet and water in an ambient environment with 1/2 daylight/1/2 day dark cycle at $25 \pm 3^\circ\text{C}$ for 15 days for adaptation. The animals in the control group were maintained on a standard rodent pellet diet and water ad libitum without any treatment during the 8-week experimental period. The animals in the CO group were administered a daily dose of 2 ml of CO by oral gavage for 8 weeks. The VD group was given a dose of VD 200 IU/day via an oral dropper for 8 weeks. The IMC group was administered orally with 20 mg/kg IMC dissolved in CO (a total of 2 ml volume) per day for 8 weeks, while the IMC + VD group was administered orally 20 mg/kg IMC dissolved in CO (a total of 2 ml volume) and 200 IU/day VD via an oral dropper for 8 weeks. Testis tissues obtained from rats, which were anesthetized with ketamine and xylazine (75 and 10 mg/kg, respectively), were fixed in 10% buffered formalin. Some fresh tissue samples were stored at -20°C for total antioxidant status (TAS) and total oxidant status (TOS) measurements. This animal study was approved by the Ethical Committee of Experimental Animals of Firat University (Protocol number: 2017/67).

Biochemical methods

The testis tissues were washed with cold PBS buffer, homogenized in ice with 0.1 M phosphate buffer (1/10 ratio at pH 7.4) and centrifuged at 1800 rpm for 5 minutes for evaluating TAS and TOS levels. Tissue TAS [25] and TOS [26] levels were measured by using TAS and TOS Assay Test Kits (Rel Assay Diagnostic, Gaziantep, Turkey) with an autoanalyzer (Olympus AU2700) to assess the degree of oxidant damage.

Histopathological method

Formalin fixed tissue samples were processed and blocked with paraffin, and then sectioned into 5- μm sections and stained with hematoxylin and eosin (HE). These slides were blindly examined under a light microscope (Leica DM500 microscope –Leica DFC295).

Immunohistochemical method

Briefly, 4–6 μm thick sections taken from paraffin blocks were transferred on poly-L-lysine coated slides. Deparaffinized and re-hydrated sections were boiled in a microwave oven and placed at room temperature for about 20 min and transferred into hydrogen peroxide for blocking endog-

enous peroxides. Subsequently the sections were washed with PBS for 3 × 5 min and nonspecific immunoglobulins were blocked with Ultra V Block solution for 5 min. In the next step, irisin primary antibody (Irisin Rabbit Polyclonal H-067-17, Phoenix Pharmaceuticals, Inc., California, USA) (at a ratio of 1/200) was applied on the sections for 60 min. Subsequently, an anti-mouse/rabbit IgG secondary antibody was applied for 30 min at room temperature. These sections were next washed with PBS and were incubated with streptavidin peroxidase. After these processes the sections were visualized with 3-amino-9-ethylcarbazole (AEC) and the background was colorized with Mayer's Hematoxylin. The sections were covered with aqua medium. Manufacturers' proposed methods were applied for positive and negative control stains. The stained sections were stored at room temperature in the dark until examination. Histopathological scores for the immunoreactivity were counted using the prevalence interval (0.1: < 25%, 0.4: 26–50%, 0.6: 51–75%, 0.9: 76–100%) and intensity (0: absent, +0.5: very low, +1: low, +2: moderate, +3: severe). The histopathological score was determined as prevalence x intensity.

Statistical analysis

Statistical evaluations were performed by SPSS software, version 15.0. The one-sample Kolmogorov-Smirnov test was used to determine whether the data were distributed normally. One-way analysis of variance was used to compare TAS, TOS and immunoreactivity variables. The differences between the variables of groups were determined using the Tukey HSD multiple comparison test. The obtained results were exhibited as mean ± standard deviation (SD). The significance level was considered as $p < 0.05$.

Results

Biochemical results

In biochemical analysis, similar TAS (Table I) and TOS (Table II) levels were seen in the control,

VD, and CO groups. A statistically significantly lower TAS level and a statistically significantly higher TOS level were detected in the IMC group ($p < 0.05$) compared to the control group. Compared to the IMC group, the higher TAS level and lower TOS level in the IMC + VD group were insignificant ($p < 0.05$).

Histopathological results

The testes of the control group (Figure 1. A 1), VD group (Figure 1. A 2) and CO (Figure 1. A 3) group had normal histological structures. The testes of the IMC group (Figure 1. A 4) had degenerative changes in the seminiferous tubule epithelium and interstitial edema compared to the control group. In comparison to the IMC group, degenerative changes in the seminiferous tubule epithelium and interstitial edema were at a lower level in the IMC + VD group (Figure 1. A 5).

Immunohistochemical results

Immunohistochemically similar irisin immunoreactivity (Table III) was observed in the interstitial space of testis tissue in the control group (Figure 1. B 1), VD group (Figure 1. B 2) and CO group (Figure 1. B 3). Irisin immunoreactivity was significantly higher in the IMC group (Figure 1. B 4) compared to the control group ($p < 0.05$). Conversely, irisin immunoreactivity was insignificantly lower in the IMC + VD group (Figure 1. B 5) compared to the IMC group ($p > 0.05$).

Discussion

Organophosphate compounds are commonly known as pesticides and/or insecticides. Insecticides and pesticides induce toxic effects causing atrophy in the testis of rats [27]. ROS are effective in the pathogenesis of testicular damage induced by organophosphate compounds in rat models [28] and ROS is associated with the defective morphology of sperm cells [29].

IMC-like insecticides are reproductive toxic substances in male animals, causing disruption of tes-

Table I. TAS levels of testis tissues

Groups	TAS levels Mean ± SD	P-value*
Control	27.41 ± 1.4 ^b	
IMC	21.31 ± 1.31 ^a	< 0.001
IMC + VD	22.48 ± 1.64 ^a	
VD	27.57 ± 1.78 ^b	
CO	29.16 ± 1.48 ^b	

*One-way ANOVA. ^{a,b}Mean values of groups with different superscripts differ significantly from each other ($p < 0.05$).

Table II. TOS levels of testis tissues

Groups	TOS levels Mean ± SD	P-value*
Control	14.87 ± 1.304 ^b	
IMC	24.84 ± 1.64 ^a	< 0.001
IMC + VD	22.75 ± 2.39 ^a	
VD	16.02 ± 0.82 ^b	
CO	16.05 ± 1.15 ^b	

*One-way ANOVA. ^{a,b}Mean values of groups with different superscripts differ significantly from each other ($p < 0.05$).




Figure 1. A – Testis histopathology of the control, VD, CO, IMC, and IMC + VD groups. Scale bar is 100 μ m in the HE stained figures. Arrowheads indicate degeneration. **A 1** – Control group: normal histological view of testis. **A 2** – VD group: normal histological view of testis. **A 3** – CO group: normal histological view of testis. **A 4** – IMC group: severe degeneration, edema, and desquamation in testis. **A 5** – IMC + VD group: light degeneration and edema in testis. **B** – Irisin immunohistochemistry of the control, VD, CO, IMC, and IMC + VD groups, respectively. Scale bar is 200 μ m in the iris in stained figures. Arrows indicate immunopositivity. **B 1** – Control group: slight interstitial positivity. **B 2** – VD group: slight interstitial positivity. **B 3** – CO group: slight interstitial positivity. **B 4** – IMC group: severe interstitial positivity. **B 5** – IMC + VD group: slight interstitial positivity

ticular tissues, spermatogenic cells and decreased sperm motility [30] and may induce necrosis in the seminiferous tubules along with histopathological alterations [27]. In this study, significantly increased TOS and decreased TAS levels, as well as degeneration in the germinal epithelium of seminiferous tubules and interstitial edema, were observed only in the IMC group. Therefore, the negative effect of IMC on testicular tissue of male rats is thought to be related to oxidative stress.

In previous studies, decreased spermatocytes, reduced or lack of spermatogenesis, weak or arrested spermatogenesis in some seminiferous tubules and mild edema in the interstitial space, as well as increased apoptosis of germ cells and fragmentation of seminal DNA, have been reported in IMC-treated rats [31, 32]. In IMC-treated rabbits, it was observed that the seminiferous tubule lumen was empty and interstitial tissue widened probably as a result of the elimination of Leydig cells and in some tubules this space was compressed. Furthermore, vacuolization in the seminiferous tubules and pyknosis in the spermatocytes have been reported [33]. In the present study, histopathological observations confirmed that IMC caused testicular damage including degeneration in the germinal epithelium of seminiferous tubules and edema in the interstitial space in the testis tissue, similar to previous studies [31–34].

It has been reported that VD deficiency disrupted both testicular development and spermatogenesis in mice and that mature seminiferous tubules remain at a lower percentage [34]. VD-deficient rats showed degenerative changes in the germinal epithelium and significantly reduced Leydig cell and sperm numbers, while in the normal VD diet group, only body weight decreased [35]. In the current study, depending on the IMC administration, we observed histopathological damage, decreased TAS and increased TOS levels. In the histopathological examination of the IMC + VD-treated group, it was observed that VD ameliorated testis injuries by decreasing degenerative tubules and interstitial edema. However, VD showed an insignificant effect on the oxidative state. Taking into consideration these positive effects of VD on testis morphology, VD is thought to be useful in decreasing IMC-induced testicular tissue damage.

Depleted antioxidant levels were reported on IMC-treated male rats [31]. Decreased TAS and increased TOS levels were reported in the testis tissue of diabetic rats and VD administration was observed to improve these biochemical parameters by increasing TAS and decreasing TOS levels [36]. In the current study, while a significant decrease in TAS and increase in TOS levels were observed in the IMC group, an insignificant increase in TAS levels and decrease in TOS levels were detected in the IMC + VD group.

Table III. Histoscore of irisin immunoreactivity in testes

Groups	Irisin immunoreactivity Mean \pm SD	P-value*
Control	0.95 \pm 0.19 ^b	
IMC	1.67 \pm 0.54 ^a	< 0.001
IMC + VD	1.1 \pm 0.34 ^{ab}	
VD	0.95 \pm 0.32 ^b	
CO	0.90 \pm 0.24 ^b	

*One-way ANOVA. ^{a,b}Mean values of groups with different superscripts differ significantly from each other ($p < 0.05$).

Irisin is a peptide that has been studied with increasing interest recently. However, its importance is still not fully understood. The localization of this peptide has been determined in many tissues [37–39]. Gur *et al.* [40] determined that irisin is located in the cytoplasm of the cells of all examined tissues including testis. Furthermore, irisin immunoreactivity in seminiferous tubules, fetal spermatogenic cells, ductus epididymis in fetal human epididymis, Leydig cells in fetal and adult testis has been observed [23]. In our study, however, irisin immunoreactivity was observed only in the interstitial area. Additionally, irisin is thought to be associated with the antioxidant mechanism. It has been reported that irisin might protect cells from ROS-induced severe cellular oxidative damage via the activation of antioxidative mechanisms [41]. In another study, it was reported that hyper- and hypothyroidism are associated with up-regulation of serum irisin in male rats, probably as a reaction to oxidative damage or myopathy observed under both conditions [42]. Irisin levels have been observed to be increased in type 1 diabetes [39] and in hypothyroid patients [37]. In the current study, similar to previous studies [35, 39, 42], irisin immunoreactivity increased in the IMC-treated group compared to the control group. Increased immunoreactivity was insignificantly reduced in the IMC + VD group compared to the IMC group. This increase in irisin immunoreactivity may be a response to oxidative stress caused by IMC in testis tissue.

In conclusion, in this study, it was found that Vitamin D administered at a dose of VD 200 IU/day prevents testicular damage in rats given 20 mg/kg IMC daily for 8 weeks. ROS is thought to be a cause of testicular damage induced by IMC. Furthermore, VD was determined to have an effect in improving oxidative stress and especially testicular damage induced by IMC. Therefore, VD is thought to contribute to the improvement of testicular injury caused by IMC in male rats. The annual use of IMC and similar insecticides world-

wide reaches thousands of tonnes. Both field observations and experimental studies have shown that these compounds have significant negative effects on male fertility as well as many different side effects. In addition to their secondary effects on fertility, primary damage to the testis and spermatogenic cells can range from insufficient motility to atrophy. In this context, further studies on vitamin D, which demonstrated protective properties in terms of the testis and hence fertility, should be conducted and the specific dosage and administration method should be well studied. This is important because high doses of vitamin D may be harmful even for individuals with vitamin D deficiency.

Conflict of interest

The authors declare no conflict of interest.

References

- Grisham MB, Jourd'Heuil D, Wink DA. Nitric oxide. I. Physiological chemistry of nitric oxide and its metabolites: implications in inflammation. *Am J Physiol* 1999; 276: G315-21.
- Kanbur M, Liman BC, Eraslan G, Altinordulu S. Effects of cypermethrin, propetamphos, and combination involving cypermethrin and propetamphos on lipid peroxidation in mice. *Environ Toxicol* 2008; 23: 473-9.
- Muniz JF, McCauley L, Scherer J, et al. Biomarkers of oxidative stress and DNA damage in agricultural workers: a pilot study. *Toxicol Appl Pharmacol* 2008; 227: 97-107.
- Matsuda K, Buckingham SD, Kleier D, Rauh JJ, Grauso M, Sattelle DB. Neonicotinoids: insecticides acting on insect nicotinic acetylcholine receptors. *Trends Pharmacol Sci* 2001; 22: 573-80.
- Jeschke P, Nauen R, Schindler M, Elbert A. Overview of the status and global strategy for neonicotinoids. *J Agric Food Chem* 2011; 59: 2897-908.
- Bagri P, Kumar V, Sikka AK. Assessment of imidacloprid-induced mutagenic effects in somatic cells of Swiss albino male mice. *Drug Chem Toxicol* 2016; 39: 412-7.
- Duzguner V, Erdogan S. Acute oxidant and inflammatory effects of imidacloprid on the mammalian central nervous system and liver in rats. *Pestic Biochem Physiol* 2010; 97: 13-8.
- Kara M, Yumrutas O, Demir CF, et al. Insecticide imidacloprid influences cognitive functions and alters learning performance and related gene expression in a rat model. *Int J Exp Pathol* 2015; 96: 332-7.
- Mikolić A, Karačonji IB. Imidacloprid as reproductive toxicant and endocrine disruptor: investigations in laboratory animals. *Arh Hig Rada Toksikol* 2018; 69: 103-8.
- McCarty DE. Resolution of hypersomnia following identification and treatment of vitamin D deficiency. *J Clin Sleep Med* 2010; 6: 605-8.
- Caprio M, Mammi C, Rosano GM. Vitamin D: a novel player in endothelial function and dysfunction. *Arch Med Sci* 2012; 8: 4-5.
- Timar A, Saberi-Karimian M, Ghazizadeh H, et al. Evaluation of the serum prooxidant-antioxidant balance before and after vitamin D supplementation in adolescent Iranian girls. *Adv Med Sci* 2019; 64: 174-80.
- Makariou S, Liberopoulos E, Florentin M, et al. The relationship of Vitamin D with non-traditional risk factors for cardiovascular disease in subjects with metabolic syndrome. *Arch Med Sci* 2012; 8: 437-43.
- Ilinčić B, Stokić E, Stošić Z, et al. Vitamin D status and circulating biomarkers of endothelial dysfunction and inflammation in non-diabetic obese individuals: a pilot study. *Arch Med Sci* 2017; 13: 53-60.
- Dziedzic EA, Gašior JS, Pawłowski M, et al. Vitamin D level is associated with severity of coronary artery atherosclerosis and incidence of acute coronary syndromes in non-diabetic cardiac patients. *Arch Med Sci* 2019; 15: 359-68.
- Imga NN, Berker D, Can B, Guler S. The effects of three regimens of cholecalciferol (vitamin D3) supplementation on vitamin D deficiency in non-obese and obese females. *Arch Med Sci Atheroscler Dis* 2018; 3: e60-7.
- Costanzo PR, Knoblovits P. Vitamin D and male reproductive system. *Horm Mol Biol Clin Investig* 2016; 28: 151-9.
- de Angelis C, Galdiero M, Pivonello C, et al. The role of Vitamin D in male fertility: a focus on the testis. *Rev Endocr Metab Disord* 2017; 18: 285-305.
- Ozkaya F, Demirel A. Vitamin D deficiency in infertile patients. *Arch Esp Urol* 2018; 71: 850-5.
- Merino O, Sánchez R, Gregorio MB, Sampaio F, Risopatrón J. Effect of high-fat and vitamin D deficient diet on rat sperm quality and fertility. *Theriogenology* 2019; 125: 6-11.
- Boström P, Wu J, Jedrychowski MP, et al. A PGC1- α -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature* 2012; 481: 463-8.
- Gouni-Berthold I, Berthold HK, Huh JY, et al. Effects of lipid-lowering drugs on irisin in human subjects in vivo and in human skeletal muscle cells ex vivo. *PLoS One* 2013; 8: e72858.
- Aydin S, Kuloglu T, Aydin S, et al. A comprehensive immunohistochemical examination of the distribution of the fat-burning protein irisin in biological tissues. *Peptides* 2014; 61: 130-6.
- Askari H, Rajani SF, Poorebrahim M, Haghi-Aminjan H, Raeis-Abdollahi E, Abdollahi M. A glance at the therapeutic potential of irisin against diseases involving inflammation, oxidative stress, and apoptosis: an introductory review. *Pharmacol Res* 2018; 129: 44-55.
- Erel O. A novel automated direct measurement method for total antioxidant capacity using a new generation, more stable ABTS radical cation. *Clin Biochem* 2004; 37: 277-85.
- Erel O. A new automated colorimetric method for measuring total oxidant status. *Clin Biochem* 2005; 38: 1103-11.
- Bustos-Obregón E, González-Hormazabal P. Effect of a single dose of malathion on spermatogenesis in mice. *Asian J Androl* 2003; 5: 105-7.
- Gultekin F, Ozturk M, Akdogan M. The effect of organophosphate insecticide chlorpyrifos-ethyl on lipid peroxidation and antioxidant enzymes (in vitro). *Arch Toxicol* 2000; 74: 533-8.
- Kessopoulou E, Tomlinson MJ, Barratt CL, Bolton AE, Cooke ID. Origin of reactive oxygen species in human semen: spermatozoa or leucocytes? *J Reprod Fertil* 1992; 94: 463-70.
- Karabay NU, Oguz MG. Cytogenetic and genotoxic effects of the insecticides, imidacloprid and methamidophos. *Genet Mol Res GMR* 2005; 4: 653-62.
- Bal R, Turk G, Tuzcu M, et al. Assessment of imidacloprid toxicity on reproductive organ system of adult male rats. *J Environ Sci Health B* 2012; 47: 434-44.

32. Lonare M, Kumar M, Raut S, et al. Evaluation of ameliorative effect of curcumin on imidacloprid-induced male reproductive toxicity in wistar rats. *Environ Toxicol* 2016; 31: 1250-63.
33. Memon SA, Memon N, Mal B, Shaikh SA, Shah MA. Histopathological changes in the gonads of Male rabbits (*Oryctolagus cuniculus*) on exposure to imidacloprid insecticide. *J Entomol Zool Stud* 2014; 2: 159-63.
34. Fu L, Chen YH, Xu S, et al. Vitamin D deficiency impairs testicular development and spermatogenesis in mice. *Reprod Toxicol* 2017; 73: 241-9.
35. Sood S, Marya RK, Reghunandan R, Singh GP, Jaswal TS, Gopinathan K. Effect of Vitamin D deficiency on testicular function in the rat. *Ann Nutr Metab* 1992; 36: 203-8.
36. Yalcin MH. Investigation of effects of Vitamin D on irisin and apoptosis in experimental diabetic rat's testicular tissue. PhD thesis. Firat University Health Science Institute. Elazig, 2018. [in Turkish with English abstract].
37. Ates I, Altay M, Topcuoglu C, Yilmaz FM. Circulating levels of irisin is elevated in hypothyroidism, a case-control study. *Arch Endocrinol Metab* 2016; 60: 95-100.
38. Erdogan MA, Yalcin A. Protective effects of benfotiamine on irisin activity in methotrexate-induced liver injury in rats. *Arch Med Sci* 2018; <https://doi.org/10.5114/aoms.2018.80002>.
39. Faienza MF, Brunetti G, Sanesi L, et al. High irisin levels are associated with better glycemic control and bone health in children with type 1 diabetes. *Diabetes Res Clin Pract* 2018; 141: 10-7.
40. Gur FM, Timurkaan S, Yalcin MH, Girgin A, Gencer Tarakci B. Immunohistochemical localization of irisin in mole rats (*Spalax leucodon*). *Biotech Histochem Comm* 2017; 92: 245-51.
41. Mazur-Bialy AI, Kozłowska K, Pochec E, Bilski J, Brzozowski T. Myokine irisin-induced protection against oxidative stress in vitro. Involvement of heme oxygenase-1 and antioxidizing enzymes superoxide dismutase-2 and glutathione peroxidase. *J Physiol Pharmacol* 2018; 69: 117-25.
42. Samy DM, Ismail CA, Nassra RA. Circulating irisin concentrations in rat models of thyroid dysfunction – effect of exercise. *Metabolism* 2015; 64: 804-13.