Identification of Urothelial Cancer-Associated 1 (UCA1) as a diagnostic biomarker of pre-eclampsia via regulating microRNA-16 and its downstream signaling pathway

Type

Research paper

Keywords

miR-16, UCA1, Pre-eclampsia (PE), PAPP-A

Abstract

Introduction

In this study, we studied the clinical value of using urothelial cancer-associated 1 (UCA1) and microRNA-16 (miR-16) as biomarkers for the diagnosis of pre-eclampsia (PE). Also, we compared the diagnostic values of miR-16, UCA1 and pregnancy-associated plasma protein-A (PAPP-A) in PE. Furthermore, we investigated the interaction between miR-16 and UCA1/PAPP-A.

Material and methods

128 PE patients and 172 healthy pregnant women were enrolled in this study. Receiver operating characteristic (ROC) analysis was carried out to predict the diagnostic values of UCA1, miR-16 and PAPP-A in PE. Enzyme-linked immunosorbent assay (ELISA), real-time polymerase chain reaction (PCR), Western blot analysis, immunohistochemistry (IHC) assay, computational analysis, and luciferase assay were conducted to measure the differential expression of UCA1, miR-16, and PAPP-A while establishing a signaling pathway of UCA1/miR-16/PAPP-A.

Results

Compared with miR-16 and PAPP-A, UCA1 exhibited a better value in the diagnosis of PE. The expression of PAPP-A and UCA1 was down-regulated while the expression of miR-16 was up-regulated in patients with PE, especially in patients with HELLP pregnancies. Moreover, UCA1 was identified as a sponge of miR-16, while PAPP-A mRNA was identified as a virtual target gene of miR-16. Finally, a negative regulatory relationship was observed between the expression of miR-16 and UCA1 or PAPP-A, while the expression of UCA1 and PAPP-A were positively related.

Conclusions

Taken together, UCA1 could be used as a more valuable biomarker in the diagnosis of PE. Meanwhile, the reduced expression of UCA1 could exert a positive effect by reducing the expression of PAPP-A in the pathogenesis of PE.

1 Identification of Urothelial Cancer-Associated 1 (UCA1) as a diagnostic biomarker of 2 pre-eclampsia via regulating microRNA-16 and its downstream signaling pathway

3 Ye-hong Xv¹, Wei Gao², Liao-jv Wang^{2*}

4 1. Department of Obstetrics, Northwest Women's and Children's Hospital, Xi'an,
5 Shaanxi, China 710061

6 2. Department of Obstetrics, The First People's Hospital of Xianyang City, Xianyang,
7 Shaanxi, China 712000

8 *Corresponding author: Liao-jv Wang

9 Institution: Department of Obstetrics, The First People's Hospital of Xianyang City,
10 Xianyang, Shaanxi, China 712000

11 Address: No.10 Biyuan Rd, Xianyang, Shaanxi, China 712000

12 Email: PECXYH@yeah.net

13 Abstract

14 **Background:** In this study, we studied the clinical value of using urothelial cancer-
15 associated 1 (UCA1) and microRNA-16 (miR-16) as biomarkers for the diagnosis of pre-
16 eclampsia (PE). Also, we compared the diagnostic values of miR-16, UCA1 and pregnancy-
17 associated plasma protein-A (PAPP-A) in PE. Furthermore, we investigated the interaction
18 between miR-16 and UCA1/PAPP-A. **Method:** 128 PE patients and 172 healthy pregnant
19 women were enrolled in this study. Receiver operating characteristic (ROC) analysis was
20 carried out to predict the diagnostic values of UCA1, miR-16 and PAPP-A in PE. Enzyme-
21 linked immunosorbent assay (ELISA), real-time polymerase chain reaction (PCR), Western
22 blot analysis, immunohistochemistry (IHC) assay, computational analysis, and luciferase
23 assay were conducted to measure the differential expression of UCA1, miR-16, and PAPP-
24 A while establishing a signaling pathway of UCA1/miR-16/PAPP-A. **Results:** Compared
25 with miR-16 and PAPP-A, UCA1 exhibited a better value in the diagnosis of PE. The
26 expression of PAPP-A and UCA1 was down-regulated while the expression of miR-16 was
27 up-regulated in patients with PE, especially in patients with HELLP pregnancies. Moreover,
28 UCA1 was identified as a sponge of miR-16, while PAPP-A mRNA was identified as a virtual

target gene of miR-16. Finally, a negative regulatory relationship was observed between the expression of miR-16 and UCA1 or PAPP-A, while the expression of UCA1 and PAPP-A were positively related. **Conclusion:** Taken together, UCA1 could be used as a more valuable biomarker in the diagnosis of PE. Meanwhile, the reduced expression of UCA1 could exert a positive effect by reducing the expression of PAPP-A in the pathogenesis of PE.

Running title: MiR-16, UCA1, and PAPP-A value in PE diagnosis

Keywords: Pre-eclampsia (PE), UCA1, miR-16, PAPPA

Introduction

As a complication of pregnancy, pre-eclampsia (PE) is featured by proteinuria and increased blood pressure [1]. In Western countries, the incidence of PE is about 5%, although the incidence of PE is much higher in developing countries [2]. Worldwide, PE is attributed to over one hundred thousand maternal deaths every year and is considered as a key reason for premature delivery, i.e., children born with <1500 g of birth weight [1, 3, 4]. Moreover, PE has been shown to cause cardiovascular morbidity in pregnant women [5].

Initially isolated from the serum of pregnant women, pregnancy-associated plasma protein-A (PAPP-A) contains a unique metalloproteinase zinc-binding motif in its amino acid structure, although the role of this motif remains unclear. PAPP-A is increasingly used for the prenatal screening of trisomy 21, either alone or in combination with ultrasound measurement of nuchal translucency (NT) [6, 7], while recently, a study showed that PAPP-A is associated with a proteolytic activity [8-11]. In fact, an abnormally low level of PAPP-A has been used as a biomarker to predict the tendency to develop small-for-gestational-age (SGA) [12]. Also, it was shown that the ultrasound examination of the uterine artery can provide useful information to predict the onset of PE even in the first trimester [13]. Currently, it is believed that an abnormally low level of PAPP-A plays a role as essential as that of abnormal ultrasound results in the prediction of fetuses with SGA, although the role of a low level of PAPP-A in the prediction of PE remains controversial

[12]. These discrepancies may be due to the different molecular mechanisms implicated in the development of SGA and PE, such that an abnormal ultrasound result may indicate the invasion of trophoblasts in maternal spiral arteries [14], whereas a low value of PAPP-A may indicate that the placenta can no longer support the further growth of the fetus.

The expression of transcriptome is regulated by a complex network involving the interaction among a wide range of molecules, including messenger RNA (mRNA), microRNA (miRNA), and long non-coding RNA (lncRNA) [15]. Among them, miRNA plays an essential role to maintain the stability of transcriptome and to regulate the homeostasis in the body. Recently, the molecular interaction between lncRNA/miRNA and mRNAs has gained special interests because it has been revealed that both miRNAs, a type of small non-coding RNAs (ncRNAs) with a length of about 22 nucleotides, and lncRNAs, a type of ncRNAs with a length of over 200 nucleotides, are involved in the translation regulation of their target mRNAs [16, 17]. At the same time, mRNAs were also shown to regulate the expression of ncRNAs in a certain way [18, 19].

In a previous study, it was found that increased expression of urothelial cancer-associated 1 (UCA1) was positively associated with the increase in imatinib (IM) resistance [20]. Further investigation showed that UCA1 competitively binds to miR-16 and suppressing the function of miR-16. In another study, it was demonstrated that UCA1 functions as competing endogenous (ceRNA) of miR-16 [20]. In bladder cancer cells, UCA1 could also function as an endogenous sponge of miR-16 to regulate GLS2 expression and glutamine metabolism [21].

It has been shown that PAPP-A could function as a biomarker in the diagnosis of PE [22]. Meanwhile, using an online miRNA database, we found that miR-16 might act as a potential regulator of PAPP-A. Furthermore, UCA1 has been identified as a competing endogenous RNA (ceRNA) of miR-16 [21]. In this study, we studied the value of using UCA1 and miR-16 as biomarkers in the diagnosis of PE. In addition, we compared the diagnostic values of miR-16, UCA1 and PAPP-A in PE.

Materials and Methods

Human subjects and sample collection

In this study, PE patients (N=128) and healthy pregnant women (N=172) were enrolled in the case group and the control group, respectively. Subsequently, according to the type of PE, the case group was further divided into a MILD group (patients with mild PE, N=88), a SEVERE group (patients with severe PE, N=32) and a HELLP group (patients with HELLP pregnancies, which was a complication of pregnancy characterized by hemolysis, elevated liver enzymes, and a low platelet count. N=8). Demographic and clinical characteristics of all subjects were collected and analyzed and demonstrated in Table 1. Moreover, peripheral blood samples and placenta samples were collected from all subjects for subsequent receiver operating characteristic (ROC) analyses. All patients have signed the written informed consent before their participation in this study. In addition, this study was approved by the Ethics Committee of our institute and was carried out in accordance with the Helsinki declaration.

RNA isolation and real-time PCR

A miRNeasy Mini Kit (Qiagen, Hilden, Germany) was used to extract the total RNA from tissue and cell samples. Isolated RNA was reversely transcribed into complementary DNA (cDNA) templates using a reverse transcription kit (Thermo Fisher Scientific, Waltham, MA). The expression of UCA1, miR-16, and PAPP-A mRNA was measured using quantitative real-time PCR and an SYBR Premix Ex Taq II reagent kit (Takara, Tokyo, Japan) following the instruction of the kit. The reaction system of quantitative real-time PCR contained a volume of 20 μ l, including 10.0 μ l of One Step SYBR RT-PCR Buffer III, 0.4 μ l of TaKaRa Ex Taq HS, 0.4 μ l of Prime Script RT Enzyme Mix II, 0.3 μ l of PCR forward primer, 0.3 μ l of PCR reverse primer, 0.4 μ l of ROX Reference Dye (50 $\times$), 2 μ l of total RNA, and 6.2 μ l of RNase Free ddH₂O. An ABI7500 quantitative PCR instrument (ABI, Foster City, CA) was used to carry out the quantitative real-time PCR. The reaction conditions were set as follows: pre-denaturation at 95 °C for 10 s, followed by 40 cycles of denaturation at 95 °C for 5 s and annealing at 60 °C for 25 s, and a final extension step. U6 was used as the internal reference to quantify the relative expression of UCA1, miR-16, and PAPP-A mRNA using the $2^{-\Delta\Delta Ct}$ method. Each experiment was repeated three times.

Cell culture and transfection

Trophoblast and Hela cells were acquired from ATCC (American Type Culture Collection, Manassas, VA) and maintained in an environment of saturated humidity at 37 °C and under 5% CO₂. DMEM (Gibco, Thermo Fisher Scientific, Waltham, MA) containing 10% fetal bovine serum, 100 unit/mL streptomycin, and 100 mg/mL penicillin was used as the culture medium. For transfection experiments, trophoblast and Hela cells were seeded into 96-well plates at a density of 10⁴ cells per well and incubated overnight. On the next day, the cells were transfected with pcDNA-UCA1, pcDNA-anti-miR-16 using Lipofectamine 2000 (Invitrogen, Carlsbad, CA) following the instruction of the manufacturer. The transfected cells were harvested at 48 h post-transfection for subsequent analyses. Each experiment was repeated three times.

Vector construction, mutagenesis, and luciferase assay

The putative target genes of miR-16 were determined using online bioinformatics tools. The results identified the presence of miR-16 binding sites in the 3' UTR of PAPP-A mRNA and the promoter of UCA1, respectively. Subsequently, the 3' UTR of PAPP-A mRNA and the promoter of UCA1 containing the miR-16 binding sites were amplified by PCR and cloned into pcDNA vectors (Promega, Madison, WI) to generate wild type vectors for the 3' UTR of PAPP-A mRNA and the promoter of UCA1, respectively. At the same time, site-directed mutagenesis was also carried out at the miR-16 binding sites in the 3' UTR of PAPP-A mRNA and the promoter of UCA1, respectively. Subsequently, the mutant sequences were also amplified by PCR and cloned into pcDNA vectors to generate mutant vectors for the 3' UTR of PAPP-A mRNA and the promoter of UCA1, respectively. In the next step, Trophoblast and Hela cells were co-transfected with UCA1 and miR-16, or miR-16 and PAPP-A mRNA, using Lipofectamine 2000. At 48 h after transfection, the transfected cells were collected and the luciferase reporter activity in transfected cells was detected by a Dual Luciferase Reporter Assay System (Promega, Madison, WI). The Renilla internal control was used in the assay to normalize the relative firefly luciferase activity of UCA1 and PAPP-A mRNA, respectively. Each experiment was repeated three times.

Western blot analysis

Collected tissue and cell samples were lysed using a radioimmunoprecipitation assay (RIPA) buffer (Invitrogen, Thermo Fisher Scientific, Waltham, MA) supplemented with 1% sodium deoxycholate, 1% NP-40, 0.1% SDS, 150 mM NaCl and 50 mM Tris-HCl (pH 8.8). The lysis treatment was performed at 4°C for 30 min following the guideline provided by the supplier. In the next step, the cell lysate was centrifuged at 4°C and 15,000 rpm for 10 min. Subsequently, the bicinchoninic acid method was used to measure the concentration of protein in the supernatant. Thereafter, 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was used to separate the protein in each sample, and the separated proteins were then blotted onto a polyvinylidene difluoride (PVDF) membrane (Immobilon-P; Millipore, Billerica, MA) for 60 min at 180 mA using a transfer buffer containing 0.2 M glycine, 20% methanol, and 25 mM Tris (Promega, Madison, WI) following a standard protocol. Subsequently, the membrane was blocked in 5% skim milk (Merck, Darmstadt, Germany) and then incubated with anti-PAPP-A (1:1000 dilution, Abcam, Cambridge, MA) and anti- β -actin primary antibodies (1:10,000 dilution, Abcam, Cambridge, MA) overnight at 4°C, followed by another 60 min of incubation with horseradish peroxidase (HRP)-conjugated goat anti-rabbit immunoglobulin G secondary antibodies (1:10,000 dilution, Abcam, Cambridge, MA). In the next step, the chemiluminescent signals of bound antibodies were detected using an Electrochemiluminescence Plus Western Blotting Detection System (GE Healthcare Bio-Sciences, Pittsburgh, PA) in conjunction with a high-performance chemiluminescence film (GE Healthcare Bio-Sciences, Pittsburgh, PA) in accordance with the manufacturer's recommendation. The relative expression of PAPP-A protein in each sample was normalized to the protein expression of the internal control β -actin. Each experiment was repeated three times.

Immunohistochemistry

Sample sections were dewaxed twice in xylene (10 min each time) and dehydrated in graded ethanol (100%, 95%, 80%; and 70%, 2 min each time). Subsequently, the sections were placed on a shaker and washed twice in distilled water (5 min each time). In the next step, the sections were incubated in 3% H₂O₂ for 10 min and washed by distilled water,

followed by antigen repair at high pressure for 90 s. After being cooled down to room temperature, the sections were washed with phosphate buffer saline (PBS), blocked with 5% bovine serum albumin (BSA) for 30 min at 37°C, and then incubated overnight at 4°C with anti-PAPP-A primary antibodies (1 : 100, Abcam, Cambridge, MA). After PBS washing, the sections were further incubated with biotinylated secondary antibodies (1: 100, Abcam, Cambridge, MA) at 37°C for 30 min. Subsequently, after being fully washed, the sections were incubated with horseradish peroxidase (HRP)-conjugated working fluid, stained by 3,3-diaminobenzidine (DAB, chromogenic agent), and counterstained for 5 min with hematoxylin. During this experiment, PBS was used as the negative control. After staining, the cells showing positive expression of PAPP-A protein were analyzed under a microscope. In brief, five fields of vision were chosen randomly from each section under a high power lens to detect the localization of positive PAPP-A expression. Each experiment was repeated three times.

ELISA

The level of PAPP-A protein expression in collected samples was measured using an ELISA kit (Thermo Fisher Scientific, Waltham, MA) following the instruction of the kit. Each experiment was repeated three times.

Statistical analysis

All statistical analysis was carried out using SPSS 19.0 software (SPSS, Chicago, IL). All data were shown as mean $\pm$ standard deviations (SD). The differences between two different groups were compared using Student's *t*-tests, while the differences between multiple groups were compared using one-way analysis of variance (ANOVA). The ROC curves were plotted using NCSS Statistical Software (NCSS, Kaysville, UT) to calculate the area under curve (AUC) values of different markers. At least three independent tests were performed for each experiment. A P value of < 0.05 was considered statistically significant.

Results

UCA1 showed a better value than miR-16 and PAPP-A in the diagnosis of PE

In this study, demographic and clinical characteristics of all subjects were collected and analyzed, as shown in Table 1. And there were no significant differences between

different groups of PE patients as well as the case group with respect to the above indexes. Moreover, as shown in Fig.1, upon the calculation of the diagnostic values of UCA1, miR-16 and PAPP-A in the diagnosis of PE, the AUC of miR-16 and PAPP-A was very similar, while the AUC of UCA1 was evidently higher, suggesting that UCA1 can be used as a better marker in the diagnosis of PE.

UCA1/PAPP-A expression was inhibited in PE cases along with an enhanced miR-16 expression

As shown in Fig.2A, PAPP-A expression was similarly down-regulated in the MILD group and SEVERE group compared with that in the control group, which showed the highest PAPP-A expression. On the contrary, the HELLP group showed the lowest PAPP-A expression. Besides, the relative expression of UCA1 in the plasma (Fig.2C) showed the same trend as that of log10 MoM PAPP-A. However, the relative expression of miR-16 in the plasma (Fig.2B) showed the opposite trend: The expression of miR-16 was the highest in the HELLP group and the lowest in the control group. Furthermore, the expression of PAPP-A (Fig.3A), miR-16 (Fig.3B) and UCA1 (Fig.3C) in placenta samples collected from different groups also showed a similar trend as that in the peripheral blood samples. Finally, the trend in PAPP-A expression among different groups was also validated using IHC assays (Fig.3D). Therefore, PE patients showed inhibited miR-16 expression and increased UCA1/PAPP-A expression, while the patients with HELLP pregnancies showed the lowest UCA1/PAPP-A expression and the highest miR-16 expression.

UCA1 and PAPP-A mRNA functioned as a sponge and a target of miR-16, respectively

Based on a computational analysis carried out using an online target prediction tool (www.targetscan.org), a binding sequence of miR-16 was identified in UCA1 (Fig.4A). Subsequently, a luciferase assay was performed to evaluate the interaction between miR-16 and UCA1, and the results demonstrated evidently reduced luciferase activity in Trophoblast (Fig.4B) and Hela (Fig.4C) cells co-transfected with miR-16 and wild-type UCA1, suggesting that UCA1 acts as a sponge of miR-16. Additionally, a 'seed sequence' of miR-16 was identified in the 3'UTR of PAPP-A mRNA (Fig.4D), and the luciferase activity

of Trophoblast (Fig.4E) and Hela (Fig.4F) cells was significantly reduced upon the co-transfection with miR-16 and wild-type PAPP-A 3'UTR, indicating that PAPP-A mRNA acts as a virtual target gene of miR-16.

The establishment of the UCA1/miR-16/PAPP-A signaling pathway

To further investigate the regulatory relationships among UCA1, miR-16, and PAPP-A, Trophoblast cells were transfected with plasmids carrying UCA1 or anti-miR-16. As shown in Fig.5, the relative expression of miR-16 (Fig.5A) was significantly suppressed in the presence of UCA1 or anti-miR-16, along with significantly increased expression of PAPP-A mRNA. Also, the results of Western blot analysis showed significantly increased expression of PAPP-A protein (Fig.5C) in the UCA1 and anti-miR-16 groups, verifying the role of UCA1 as a negative regulator of miR-16 expression. Unlike miR-16, a positive relationship was confirmed between UCA1 and PAPP-A. Similar results were also obtained from Hela cells (Fig.6), thus establishing a UCA1/miR-16/PAPP-A signaling pathway. In summary, it can be concluded that when the expression of UCA1 is reduced, the expression of PAPP-A is accordingly suppressed, thus exhibiting positive effects in the pathogenesis of PE.

Discussion

As a major contributor to neonatal and maternal mortality and morbidity, PE is known for its role as a trigger of maternal proteinuria, hypertension, imbalance of angiogenic factors, and placental dysfunction [23]. During PE, the decrease in uteroplacental perfusion may cause ischemia, which in turn promotes the release of soluble factors from the ischemic placenta and subsequently causes hypertension and endothelial dysfunction [24]. The physiological alterations observed in PE also promote the synthesis of proinflammatory cytokines, including IL-6 and tumor necrosis factor-alpha (TNF- α), from the placenta [25]. Recently, many studies showed that the dysregulation in the production of placental growth factor (PlGF), vascular endothelial growth factor (VEGF), and soluble fms-like tyrosine kinase I (sFlt-1) plays an essential role in the pathology of PE [26, 27].

UCA1 was first discovered in human bladder carcinoma [20]. Previous studies have reported that UCA1 can promote the development of drug resistance, metastasis, and proliferation in many types of cancer cells, including the cells of breast cancer and bladder cancer [28-30]. UCA1 was recently found to function as a miRNA sponge. For example, UCA1 can activate mTOR to regulate the expression of hexokinase 2 (HK2) and glycolysis by activating STAT3 and by inhibiting the expression of miR-143 [31]. UCA1 also increases the invasion and migration ability of epithelial ovarian cancer cells by directly binding to miR-485-5p [32]. In this study, the AUC of miR-16 and PAPP-A was evidently lower than that of UCA1, suggesting that UCA1 is associated with a better value in the diagnosis of PE.

In both plasma and placenta samples collected from the subjects of this study, the expression of PAPP-A and UCA1 was similarly down-regulated in the MILD and SEVERE groups, while the patients in the HELLP group showed the lowest expression of PAPP-A and UCA1. Meanwhile, the relative expression of miR-16 was the highest in the HELLP group and the lowest in the control group, verifying the role of UCA1, miR-16 and PAPP-A dysregulation in the pathogenesis of PE, especially in the pathogenesis of HELLP pregnancies.

In addition, it was shown that miR-16 acts as a tumor suppressor by regulating the apoptosis and proliferation of many types of cancer cells, such as bladder cancer cells [33]. It was postulated that UCA1 can bind to miR-16 to impair its functions in bladder cancer. Past results also indicated that UCA1 exerts a dose-dependent sponge effect on miR-16. Besides, it was shown using luciferase assays that miR-16 can directly bind to the 3'UTR of GLS2 mRNA, suggesting that UCA1 acts as a tumor suppressor and sponge of miR-16 to block its role in promoting the tumorigenesis of bladder cancer. In this study, miR-16 binding sites were identified on UCA1 and the 3'UTR of PAPP-A mRNA, respectively. Also, the luciferase activity of cells was evidently reduced when the cells were co-transfected with miR-16 and wild-type UCA1/PAPP-A mRNA. These results suggested that UCA1 functions as a sponge of miR-16, while PAPP-A mRNA acts as a virtual target gene of miR-16. In humans, miR-16 targets WNT3A, BCL2 and Cyclin D1 (CCND1),

all of which can enhance cell cycle arrest at the G1/S checkpoint and suppress the invasion, proliferation, and survival of cancer cells [34]. In fact, it has been shown that, upon DNA damage, the expression of miR-16 is quickly increased before returning to a normal value. Altered expression of miR-16 can dramatically change the activity of Wip1. For example, the silencing of miR-16 expression promoted the expression of Wip1, while the over-expression of miR-16 significantly inhibited the expression of Wip1 upon DNA damage. Some of these miR-16 potential targets like BCL2, Wip1, and CCND1 have been observed to be associated or involved in the pathogenesis of PE [35, 36, 37]. An analysis carried out on the miRNA/miRNA GO network produced most connections for miR-16. Also, the over-expression of miR-16 suppressed the migration and proliferation of decidua derived mesenchymal stem cells (dMSCs) while inducing cell-cycle arrest by binding to CCNE1. Furthermore, the over-expression of miR-16 also decreased the angiogenesis potential of HUVECs and blocked the migration of trophoblast cells, and such effect of miR-16 may be due to its activity in inhibiting the production of VEGF-A by dMSCs. Interestingly, the expression of VEGF-A and CCNE1 is negatively related to the expression of miR-16 in dMSCs, suggesting that the change in miR-16 expression in dMSCs may be implicated in the pathogenesis of PE. In this study, compared with that in cells transfected with the negative control, the relative expression of miR-16 was suppressed in cells transfected with UCA1 or anti-miR-16. In addition, the expression of PAPP-A mRNA/protein was elevated in cells transfected with UCA1 or anti-miR-16. Therefore, in the signaling pathway of UCA1/miR-16/PAPP-A, UCA1 acted as a negative regulator of miR-16 expression and an enhancer of PAPP-A mRNA/protein expression.

As a protein with high concentrations in the circulation of pregnant women, PAPP-A was initially discovered in 1974, although its biological role remained unclear for several decades. Nevertheless, PAPP-A1 has been used clinically as a biomarker in the screening of Down's syndrome [8, 38]. In the 1990s, it was reported that PAPP-A is associated with a protease activity against insulin-like growth factor binding protein-4 (IGFBP-4), and a unique feature of this protease activity is that the presence of IGF-II or IGF-I is required to cleave IGFBP-4 [39-41]. As a protein complex, PAPP-A plays an important role and its

expression in PE patients is significantly decreased [22]. Therefore, PAPP-A may be used as a potential biomarker for the early diagnosis of PE. Besides, when the serum level of PAPP-A is used in conjunction with other serum biomarkers or imaging examinations, including the ultrasound examination of the uterine artery in the first trimester [6, 7], it may provide more valuable information for the early diagnosis of PE. Moreover, Canini et al have shown that PAPP-A can be used as a more useful marker in the prediction of fetal growth restriction than in the prediction of PE [42]. Spencer et al also showed that a low level of serum PAPP-A in pregnant women is associated with an elevated risk of SGA, regardless of the karyotype of the infants [43].

Conclusion

In summary, the findings of this study demonstrated that, compared with miR-16 and PAPP-A, UCA1 can be used as a biomarker of higher diagnostic value in the diagnosis of PE. Meanwhile, miR-16 was validated to act as a potential regulator of PAPP-A expression while UCA1 was demonstrated to sponge the expression of miR-16. Therefore, the signaling pathway of UCA1/miR-16/PAPP-A was established, in which the expression of PAPP-A was positively regulated by UCA1. Consequently, the reduced expression of UCA1 could exert a positive effect by reducing the expression of PAPP-A in the pathogenesis of PE.

Conflict of interest

None

Figure legends

Table 1

Demographic and clinical characteristics of subjects without PE (N=172), subjects with mild PE (N=88), subjects with severe PE (N=32) and subjects with HELLP pregnancies (N=8).

Figure 1.

Diagnostic values of UCA1, miR-16 and PAPP-A were determined by calculating the AUC of each index.

Fig.2

- A. Log10 MoM PAPP-A in the plasma of MILD, SEVERE, HELLP and control groups;
- B. Relative expression of miR-16 in the plasma of MILD, SEVERE, HELLP and control groups;
- C. Relative expression of UCA1 in the plasma of MILD, SEVERE, HELLP and control groups.

Fig.3

- A. Log10 MoM PAPP-A in the placenta samples of MILD, SEVERE, HELLP and control groups;
- B. Relative expression of miR-16 in the placenta samples of MILD, SEVERE, HELLP and control groups;
- C. Relative expression of UCA1 in the placenta samples of MILD, SEVERE, HELLP and control groups;
- D. Expression of PAPP-A protein in the placenta samples of MILD, SEVERE, HELLP and control groups were detected using the IHC assay.

Fig.4

- A. Computational analysis on the sequences of miR-16 and UCA1;
- B. Luciferase assay of Trophoblast cells co-transfected with wild-type/mutant UCA1 and miR-16/negative controls;
- C. Luciferase assay of Hela cells co-transfected with wild-type/mutant UCA1 and miR-16/negative controls;
- D. Computational analysis on the sequences of miR-16 and PAPP-A mRNA;
- E. Luciferase assay of Trophoblast cells co-transfected with wild-type/mutant PAPP-A mRNA and miR-16/negative controls;

364 F. Luciferase assay of Hela cells co-transfected with wild-type/mutant PAPP-A mRNA and
365 miR-16/negative controls.

366 **Fig.5**

367 A. Relative expression of miR-16 in Trophoblast cells transfected with negative controls,
368 UCA1, or anti-miR-16;

369 B. Relative expression of PAPP-A mRNA in Trophoblast cells transfected with negative
370 controls, UCA1, or anti-miR-16;

371 C. Relative expression of PAPP-A protein in Trophoblast cells transfected with negative
372 controls, UCA1, or anti-miR-16;

373 D. The relative density of the PAPP-A protein band in Trophoblast cells transfected with
374 negative controls, UCA1, or anti-miR-16.

375 **Fig.6**

376 A. Relative expression of miR-16 in Hela cells transfected with negative controls, UCA1, or
377 anti-miR-16;

378 B. Relative expression of PAPP-A mRNA in Hela cells transfected with negative controls,
379 UCA1, or anti-miR-16;

380 C. Relative expression of PAPP-A protein in Hela cells transfected with negative controls,
381 UCA1, or anti-miR-16;

382 D. The relative density of the PAPP-A protein band in Hela cells transfected with negative
383 controls, UCA1, or anti-miR-16.

384 **References**

- 385 1 Sibai B, Dekker G, Kupferminc M: Pre-eclampsia. Lancet 2005;365:785-799.
386 2 Moodley J: Maternal deaths due to hypertensive disorders in pregnancy. Best
387 Pract Res Clin Obstet Gynaecol 2008;22:559-567.

- 388 3 Wright A, Wright D, Syngelaki A, Georgantzi A, Nicolaides KH: Two-stage screening
389 for preterm preeclampsia at 11-13 weeks' gestation. *Am J Obstet Gynecol*
390 2018;10.1016/j.ajog.2018.10.092
- 391 4 Shennan AH: Recent developments in obstetrics. *BMJ* 2003;327:604-608.
- 392 5 Bellamy L, Casas JP, Hingorani AD, Williams DJ: Pre-eclampsia and risk of
393 cardiovascular disease and cancer in later life: systematic review and meta-
394 analysis. *BMJ* 2007;335:974.
- 395 6 Spencer K, et al.: First-Trimester Markers of Aneuploidy and the Prediction of Small-
396 for-Gestational Age Fetuses. *Obstetrical & Gynecological Survey* 63.4(2008):215-
397 217.
- 398 7 Spencer, Kevin: Screening for trisomy 21 in twin pregnancies in the first trimester
399 using free β -hCG and PAPP-A, combined with fetal nuchal translucency thickness.
400 *Prenat Diagn* 20.2(2015):91-95.
- 401 8 Lin TM, Galbert SP, Kiefer D, Spellacy WN, Gall S: Characterization of four human
402 pregnancy-associated plasma proteins. *Am J Obstet Gynecol* 1974;118:223-236.
- 403 9 Kristensen T, Oxvig C, Sand O, Moller NP, Sottrup-Jensen L: Amino acid sequence
404 of human pregnancy-associated plasma protein-A derived from cloned cDNA.
405 *Biochemistry* 1994;33:1592-1598.
- 406 10 Stocker W, Grams F, Baumann U, Reinemer P, Gomis-Ruth FX, McKay DB, Bode W:
407 The metzincins--topological and sequential relations between the astacins,
408 adamalysins, serralysins, and matrixins (collagenases) define a superfamily of zinc-
409 peptidases. *Protein Sci* 1995;4:823-840.
- 410 11 Lawrence JB, Oxvig C, Overgaard MT, Sottrup-Jensen L, Gleich GJ, Hays LG, Yates
411 JR, 3rd, Conover CA: The insulin-like growth factor (IGF)-dependent IGF binding
412 protein-4 protease secreted by human fibroblasts is pregnancy-associated plasma
413 protein-A. *Proc Natl Acad Sci U S A* 1999;96:3149-3153.
- 414 12 Pihl, Kasper, et al.: First trimester maternal serum PAPP-A, β -hCG and ADAM12 in
415 prediction of small-for-gestational-age fetuses. *Prenatal Diagnosis*
416 28.12(2008):1131-1135.

417 13 Kuc, Sylwia, et al.: Evaluation of 7 Serum Biomarkers and Uterine Artery Doppler
418 Ultrasound for First-Trimester Prediction of Preeclampsia: A Systematic Review.
419 Obstetrical & Gynecological Survey 66.4(2011):225-239.

420 14 Meekins JW, et al.: A study of placental bed spiral arteries and trophoblast invasion
421 in normal and severe pre-eclamptic pregnancies. British Journal of Obstetrics and
422 Gynaecology 101.8(1994):669-674.

423 15 Vassart G and Dumont JE: The thyrotropin receptor and the regulation of thyrocyte
424 function and growth. Endocr Rev 13: 596-611, 1992.

425 16 Marques AC and Ponting CP: Intergenic lncRNAs and the evolution of gene expression.
426 Curr Opin Genet Dev 27: 48-53, 2014.

427 17 Chen G, Yin K, Shi L, Fang Y, Qi Y, Li P, Luo J, He B, Liu M and Shi T: Comparative
428 analysis of human protein-coding and noncoding RNAs between brain and 10
429 mixed cell lines by RNA-Seq. PLoS One 6: e28318, 2011.

430 18 Martens-Uzunova ES, Bottcher R, Croce CM, Jenster G, Visakorpi T, Calin GA: Long
431 noncoding RNA in prostate, bladder, and kidney cancer. Eur Urol 2014;65:1140-
432 1151.

433 19 Ribeiro AO, Schoof CR, Izzotti A, Pereira LV, Vasques LR: MicroRNAs: modulators
434 of cell identity, and their applications in tissue engineering. Microna 2014;3:45-
435 53.

436 20 Xiao Y, Jiao C, Lin Y, Chen M, Zhang J, Wang J, Zhang Z: lncRNA UCA1 Contributes
437 to Imatinib Resistance by Acting as a ceRNA Against miR-16 in Chronic Myeloid
438 Leukemia Cells. DNA Cell Biol 2017;36:18-25.

439 21 Li HJ, Li X, Pang H, Pan JJ, Xie XJ, Chen W: Long non-coding RNA UCA1 promotes
440 glutamine metabolism by targeting miR-16 in human bladder cancer. Jpn J Clin
441 Oncol 2015;45:1055-1063.

442 22 De Villiers CP, Hedley PL, Placing S, Wojdemann KR, Shalmi AC, Carlsen AL, Rode L,
443 Sundberg K, Tabor A, Christiansen M: Placental protein-13 (PP13) in combination
444 with PAPP-A and free leptin index (fLI) in first trimester maternal serum screening
445 for severe and early preeclampsia. Clin Chem Lab Med 2017;56:65-74.

446 23 Duley L: The global impact of pre-eclampsia and eclampsia. *Semin Perinatol*
447 2009;33:130-137.

448 24 Spradley FT, Palei AC, Granger JP: Immune Mechanisms Linking Obesity and
449 Preeclampsia. *Biomolecules* 2015;5:3142-3176.

450 25 Shaw J, Tang Z, Schneider H, Salje K, Hansson SR, Guller S: Inflammatory processes
451 are specifically enhanced in endothelial cells by placental-derived TNF-alpha:
452 Implications in preeclampsia (PE). *Placenta* 2016;43:1-8.

453 26 Herraiz I, Simon E, Gomez-Arriaga PI, Martinez-Moratalla JM, Garcia-Burguillo A,
454 Jimenez EA, Galindo A: Angiogenesis-Related Biomarkers (sFlt-1/PLGF) in the
455 Prediction and Diagnosis of Placental Dysfunction: An Approach for Clinical
456 Integration. *Int J Mol Sci* 2015;16:19009-19026.

457 27 Liu Z, Afink GB, Dijke PT: Soluble fms-like tyrosine kinase 1 and soluble endoglin
458 are elevated circulating anti-angiogenic factors in pre-eclampsia. *Pregnancy*
459 *Hypertens* 2012;2:358-367.

460 28 Wang F, Li X, Xie X, Zhao L, Chen W: UCA1, a non-protein-coding RNA up-regulated
461 in bladder carcinoma and embryo, influencing cell growth and promoting invasion.
462 *FEBS Lett* 2008;582:1919-1927.

463 29 Huang J, Zhou N, Watabe K, Lu Z, Wu F, Xu M, Mo YY: Long non-coding RNA UCA1
464 promotes breast tumor growth by suppression of p27 (Kip1). *Cell Death Dis*
465 2014;5:e1008.

466 30 Fan Y, Shen B, Tan M, Mu X, Qin Y, Zhang F, Liu Y: Long non-coding RNA UCA1
467 increases chemoresistance of bladder cancer cells by regulating Wnt signaling.
468 *FEBS J* 2014;281:1750-1758.

469 31 Li Z, Li X, Wu S, Xue M, Chen W: Long non-coding RNA UCA1 promotes glycolysis
470 by upregulating hexokinase 2 through the mTOR-STAT3/microRNA143 pathway.
471 *Cancer Sci* 2014;105:951-955.

472 32 Yang Y, Jiang Y, Wan Y, Zhang L, Qiu J, Zhou S, Cheng W: UCA1 functions as a
473 competing endogenous RNA to suppress epithelial ovarian cancer metastasis.
474 *Tumour Biol* 2016;37:10633-10641.

475 33 Aqeilan RI, Calin GA, Croce CM: miR-15a and miR-16-1 in cancer: discovery,
476 function and future perspectives. *Cell Death Differ* 2010;17:215-220.

477 34 Bonci D, Coppola V, Musumeci M, Addario A, Giuffrida R, Memeo L, D'Urso L,
478 Pagliuca A, Biffoni M, Labbaye C, Bartucci M, Muto G, Peschle C, De Maria R: The
479 miR-15a-miR-16-1 cluster controls prostate cancer by targeting multiple
480 oncogenic activities. *Nat Med* 2008;14:1271-1277.

481 35 Füsün Varol, et al.: VEGFR-1, Bcl-2, and HO-1 ratios in pregnant women with
482 hypertension. *Clinical and Applied Thrombosis/Hemostasis* 21.3(2013).

483 36 Nuzzo AM, et al.: JunB/Cyclin-D1 imbalance in placental mesenchymal stromal cells
484 derived from preeclamptic pregnancies with fetal-placental compromise. *Placenta*
485 35.7(2014):483-490.

486 37 Tan B, et al.: The Regulation of Trophoblastic p53 Homeostasis by the p38-Wip1
487 Feedback Loop is Disturbed in Placentas from Pregnancies Complicated by
488 Preeclampsia. *Cell Physiol Biochem*. 2019;52(2):315-335

489 38 Wald NJ, Watt HC, Hackshaw AK: Integrated screening for Down's syndrome based
490 on tests performed during the first and second trimesters. *N Engl J Med*
491 1999;341:461-467.

492 39 Conover CA, Kiefer MC, Zapf J: Posttranslational regulation of insulin-like growth
493 factor binding protein-4 in normal and transformed human fibroblasts. Insulin-like
494 growth factor dependence and biological studies. *J Clin Invest* 1993;91:1129-1137.

495 40 Durham SK, Kiefer MC, Riggs BL, Conover CA: Regulation of insulin-like growth
496 factor binding protein 4 by a specific insulin-like growth factor binding protein 4
497 proteinase in normal human osteoblast-like cells: implications in bone cell
498 physiology. *J Bone Miner Res* 1994;9:111-117.

499 41 Parker A, Gockerman A, Busby WH, Clemmons DR: Properties of an insulin-like
500 growth factor-binding protein-4 protease that is secreted by smooth muscle cells.
501 *Endocrinology* 1995;136:2470-2476.

502 42 Canini S, Prefumo F, Pastorino D, Crocetti L, Afflitto CG, Venturini PL, De Biasio P:
503 Association between birth weight and first-trimester free beta-human chorionic

504 gonadotropin and pregnancy-associated plasma protein A. Fertil Steril
505 2008;89:174-178.

506 43 Spencer K, Cowans NJ, Avgidou K, Molina F, Nicolaides KH: First-trimester
507 biochemical markers of aneuploidy and the prediction of small-for-gestational age
508 fetuses. Ultrasound Obstet Gynecol 2008;31:15-19.

509

510

Preprint

Characteristics	Control (N=172)	Mild PE (N=88)	Severe PE (N=32)	HELLP (N=8)	P value
Age, years	32.5 ± 5.3	33.1 ± 6.2	33.0 ± 5.8	31.5 ± 4.8	ns
BMI (kg/m ²)	21.5 ± 5.1	23.8 ± 6.5	24.4 ± 5.2*	25.4 ± 7.2*	<0.001
Gravida					ns
1 (%)	92 (53.5)	48 (54.5)	17 (53.1)	4 (50.0)	
2 (%)	45 (26.2)	24 (27.3)	9 (28.1)	2 (25.0)	
3 (%)	19 (11.0)	9 (10.2)	3 (9.4)	1 (12.5)	
>3 (%)	16 (9.3)	7 (8.0)	3 (9.4)	1 (12.5)	
Birth weight, g	3654.6 ± 167.7	3218.6 ± 182.4*	3239.7 ± 176.8*	3215.4 ± 147.9*	<0.001
GA at birth, days	285.6 ± 18.4	269.1 ± 17.9*	268.4 ± 17.5*	272.4 ± 21.5*	<0.001
GA sampling, days	92.6 ± 8.4	92.4 ± 6.8	91.9 ± 10.4	91.8 ± 6.9	ns

Table 1. demographic and clinicopathological characteristics of the participants of this study (*P<0.05 as compared with control)