

Clinical value of microRNA-19a-3p and microRNA-99a-5p in bladder cancer

Edyta M. Borkowska¹, Piotr Kutwin², Dorota Rolecka¹, Tomasz Konecki², Maciej Borowiec¹, Zbigniew Jabtonowski²

¹Chair of Laboratory and Clinical Genetics, Department of Clinical Genetics, Medical University of Lodz, Lodz, Poland

²1st Clinic of Urology, Medical University of Lodz, Lodz, Poland

Submitted: 13 August 2019

Accepted: 18 September 2019

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms.2019.89700>

Copyright © 2019 Termedia & Banach

Corresponding author:

Edyta M. Borkowska PhD

Chair of Clinical and

Laboratory Genetics

Department

of Clinical Genetics

Medical University

of Lodz

251 Pomorska St

92-213 Lodz, Poland

Phone: +48 42 272 53 58

E-mail: [edyta.borkowska@](mailto:edyta.borkowska@umed.lodz.pl)

[umed.lodz.pl](mailto:edyta.borkowska@umed.lodz.pl)

Abstract

Introduction: MicroRNAs (miRNA) are small (approximately 17 to 25 nucleotides in length), single stranded, non-coding RNAs that play an important role in the control of gene expression at the post-transcriptional stage, by inhibiting protein translation or promoting mRNA degradation. The main aim of the study was to evaluate the clinical utility of the tested markers (miRNAs 19a-3p and 99a-5p), which might be important in the diagnostics of non-invasive bladder cancer (BC).

Material and methods: The study involved a group of 60 patients suffering from BC (histopathologically confirmed), in which 20 patients were diagnosed with muscle invasive BC (INBC) and 40 patients with non-muscle invasive BC (NINBC). The control group consisted of 20 samples of normal urothelium, which did not show any cancerous changes during histopathological examination. We assessed the expression of microRNA, using real-time PCR and the miRCURY LNA Universal RT microRNA PCR Kit by Exiqon, Denmark.

Results: Reduced expression of both analyzed markers was observed in most cases: miR-19a-3p in 51.8% and miR-99a-5p in 65.5% (as follows Mann-Whitney *U* test $p < 0.000001$ and Student's *t* test $p = 0.034262$). Moreover, miR-19a-3p in our tested group was useful to differentiate between low and high grade disease in non-invasive stages (*t* test $p = 0.0315435$). Furthermore, miR-19a-3p and miR-99a-5p were able to discriminate patients in low grade for groups with or without recurrence.

Conclusions: Our data indicated that miR-19a-3p and miR-99a-5p were significantly altered in bladder cancer samples and useful as diagnostic markers.

Key words: bladder cancer, microRNA, genetic marker, early detection.

Introduction

MicroRNAs belong to the category of small, 17-25-nucleotide, single-stranded, non-coding RNAs, whose main function is post-transcriptional regulation of gene expression. These molecules are regarded as the “key” ones in the gene regulatory network [1]. The influence of microRNAs on the profile of gene expression cannot be ignored in the tumor transformation process. Various levels of microRNA expression have been observed in many cancers, including bladder cancer [1–4]. Molecular analyses indicated the existence of two main pathways visible in bladder cancer pathogenesis, where microRNA has a crucial role. These

are the signaling pathways of genes *FGFR3/RAS* and *TP53* [2, 5].

Receptor 3 of the fibroblast growth factor, *FGFR3*, belongs to the group of receptor tyrosine kinases. It plays a significant role in cell diversification and division as well as in angiogenesis. *FGFR3* gene mutations are rarely found in cancers and their occurrence has only been detected in bladder and cervical cancers [6]. In bladder cancer (BC), *FGFR3* gene mutations are found in about 80% of patients in non-invasive stages. These mutations are much rarer in invasive cancers, constituting approximately 16% of cases. The study has shown that *FGFR3* oncogenic function is associated with the stimulation of signaling pathways such as *RAS/MAPK*, the activation of which depends on adaptor proteins *GRB2* and *SOS*. *SOS* is the factor changing the guanine nucleotide *GDP* into *GTP*, thereby activating protein *RAS*. MicroRNA 99a-5p (miR-99a-5p) has its anchor point at the mRNA level, which is this oncogene. About 60% of patients with invasive bladder cancer have changes in the gene *TP53* [7]. It is one of the most important tumor suppressors, described as the guardian of the genome. Protein *P53* coded by this gene is an inhibitor of cell growth, which leads to arresting the cell cycle and/or apoptosis. In both mentioned pathways, microRNA plays a significant role as a negative regulator of gene expression at the post-transcriptional level. Besides other genes, microRNA 19a-3p

(miR-19a-3p) targets the suppressor gene *PTEN* (phosphatase and tensin homolog). The product of this gene is a negative regulator of the *PI3K/Akt* pathway, and therefore it is involved in regulating basic cell processes, including proliferation and apoptosis. It is the result of cell cycle control at the transfer point from stage *G1* to *S*.

The purpose of the study was to analyze changes in miR-99a-5p and miR-19a-3p expression in samples of BC patients' tumor tissues and to determine whether the observed changes in the expression level of the above-mentioned microRNAs can be used as a diagnostic and prognostic marker in bladder cancer.

Material and methods

The tested group consisted of tumor tissue samples stored in the tissue bank in the Department of Clinical Genetics, the Chair of Clinical and Laboratory Genetics, Medical University of Lodz. The tumor tissues were taken during the TURBT examination (transurethral resection of bladder tumor) at the Urology Ward of the University Clinical Hospital, Military Memorial Medical Academy in Lodz. The official permission to conduct examinations was granted by the Bioethics Advisory Commission at Lodz Medical University, No. RNN/62/15/ KE/M, and the patients signed the formal consent. The fragments selected for RNA isolation were submerged in the special stabilization RNAlater solution (Sigma) and stored at -20°C until the isolation time. The tests were carried out on a group of 60 patients with diagnosed bladder cancer (details presented in Table I). The age range was 40–89, with the average 69.4. Most of the patients were male (49/60 – 81.7%). All the tumors were of urothelial origin. Forty (66.7%) patients in the group suffered from non-invasive bladder cancer, among whom 20 (33.35%) patients were in stage Ta and 20 (33.35%) patients in stage T1. The last 20 (33.3%) patients were diagnosed with invasive bladder cancer in stage T2. The criterion which decided about the patients' selection for the tested group was the histopathological recognition, confirmed by a clinical physician. The additional criteria were: primary tumor patients and patients with no oncology treatment history. Tumors were graded according to the WHO classification and staged using the TNM criteria [8–10]. Progressive disease was defined as progression to stage T2 or higher stage of disease, development of nodal or distant metastases. The control group (C) consisted of 20 patients admitted to the Urology Ward. They underwent control cystoscopy, the aim of which was to confirm or to exclude tumor changes in the bladder. The examination did not reveal any tumor changes. All the controls were age and sex matched and had no history of any tumors.

Table I. Characteristics of patients

Parameter	Results
Age, mean (range) [years]	69.4 (40–89)
Follow-up, median (range) [months]	31 (5–34)
Gender:	
Male	49 (81.7%)
Female	11 (18.3%)
Smoking (ex-or current)	52 (86.7%)
Occupational exposure	21 (35%)
Stage:	
pTa	20 (33.35%)
pT1	20 (33.35%)
pT2	20 (33.35%)
Grade:	
LG	71 (51.4%)
HG	34 (24.6%)
Recurrence	14 (23.3%)
Progression	26 (43.3%)

Methods

The MirVana miRNA Isolation Kit (Life Technologies, Cat No 1556) was used to isolate microRNA from the tumor tissues. The whole procedure was followed in accordance with the producer's guidelines. After the completion of the isolation procedure, the amount and purity of the obtained microRNA were determined, applying the spectrophotometric method on the NanoDrop ND-1000 instrument. The purity measurement of the obtained extracts is the relationship A260/230 and A260/280. It is accepted that for good quality nucleic acids it is respectively 1.8–2.2 and 1.8–2.0. The measurement results of the samples selected for further analysis met the required criteria.

The next step was to conduct the synthesis reaction of cDNA on the microRNA transcript using the miRCURY LNA Universal RT microRNA PCR set (Exiqon, Cat No. 203351, T100 Thermal Cycler, Bio-Rad), following the manufacturer's protocol (briefly: 5× Reaction Buffer 2 µl, Reverse Transcriptase, 50 U/µl 1 µl, Synthetic RNA Spike ins 0.5 µl, microRNA 5 ng/µl 2 µl, Nuclease-free Water 4.5 µl). First, all the samples were diluted until the final concentration of 5 ng/µl in a 10 µl volume was obtained. The RNA Spike-in synthetic transcript, attached to the set, was used for quality control of the described reaction. The cDNA obtained in this way was stored at –20°C, and then it was used as a transcript for the real time-PCR (CFX96 Real-Time PCR Detection System, Bio-Rad). The expression level of the selected microRNAs was assessed using the miRCURY LNA Universal RT microRNA PCR set (Exiqon, Cat No. 203403). The first step involved preparing the 1/80 dilution of the obtained cDNA by suspending 5 µl of the reaction mixture containing cDNA in 395 µl of the Ultra Pure Water. Validated hsa-mir-99a-5p and hsa-mir-19a-3p were used for the amplification (hsa-mir-19a-3p: 5'UGUGCAAUCUAUGCAAACUGA3' and hsa-mir-99a-5p: 5'AACCCGUAGAUCCGAUCUUGUG3'). In addition, the commercial set which was used in our tests provided double control. Firstly, there was internal control as the so-called RNA Spike-in UniSp6, the synthetic transcript, enabled validity and quality control of all the stages of our analysis, starting from the microRNA isolation. Secondly, there was also endogenous control. Reference microRNA hsa-mir-103a-5p was used to normalize the measures of the expression level of the selected markers. Each sample was run in duplicate at a final volume of 10 µl containing 5 µl of ExiLent SYBR Green master mix, 2× concentrated, PCR primer mix 1 µl, diluted cDNA 4 µl. Each PCR included no template control and all of them were negative. The PCR reaction was heated to 95°C for 10 min; after that there were 40 cycles of 95°C for 10 s and 60°C for 1 min. The specificity of the

reaction was checked through the analysis of the melting curve of PCR reaction products.

Methods of analysis of obtained results

The $\Delta\Delta Ct$ method, also defined as the comparative method, was applied to determine the relative expression level of the examined microRNAs. The method is based on mathematical calculations which enable one to determine the relative difference in the expression level of the tested marker between unknown samples and the reference. The first stage consisted of the analysis of marked threshold cycles (Ct) in the amplification reaction of the examined microRNAs and the control microRNAs for both the tested and control group. The calculated expression level of each patient was normalized against the endogen control – miR-103a-5p. After that, the difference of the tested and the control microRNAs (ΔCt) was calculated for individual samples. The calculations were performed using the formula $FC = 2^{-\Delta\Delta Ct}$. The results obtained in this way are expressed as the multiple of the control trial [11]. The obtained values were divided into the classes of increased, decreased or unchanged expression levels on the basis of the calculated average ± 2 SEM (standard error of the mean).

Statistical analysis

The program Statistica 13, Stat-Soft Inc was used to make statistical calculations. The normality of the distribution of variables was verified by the Shapiro-Wilk test and parametric test. The analysis of the unrelated variables was performed with the t test and Mann-Whitney U test. The analysis of relationships between the expression of the selected microRNAs and the selected clinical parameters was carried out by means of the accurate Pearson's χ^2 test. The value $p < 0.05$ was accepted as the threshold of statistical difference or correlation significance. The area under the receiver operating characteristic curve (ROC) was calculated (results presented in Table II and Figure 1).

Results

The statistical analysis using the non-parametric Mann-Whitney U test indicated significant statistical differences in the expression level of the two selected microRNAs, miR-19a-3p and miR-99a-5p, between the tumor tissue and the normal urothelium (miR-99a-5p $p = 0.34262$, miR-19a-3p $p < 0.001$).

Analysis of microRNA-19a-3p expression level

Fifty-six patients were subjected to analysis (37 with NINBC and 19 with INBC); 4 patients

Table II. 7 test results of miRs for the appropriate subgroup of patients

t test for HG	Progression (yes/no: 26/11)	Recurrence (yes/no: 13/24)	Stage (TaT1/T2: 18/19)
miR19a	0.2780225	0.37691	0.014332
miR99	0.228834	0.281592	0.451997
t test for LG	Progression (yes/no: 6/13)	Recurrence (yes/no: 14/5)	Stage (TaT1/T2 = 19/0)
miR19a	0.21001*	0.033884*	–
miR99	0.1598295	0.035877	–
t test for TaT1	Progression (yes/no: 16/21)	Recurrence (yes/no: 22/15)	Grade (HG/LG: 18/19)
miR19a	0.4116115	0.3017835	0.0315435
miR99	0.1862725	0.075472	0.4197025
t test for T2	Progression (yes/no: 16/3)	Recurrence (yes/no: 5/14)	Grade (HG/LG = 19/0)
miR19a	0.264414	0.3949525	–
miR99	0.1658645	0.2444815	–

p* – Mann-Whitney U test, **boldface** represents p-values < 0.05.




Figure 1. Receiver operating characteristic (ROC) curve analysis of miR-19a-3p: **A** – ROC curve for detecting bladder cancer compared with the control group, **B** – ROC curve differentiating the TaT1 group with the control group, **C** – ROC curve for differentiating the T2 group with the control group

were rejected due to unsuccessful amplification in the rt-PCR reaction. Decreased expression was observed in most cases – 29 (51.8%) patients. Thirteen (23.2%) patients had increased expression whereas an unchanged expression level was observed in 14 (25%) patients.

Analysis of microRNA-99a-5p expression level

Fifty-eight patients were subjected to analysis (39 with NINBC and 19 with INBC); 2 patients were rejected due to unsuccessful amplification in the rt-PCR reaction. The expression level of the tested

microRNA was mostly decreased – 38 (65.5%) patients. The other 20 patients had an unchanged expression level (17.25%) or an increased expression level (17.25%). Results presented in Figure 2.

There was an analysis of the expression levels of the two tested microRNAs, miR-19a-3p and miR-99a-5p, in terms of stage and grade. Pearson’s χ^2 test was applied for it. The analysis at the 95% significance level showed no statistically significant relationships.

No correlation was found between the expression level of the tested microRNAs and other clinical parameters, smoking or occupational ex-




Figure 2. Expression level of miR-19a-3p (A) and miR-99a-5p (B) in tumor tissue compared to the control group. Results based on normalization to miR-103a-5p. Analysis of changes in expression level according to clinical stage: miR-19a-3p (C), miR-99a-5p (D) and grade of cancer: miR-19a-3p (E), miR-99a-5p (F)

NIBC – non-invasive bladder cancer, INBC – invasive bladder cancer, LG – low grade, HG – high grade.

posure. Also no statistical relationship was found between the patients' sex or age and the expression level of the analyzed markers.

The *t* test was applied to assess the average values of individual miRs for the appropriate subgroup of patients. Only miR-19a was found to be statistically significant in all the subgroups (LG, HG, TaT1, T2) compared to the control group ($p < 0.001$ for all subgroups). Moreover, the analysis for progression and/or recurrence showed that only miR-19a differentiated patients in TaT1 and T2 stages in HG. In the group of patients in a high grade, those in clinical stage TaT1 are characterized by significantly low values of miR-19a-3p in comparison to the patients of the same grade and more advanced stage (T2). Similarly, in the group of TaT1 patients with a low grade of the disease, miR-19a-3p values are significantly higher than for patients with the same stage and higher grade. However, both miRs differentiated patients who have recurrence from those without recurrence (results presented in Table II). Receiver operating characteristic curve analysis showed that only miR-19a-3p is a classifier with a significantly different area under the curve (AUC) compared with the control group. Moreover, it helps to differentiate patients in both stage and grade (demonstrated in Table III and Figure 1).

Discussion

Bladder cancer is one of the most frequent urinary tract tumors [12]. In Poland it is the fifth cause

of death from malignant diseases diagnosed in male patients. At present, urine sediment cytology with cystoscopy is the primary tool in bladder cancer diagnosis and the stage and grade recognized in histopathological examination are the most reliable prognostic markers [10]. The high risk of recurrence and progression of the disease towards invasive changes makes it important to identify markers enabling both diagnosis of changes in the early stage and monitoring of patients. Numerous scientific reports suggest that besides *TP53* gene mutation, other molecular factors are of significant importance in bladder cancer etiopathogenesis [2, 3], that is, post-transcriptional regulation of expression of genes involved in carcinogenesis by small non-coding RNA molecules – microRNAs.

The presented analyses of the miR-19a-3p expression level showed that the expression was decreased in 51% of patients and overexpression was observed only in 23.3% of the tested group. As it is assumed in many scientific reports, there may be a relationship between this microRNA and the suppressor gene *PTEN*. For example, the research of Feng *et al.* [13] showed that at the mRNA level, the target for miR-19a-3p in bladder cancer may be the above-mentioned gene. The analyses were performed with the Western blot techniques, and later they were confirmed with the “PTEN expression plasmid” technique. There were also miR-19a-3p expression analyses performed on cell lines and on 100 tumor tissue samples obtained from bladder cancer patients using real-time poly-

Table III. Results of area under receiver operating characteristic curve (ROC)

ROC	AUC	95% CI	P-value
BC case/control = 56/20:			
miR19a	0.948	0.90–0.99	< 0.0001
miR99a	0.606	0.45–0.76	0.1762
HG (case/control = 37/20):			
miR19a	0.951	0.902–1,000	< 0.0001
miR99a	0.608	0.443–0.774	0.2006
LG (case/control = 19/20):			
miR19a	0.941	0.873–1,000	< 0.0001
miR99a	0.603	0.423–0.783	0.2638
TaT1 (case/control = 37/20):			
miR19a	0.97	0.933–1,000	< 0.0001
miR99a	0.607	0.442–0.772	0.2045
T2 (case/control = 19/20):			
miR19a	0.905	0.813–0.997	< 0.0001
miR99a	0.605	0.425–0.785	0.2511

Boldface represents *p*-values < 0.05.

merase chain reaction (rt-PCR). They revealed that the expression level of this molecule is increased considerably in bladder cancer patients, which additionally correlates with a more invasive bladder cancer phenotype. Our research pointed to the opposite tendency in terms of expression. Moreover, it did not confirm the relationship between the expression level and tumor stage or grade, but we found that miR-19a-3p is a good classifier for bladder cancer patients compared to healthy people. In addition, Feng *et al.* made an assessment of the miR-19a expression level in blood plasma, obtaining results similar to those in tissue. They concluded that the loss of function of miR-19a can promote cell growth of bladder cells. According to Shi *et al.* [14], there were some publications which also pointed to decreased miR-19a-3p expression in bladder cancer patients; however, the anchor point for this molecule was not identified at the mRNA level. It is possible that the microRNA is responsible for the regulation of other genes the relationship with which is not known yet. For instance, the research presented by Bush *et al.* [15], and confirmed by the tests with the reporter probe, showed that the direct target gene for miR-19a-3p is the gene for 5-LO (5-lipoxygenase), the enzyme included in the leukotriene biosynthesis pathway, whose deregulations can also lead to the development of tumors. Lu *et al.* [16] researched the significance of microRNA for prostate cancer using samples of patients' tumor tissues for tests. They showed that miR-19a-3p is responsible for regulation of the suppressor gene *BTG1* (B-cell translocation gene 1) and its expression is increased in tumors. The above-mentioned examples refer to piloting examinations, which are also an introduction to analysis of miR-19a-3p as a marker in tumor diagnosis, which was confirmed in our study. miR-19a-3p is useful in differentiating LG and HG patients in the TaT1 stage. A recently published study showed that miR-19a could promote the invasion and epithelial-mesenchymal transition (EMT) of bladder cancer cells through inhibiting the expression of RhoB [17].

Regarding the other microRNA 99a-5p, which was analyzed in this study, its decreased expression was observed in most of the examined patients (65%). Performing analyses on cell lines and on 100 tumor tissue samples of bladder cancer patients using the rt-PCR technique, Feng *et al.* also observed decreased expression of this microRNA, which correlated with a more malignant cancer phenotype [18]. What is more, they obtained similar results when using those patients' blood plasma for their analyses. Catto *et al.* carried out tests on 72 samples of normal and changed urothelium and on 6 cell lines [19]. Analyzing 322 microRNAs with the microarray and rt-PCR techniques, they

investigated the relationships between the level of the selected markers and the stage. Among those markers, the expression of miR-99a was significantly decreased, especially in non-invasive bladder cancer patients. Similar results were observed by Wu *et al.*, who carried out examinations on the T24 and EJ cell lines with the Western blot techniques and reporter probe tests [20]. They observed a decreased expression level of this microRNA, which related to the overexpression of its target gene – the *FGFR* oncogene. Moreover, Liu *et al.* reported that miR-99a expression inhibited bladder cell proliferation and the cell cycle [21]. Bladder cancer patients with low expression of miR-99a had a poor survival rate in their study. Tsai *et al.* confirmed that miR-99a markedly inhibits the growth of bladder cancer cells via dual inhibition of mTORC1 and mTORC2 [22]. Finally, Inamoto *et al.* found that overexpression of miR-99a is associated with high risk of an aggressive phenotype [23]. In our study miR-99a and miR-19a discriminate patients in LG with and without recurrence.

Another point of our analysis worth concentrating on is the question whether using control miR-103-5p had any effect on the ultimate results of the conducted tests. Hofbauer *et al.* used the same endogenous controls in their study. They assumed that it does not necessarily apply to any of the used miRs [24]. We analyzed the results of Peltier *et al.*, which revealed that miR-103 and miR-191 were the most stable microRNAs in terms of their expression in tissue material from lung cancer patients and in 13 other kinds of unaffected tissues [25]. The microarray technique was used for their tests. Ratert *et al.*, who also worked on the microRNA expression level in urothelial tumors with the rt-PCR technique, selected two groups of reference microRNAs using the microarray technique [26]. Based on the observed results, they stated that the combination of miR-101, miR-125a-5p, miR-148b and miR-151-5p or the combination of miR-148b, miR-181b and miR-874 is most accurate for obtaining the most reliable results. Referring to the above, in the analyses of Feng *et al.* and Han *et al.*, U6 snRNA (U6 small nuclear RNA) was used as the reference [18, 27]. In the tests of Catto *et al.*, three reference RNA markers were applied and the normalization was done using their average *Ct* [19]. The authors, however, do not give any information on the types of RNA molecules which were used. Other examples are the research of Hirata *et al.*, who chose GAPDH (glyceraldehyde 3-phosphate dehydrogenase gene) and U48 snRNA. Jia *et al.* decided on miR-92b. In our opinion, the selection of the right microRNA for expression level normalization with the rt-PCR technique is a vital element influencing

the reliability of the results [28, 29]. Most of the above-presented results were obtained in some introductory research, which was carried out on small groups of patients and on cell lines with different methods and devices.

In conclusion, microRNAs are essential molecules regulating genetic material translation. In comparison to normal urothelium, significant differences were observed in tumor tissue. Moreover, miR-19a-3p in our tested group was useful to differentiate between low and high grade disease in NINBC stages. Furthermore, miR-19a-3p and miR-99a-5p were able to distinguish patients with LG disease into groups with or without recurrence. It might be a potential non-invasive marker of recurrence. It is necessary to conduct further analysis in urine or plasma on a bigger group of patients.

Acknowledgments

Edyta M. Borkowska and Piotr Kutwin equally contributed to this work.

The authors are grateful to all participants who took part in the study and agreed to sharing of their biological samples. This study was supported by Medical University Grant No. 502-03/5-138-02/502-54-146 and Grant No. 502-03/5-138-02/502-54-133.

Conflict of interest

The authors declare no conflict of interest.

References

- Iorio MV, Croce CM. MicroRNA dysregulation in cancer: diagnostics, monitoring and therapeutics. A comprehensive review. *EMBO Mol Med* 2012; 4: 143-59.
- Golbakhsh MR, Boddouhi B, Hatami N, et al. Down-regulation of microRNA-182 and microRNA-183 predicts progression of osseosarcoma. *Arch Med Sci* 2017; 13: 1352-6.
- Liu P, Chen Y, Wang B, Wang Z, Li C, Wang Y. Expression of microRNAs in the plasma of patients with acute gouty arthritis and the effects of colchicine and etoricoxib on the differential expression of microRNAs. *Arch Med Sci* 2019; 15: 1047-55.
- Lenher SM, Tsai S, Silva Neto B, et al. MicroRNA expression profile identifies high grade, non-muscle invasive bladder tumors at elevated risk to progress to an invasive phenotype. *Genes* 2017; 8: 77.
- Dip N, Reis ST, Timoszczuk LS, et al. Stage, grade and behavior of bladder urothelial carcinoma defined by the microRNA expression profile. *J Urol* 2012; 188: 1951-6.
- Hart KC, Robertson SC, Kanemitsu MY, Meyer AN, Tynan JA, Donoghue DJ. Transformation and Stat activation by derivatives of FGFR1, FGFR3, and FGFR4. *Oncogene* 2000; 19: 3309-20.
- Guancial EA, Bellmunt J, Yeh S, Rosenberg JE, Berman DM. The evolving understanding of microRNA in bladder cancer. *Urol Oncol* 2014; 32: 41.e 31-40.
- Bircan S, Candir O, Serel TA. Comparison of WHO 1973, WHO/ISUP 1998, WHO 1999 grade and combined scoring systems in evaluation of bladder carcinoma. *Urol Int* 2004; 73: 201-8.
- Compérat EM, Burger M, Gontero P, et al. Grading of urothelial carcinoma and the new "World Health Organisation Classification of Tumours of the Urinary System and Male Genital Organs 2016". *Eur Urol Focus* 2018; 73: 560-9.
- Schmitz-Dräger BJ, Droller M, Lokeshwar VB, et al. Molecular markers for bladder cancer screening, early diagnosis, and surveillance: the WHO/ICUD consensus. *Urol Int* 2015; 94: 1-24.
- Bustin SA, Benes V, Garson JA, et al. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 2009; 55: 611-22.
- Antoni S, Ferlay J, Soerjomataram I, Znaor A, Jemal A, Bray F. Bladder cancer incidence and mortality: a global overview and recent trends. *Eur Urol* 2017; 71: 96-108.
- Feng Y, Liu J, Kang Y, He Y. miR-19a acts as an oncogenic microRNA and is up-regulated in bladder cancer. *J Exp Clin Cancer Res* 2014; 33: 67.
- Shi Z, Wei Q. MicroRNAs in bladder cancer: expression profiles, biological functions, regulation, and clinical implications. *Crit Rev Eukaryot Gene Expr* 2014; 24: 55-75.
- Busch S, Auth E, Sholl F, et al. 5-lipoxygenase is a direct target of miR-19a-3p and miR-125b-5p. *J Immunol* 2015; 194: 1646-53.
- Lu K, Liu C, Tao T, et al. MicroRNA-19a regulates proliferation and apoptosis of castration-resistant prostate cancer cells by targeting BTG1. *FEBS Lett* 2015; 589: 1485-90.
- Li Z, Li Y, Wang Y. miR-19a promotes invasion and epithelial to mesenchymal transition of bladder cancer cells by targeting RhoB. *J BUON* 2019; 24: 797-804.
- Feng Y, Kang Y, He Y, et al. MicroRNA-99a acts as a tumor suppressor and is down-regulated in bladder cancer. *BMC Urol* 2014; 14: 50.
- Catto JW, Miah S, Owen HC. Distinct microRNA alterations characterize high- and low-grade bladder cancer. *Cancer Res* 2009; 69: 8472-81.
- Wu D, Zhou Y, Pan H, Zhou J, Fan Y, Qu P. MicroRNA-99a inhibiting cell proliferation, migration and invasion by targeting fibroblast growth factor receptor 3 in bladder cancer. *Oncol Lett* 2014; 7: 1219-24.
- Liu Y, Li B, Yang X, Zhang C. MiR-99a-5p inhibits bladder cancer cell proliferation by directly targeting mammalian target of rapamycin and predicts patients survival. *J Cell Biochem* 2018; 1-8; doi: 10.1002/jcb.27318.
- Tsai TF, Lin JF, Chou KY, Lin YC, Chen HE, Hwang TIS. miR-99a-5p acts as tumor suppressor via targeting to mTOR and enhanced RAD001-induced apoptosis in human urinary bladder urothelial carcinoma cells. *Onco Targets Ther* 2018; 11: 239-52.
- Inamoto T, Uehara H, Akao Y, et al. A panel of microRNA signature as a tool for predicting survival of patients with urothelial carcinoma of the bladder. *Dis Markers* 2018; 2018: 5468672.
- Hofbauer SL, de Martino M, Lucca I, et al. A urinary microRNA (miR) signature for diagnosis bladder cancer. *Urol Oncol* 2018; 36: 531.e1-8.
- Peltier HJ, Latham GJ. Normalization of microRNA expression levels in quantitative RT-PCR assays: identification of suitable reference RNA targets in normal and cancerous human solid tissues. *RNA* 2008; 14: 844-52.
- Ratert N, Meyer HA, Jung M, et al. Reference miRNAs for miRNAome analysis of urothelial carcinomas. *PLoS One* 2012; 7: e39309.

27. Han Y, Chen J, Zhao X. MicroRNA expression signatures of bladder cancer revealed by deep sequencing. *PLoS One* 2011; 6: e18286.
28. Hirata H, Ueno K, Shahryari V, et al. Oncogenic miRNA-182-5p targets Smad4 and RECK in human bladder cancer. *PLoS One* 2012; 7: e51056.
29. Jia AY, Castillo-Martin M, Bonal DM, et al. MicroRNA-126 inhibits invasion in bladder cancer via regulation of ADAM9. *Br J Cancer* 2014; 110: 2945-54.