

Involvement of PI3K/Akt/ β -catenin signaling in schisandrin B-mitigated bone deterioration in an experimental rat model of estrogen deficiency

Yimin Liang¹, Wei Li², Xiang Li¹, Jun Nan³

¹Department of Orthopedics, The First People's Hospital of Taizhou, Taizhou, China

²The Second Clinical School of North Sichuan Medical College, Nanchong, China

³Department of Spine Surgery, Affiliated Hospital of Yanbian University, Yanji, China

Submitted: 10 July 2019

Accepted: 2 October 2019

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms.2020.92873>

Copyright © 2020 Termedia & Banach

Corresponding author:

Dr. Jun Nan

Department of

Spine Surgery

Affiliated Hospital

of Yanbian University

1327 Juzi Road

Yanji 133000, China

Phone: +86 0433-2660074

Fax: +86 0433-2660074

E-mail: nannan_junfmxm12@sohu.com

sohu.com

Abstract

Introduction: Schisandrin B (SchB) has been reported to perform a wide range of biological functions, including antioxidant activity, anti-inflammatory activity and stimulation of osteoblast proliferation. However, the function and mechanism of SchB in ovariectomy (OVX)-induced osteoporosis are still unknown. The present study was designed to investigate the anti-osteoporotic activity of SchB in an experimental rat model of estrogen deficiency, which is usually used to mimic human postmenopausal osteoporosis (PMO).

Material and methods: OVX rats were orally treated with low (10 mg/kg) or high (50 mg/kg) doses of SchB for 8 weeks. Bone metabolism-related markers were measured by ELISA. The levels of protein expression were determined by western blotting analysis. Hematoxylin and eosin (H&E) and safranin O staining were performed to analyze trabecular bone and cartilage degeneration. Tartrate-resistant acid phosphatase (TRAP) staining was used to evaluate osteoclast differentiation.

Results: SchB administration markedly increased serum Ca levels and bone Ca content and decreased urinary calcium excretion in OVX-operated rats. In addition, high-dosage SchB treatment blocked osteoclastogenesis and improved trabecular bone and cartilage degeneration in the tibia of OVX-operated rats. Furthermore, high-dosage SchB treatment dramatically elevated the protein expression of phospho-PI3K, phospho-Akt and β -catenin in OVX-operated rats.

Conclusions: SchB exerted anti-osteoporotic activity in OVX-operated rats by accelerating the phosphorylation of PI3K and Akt, subsequently upregulating the expression of β -catenin.

Key words: schisandrin B, ovariectomy, osteoporosis, PI3K/Akt.

Introduction

Postmenopausal osteoporosis (PMO) is a global healthcare problem that affects approximately 40% of postmenopausal women worldwide [1]. Estrogen deficiency is believed to be the primary cause of PMO, which is attributed to a disequilibrium between osteoblast-mediated bone formation and osteoclast-mediated bone resorption and eventually results in osteoporotic fracture [2–5]. At present, there are no ideal therapeutic drugs for the treatment of PMO. For example, hormone replacement therapy (HTR)

has been widely used in the prevention of PMO by counteracting the loss of estrogen in postmenopausal women, while emerging side effects, such as increasing the risk of endometrial cancer, cardiovascular diseases and atypical fractures, limit its clinical applications [3, 6]. There is mounting evidence that traditional Chinese herbal medicines and their bioactive components, including *Fructus ligustri lucidi* [7], *Sambucus williamsii* [8], *Epimedium* [9] and *Polygonum multiflorum* [10], exert beneficial activity against ovariectomy (OVX)-induced osteoporosis. However, the precise mechanisms regarding the anti-osteoporotic activity of traditional herbal medicines have not been completely clarified.

Schisandrin B (SchB) is the main bioactive ingredient derived from *Schisandra chinensis*, and it performs a variety of biological functions, including antioxidant, anti-inflammatory and anti-DNA oxidative damage activities [11–13]. It has also been reported that SchB prevents cartilage degeneration in a rat osteoarthritis model [14]. Intriguingly, active fractions from *Schisandra chinensis* stimulate osteoblast proliferation and increase alkaline phosphatase activity *in vitro*, suggesting that they may possess potential activity against osteoporosis [15]. However, the underlying molecular mechanisms and the effects of SchB on OVX-induced osteoporosis have not been reported in an experimental rat model of estrogen deficiency.

In the present study, we report for the first time that SchB alleviated OVX-induced osteoporosis by activating the phosphoinositide 3-kinase (PI3K)/protein kinase B (AKT) signaling pathway. In osteoporotic rats, impairment of PI3K/AKT signaling is associated with bone loss [16, 17]. *In vitro* experiments also show that upregulation of PI3K/AKT activity facilitates osteoblast differentiation and increases osteoblast-specific gene expression [18, 19]. Overactivation of PI3K/AKT signaling can also modulate expression of β -catenin protein, which is a vital component in the wnt/ β -catenin pathway and participates in osteogenesis [20]. We investigated the osteoprotective effects of SchB, and the results demonstrated that SchB could prevent Ca excretion and bone loss in an ovariectomized rat model. The underlying mechanism was mediated, at least partially, through activation of PI3K/AKT signaling in the bone. These findings suggested that SchB might serve as a potential therapeutic agent for the treatment of osteoporosis.

Material and methods

Animal treatment

Two-month-old female Sprague-Dawley rats ($n = 60$; body weight: 200 ± 20 g) were purchased from Slac Laboratory Animal, Shanghai, China and allowed to acclimate to the environment for

1 week. After 1 week of sham or OVX operation, the rats in the sham-operated (Sham) group and OVX group received sodium carboxymethylcellulose (CMC) treatment with intragastric administration; the rats in the estradiol valerate (EV) group underwent OVX operation combined with EV treatment (1 mg/kg/day), EV (1 mg/ml) was dissolved in 0.5% CMC with intragastric administration; the rats in the SchB-L group underwent OVX operation combined with low-dosage SchB (SchB-L; 10 mg/kg/day) treatment, SchB was dissolved in 0.5% CMC (10 mg/ml CMC) with intragastric administration; the rats in the SchB-H group underwent OVX operation combined with high-dosage SchB (SchB-H; 50 mg/kg/day) treatment, SchB was dissolved in 0.5% CMC (50 mg/ml CMC) with intragastric administration. Twelve rats were assigned to each group ($n = 12$ in each group). EV was obtained from Elpharm Lille SAS (Paris, France) and SchB was obtained from Shanghai PureOne Biotechnology (Shanghai, China). After 8 weeks of treatment, the rats were killed by an overdose of sodium pentobarbital (2%; 200 mg/kg; Sigma-Aldrich, Merck KGaA, Germany) 24 h after their last treatment. Serum and tibias were immediately collected and stored at -80°C for further analysis. In addition, tibias were fixed with 4% formalin at room temperature and embedded in paraffin for histomorphological analysis. The experiment was approved by the Ethics Committee of Affiliated Hospital of Yanbian University (Yanji, China) and was performed in accordance with its guidelines.

Biochemical markers in serum and urine

The levels of calcium (Ca), phosphorus (P) and creatinine (Cre) in serum and urine were measured by commercial kits (Nanjing Jiancheng Biology Engineering Institute, Nanjing, China) according to the manufacturer's protocol. Alkaline phosphatase (ALP; Elabscience Biotechnology Co., Ltd. Wuhan, China), osteocalcin (Immutopics, Inc., San Clemente, CA, USA), tartrate-resistant acid phosphatase-5b (TRAP-5b; Immunodiagnostic Systems, Scottsdale, AZ, USA), and C-terminal telopeptide of type 1 collagen (CTX; Elabscience Biotechnology Co., Ltd. Wuhan, China) were detected by rat bioactive ELISA kits according to each manufacturer's protocol. Serum intact parathyroid hormone (PTH) and osteocalcin were measured using chemiluminescence immunoassays (Roche Diagnostics, Mannheim, Germany). Serum 25(OH) vitamin D₃ assay was performed using a DiaSorin kit on a LIAISON automated immunoassay analyzer (DiaSorin, Saluggia, Italy).

Bone calcium content and bone mineral density (BMD)

To measure Ca content in the tibia, the tibia was incinerated using a muffle furnace (Ther-

mo Fisher Scientific, Inc., Waltham, MA, USA) at 800°C for 12 h, and then 10 mg of bone ash was dissolved in 1 ml of 37% HCl diluted with Milli-Q water. The calcium content was determined using a commercial kit (Nanjing Jiancheng Biology Engineering Institute) according to the manufacturer's protocol. The BMD of the femur and tibia was measured by dual-energy X-ray absorptiometry (DEXA) (Lunar DPXIQ, GE Healthcare, USA).

Histomorphology and TRAP staining

Tibias were collected immediately following sacrifice and fixed with 4% formalin at room temperature for 24 h, then decalcified in 0.5 M EDTA (pH = 8.0) and embedded in paraffin. Sections of 5 μ m were cut and stained with hematoxylin and eosin (H&E; Beyotime Institute of Biotechnology, Haimen, China) and safranin O (Nanjing SenBeiJia Biological Technology Co., Ltd., Nanjing, China) and visualized under a microscope (Leica DM 2500).

TRAP staining was used for the identification of osteoclasts according to the manufacturer's instructions (Sigma-Aldrich). The number of osteoclasts was quantified below the growth plates in the proximal tibial metaphysis and visualized under a microscope (Leica DM 2500).

Western blotting

Proteins were extracted with radio immunoprecipitation assay (RIPA) buffer (Beyotime Institute of Biotechnology, Haimen, China). Western blotting assays were performed as previously described [21]. After blocking with 5% skim milk, the membrane was incubated with primary antibodies as follows: anti-PI3K (dilution: 1 : 1,000; Cell Signaling Technology, USA), anti-p-PI3K (dilution: 1 : 1,000; Cell Signaling Technology, USA), anti-Akt (dilution: 1 : 500; Santa Cruz Biotechnology), anti-p-Akt (dilution: 1 : 500; Santa Cruz Biotechnology) and anti- β -catenin (dilution: 1 : 500; Santa Cruz Biotechnology). After incubation with primary antibodies, the membranes were incubated with the appropriate horseradish peroxidase-conjugated secondary antibody (cat. no. sc-516102; dilution:

1 : 10,000; Santa Cruz Biotechnology), followed by visualization using chemiluminescence (Thermo Fisher Scientific, Inc.). β -actin (cat. no. sc-130065; 1 : 2,000; Santa Cruz Biotechnology) was used as the loading control antibody. Signals were analyzed with Quantity One software version 4.5 (Bio-Rad).

Ethics approval

The experiment was approved by the Ethics Committee of Affiliated Hospital of Yanbian University (Yanji, China) and was performed in accordance with its guidelines.

Statistical analysis

The data from these experiments were reported as the mean \pm standard error for each group. All statistical analyses were performed using Prism version 7.0 (GraphPad Software, Inc., La Jolla, CA, USA). Student's *t*-test was used to analyze the differences between two groups. Intergroup differences were analyzed by one-way analysis of variance, followed by a post hoc Tukey test for multiple comparisons. *P* < 0.05 was considered to indicate a statistically significant difference.

Results

Effect of SchB on body weight and uterus index

Increased body weight and lipid metabolism disorders are usually accompanied by estrogen deficiency in the experimental rodent model [22]. Consistent with previous results, our findings also demonstrated that body weight was significantly increased in OVX-operated rats compared with that of rats in the sham-operated group during the experimental period, while EV administration significantly reversed the OVX-induced increase in body weight in rats. However, both low and high concentrations of SchB had no obvious effect on body weight in OVX rats (Table I). As expected, atrophy of the uterus was observed in OVX rats, while EV treatment markedly accelerated the growth of the uterus in OVX rats. There was no

Table I. Body weight of rats was recorded every two weeks during the experimental period

Week	Sham	OVX	EV	SchB-L	SchB-H
0	205.6 \pm 7.8	209.3 \pm 8.2	212.4 \pm 7.6	208.7 \pm 6.9	211.4 \pm 7.5
2	214.4 \pm 9.2	235.8 \pm 10.1	224.3 \pm 9.1	231.2 \pm 9.3	234.5 \pm 8.2
4	233.5 \pm 10.7	256.5 \pm 12.6*	231.3 \pm 12.1 [#]	258.4 \pm 11.5	269.1 \pm 14.6
6	246.1 \pm 12.4	283.4 \pm 12.5**	253.1 \pm 13.6 ^{##}	279.6 \pm 11.6	269.1 \pm 12.5
8	258.3 \pm 14.2	311.2 \pm 15.2***	272.4 \pm 13.6 ^{###}	304.8 \pm 13.7	298.4 \pm 14.2

P* < 0.05, *p* < 0.01, ****p* < 0.001 compared with sham-operated rats in the same week; [#]*p* < 0.05, ^{##}*p* < 0.01, ^{###}*p* < 0.001 compared with OVX-operated rats in the same week.




Figure 1. Effect of SchB on uterus index. Uterus index was represented as the ratio of uterus weight to body weight. $N = 12$ in each group

* $P < 0.05$ vs. sham-operated group, # $p < 0.05$ vs. OVX-operated group.

significant improvement in the uterus index in OVX rats treated with SchB (Figure 1).

Effect of SchB on Ca metabolism and BMD

Ca is the most important mineral component in the skeletal system, and Ca intake can affect bone health by increasing BMD [23]. Estrogen deficiency in animal models induces osteoclastogenesis and bone resorption, which accelerate Ca loss by urinary excretion and release of skeletal Ca into the blood [24]. Our results demonstrated that serum Ca, serum P and tibial Ca were significantly decreased in OVX rats compared with those of rats in the sham-operated group. In contrast, urinary Ca was dramatically increased in OVX rats, suggesting that Ca loss was observed in estrogen-deficient rats. Interestingly, high-dosage SchB treatment alleviated OVX-induced Ca loss but had no effect on urinary P levels (Table II). To investigate whether SchB improved Ca homeostasis by regulating calciotropic hormone levels, three mainly calciotropic hormones – 25(OH)VD₃, PTH and calcitonin – were measured in the serum. The results indicated that 25(OH)VD₃ and calcitonin were significantly decreased and PTH was significantly elevated in OVX rats compared to rats

in the sham-operated group. Conversely, the OVX rats treated with SchB at a high dosage showed an increase in serum 25(OH)VD₃ and calcitonin and a decrease in serum PTH (Figures 2 A–C), reflecting that SchB might maintain Ca homeostasis in OVX rats by regulating calciotropic hormones. Furthermore, we explored the effect of SchB on femoral and tibial BMD in OVX rats. As shown in Figures 3 A and B, compared with sham-operated rats, the BMD of the femur and tibia significantly declined from 0.166 ± 0.008 and 0.148 ± 0.010 to 0.145 ± 0.006 and 0.125 ± 0.009 , respectively, in OVX rats 8 weeks after the surgical operation. However, the BMD of the femur and tibia in the OVX rats was increased and reached 0.158 ± 0.007 and 0.146 ± 0.007 , respectively, after high-dosage SchB treatment.

Effect of SchB on bone formation and bone resorption markers

The effects of SchB on the levels of bone formation markers, ALP and osteocalcin, and bone resorption markers, TRAP-5b and CTX, are shown in Figure 4. At 8 weeks after OVX surgical operation, the levels of ALP (Figure 4 A) and osteocalcin (Figure 4 B) in the serum were significantly decreased by 45.8% and 57.1%, respectively, and serum TRAP-5b (Figure 4 C) and CTX (Figure 4 D) were markedly elevated by 37.3% and 33.2%, respectively, in the OVX-operated group compared with the sham-operated group. SchB at a high dosage had a significant effect on increasing the activity of ALP by 44.0% and osteocalcin by 76.9% and suppressing the activity of TRAP-5b by 19.6% and CTX by 17.0%, compared to the OVX-operated group.



Effect of SchB on metabolism of trabecular bone and growth plate cartilage in the tibia

To determine the effects of SchB on trabecular bone and cartilage metabolism, TRAP, H&E and safranin O staining were performed in the tibia of OVX-operated rats with or without SchB treatment. TRAP staining showed that mature osteo-

Table II. Biochemical parameters in OVX mice treated with SchB

Parameter	Sham	OVX	EV	SchB-L	SchB-H
Serum Ca [mg/dl]	10.21 ±0.28	9.05 ±0.30*	10.35 ±0.36#	9.35 ±0.35	10.06 ±0.37#
Serum P [mg/dl]	5.79 ±0.34	5.04 ±0.30*	5.94 ±0.41#	5.17 ±0.31	5.83 ±0.35#
Urinary Ca/Cre [mg/mg]	0.167 ±0.018	0.331 ±0.035*	0.194 ±0.028#	0.301 ±0.037	0.213 ±0.026#
Urinary P/Cre [mg/mg]	1.22 ±0.15	1.31 ±0.26	1.35 ±0.38	1.23 ±0.37	1.18 ±0.39
Tibial Ca/ash [mg/mg]	0.452 ±0.008	0.402 ±0.009*	0.467 ±0.011#	0.408 ±0.012	0.459 ±0.015#
Tibial Ca [mg/g]	167.5 ±10.44	146.8 ±14.8*	169.6 ±12.33#	142.78 ±14.2	159.6 ±13.56#

SchB-L – schisandrin B with low concentration, SchB-H – schisandrin B with high concentration, Ca – calcium, P – phosphorus, Cre – creatinine, EV – estradiol valerate. * $P < 0.05$ versus sham-operated group; # $p < 0.05$ versus OVX group.



clast numbers were significantly increased in the proximal metaphysis of tibia from OVX-operated rats compared to the sham-operated group. However, treatment with SchB blocked osteoclastogenesis in the tibia of OVX-operated rats (Figures 5 A, B). In addition, H&E staining was performed to assess the trabecular bone structure in the proximal metaphysis of the tibia, and we observed a notable reduction in cancellous bone mass and network connectivity in OVX-operated rats, while high-dosage SchB treatment dramatically reversed the OVX-induced bone loss (Figure 6 A). Furthermore, safranin O staining revealed that the degradation of epiphyseal plate cartilage was de-

tected in the proximal tibia of OVX-operated rats, suggesting that cartilage formation was delayed in OVX-operated rats. Intriguingly, high-dosage SchB administration protected against OVX-induced cartilage degradation (Figure 6 B).

SchB activates PI3K/Akt signaling in the bone of OVX rats

Previous studies have indicated that impaired PI3K/Akt signaling was observed in the liver, skeletal muscle and tibia in an OVX-operated animal model [17, 25]. Our results also revealed that the protein expression of phosphorylated




Figure 4. Effect of SchB on bone formation and bone resorption markers. After treatment with SchB at low or high dosage for 8 weeks, the serum levels of alkaline phosphatase (ALP; **A**); osteocalcin (**B**); tartrate-resistant acid phosphatase 5b (TRAP-5b, **C**); C-terminal telopeptide of type 1 collagen (CXT, **D**) were measured by ELISA assays. *N* = 12 in each group

****p* < 0.001 vs. sham-operated group, ##*p* < 0.01, ###*p* < 0.001 vs. OVX-operated group.




Figure 5. Effect of SchB on osteoclast differentiation. After treatment with SchB at low or high dosage for 8 weeks, tartrate-resistant acid phosphatase (TRAP) staining was used to evaluate osteoclastogenesis (**A** and **B**; magnification 50×; scale bar = 100 μm). *N* = 12 in each group

**P* < 0.05 vs. sham-operated group; #*P* < 0.05 vs. OVX-operated group. Red arrows indicate mature osteoclast.

PI3K and Akt was inhibited in the proximal tibia of OVX-operated rats compared with that of rats in the sham-operated group. SchB treatment at high dosage markedly reversed the OVX-induced suppression of phosphorylated PI3K and Akt (Fig-

ure 7 A, B). β-catenin is a vital component of the wnt/β-catenin pathway and plays an important role in osteogenesis [26]. As expected, the protein expression of β-catenin was significantly decreased in the proximal tibia of OVX-operated




Figure 6. Effect SchB on metabolism of trabecular bone and growth plate cartilage. Hematoxylin and eosin (H&E; **A**; magnification 50 \times ; scale bar = 100 μ m) and safranin O staining (**B**; magnification 50 \times ; scale bar = 100 μ m) were performed to analyze trabecular bone and cartilage degeneration, respectively




Figure 7. SchB activated PI3K/Akt/ β -catenin signaling in bone of OVX rat. After treatment with SchB at low or high dosage for 8 weeks, protein expression of PI3K and p-PI3K (**A**), Akt and p-Akt (**B**), and β -catenin (**C**) was measured by western blotting. $N = 12$ in each group

*** $P < 0.001$ vs. sham-operated group; ** $p < 0.01$, ### $p < 0.001$ vs. OVX-operated group.

rats compared with the sham group. High-dosage SchB treatment increased β -catenin protein expression in OVX rats (Figure 7 C).

Discussion

The OVX rat, as an animal model of estrogen deficiency, has been widely used for mimicking PMO in women [27]. OVX-induced decline in ovarian estrogen production leads to a rapid loss of trabecular bone mass, which eventually results in an increased risk of fragility fracture [27]. Estrogens play a crucial role in bone metabolism, but it is still difficult to expound the precise mechanisms by which estrogens modulate Ca metabolism and bone homeostasis. Notably, the canonical osteoprotegerin (OPG)/receptor activator of the nuclear factor κ B ligand (RANKL)/RANK axis has been proposed as a key signaling pathway in mediating the molecular association between estrogen deficiency and bone deterioration [28]. In addition, reactive oxygen species and inflammatory responses potentiate bone degenerative processes in the presence of estrogen deficiency [29, 30]. In the present study, we aimed to determine whether SchB could attenuate bone deterioration caused by estrogen deficiency by regulating the PI3K/Akt pathway in an experimental rat model.

A previous study indicated that the PI3K/Akt pathway is a crucial factor for bone formation [31]. Inhibition of p-AKT *in vivo* and *in vitro* is accompanied by glucocorticoid-induced osteoporosis and osteoblast apoptosis [32, 33]. Xi *et al.* showed that the protein expression of phospho-PI3K and phospho-Akt is dramatically decreased in bone tissue from osteoporotic rats, and the PI3K/Akt pathway may be involved in the inhibition of osteoporosis through acceleration of osteoblast proliferation and bone formation [16]. In OVX mice [25, 34], activation of the PI3K/Akt pathway can promote bone formation and prevent bone resorption. In accordance with previous studies, the current results showed that OVX caused downregulation of phospho-PI3K and phospho-Akt in the proximal tibia. More importantly, SchB treatment showed a significant increase in phospho-PI3K and phospho-Akt in the proximal tibia of OVX-operated rats. We also found that estrogen withdrawal-induced calcium metabolism disequilibrium, trabecular bone loss, osteoclastogenesis and epiphyseal plate cartilage degradation were blocked by treatment with SchB, and the underlying mechanism was mediated, at least partially, through activation of the PI3K/Akt pathway in the local bone.

Our findings also revealed that SchB functioned as an activator of β -catenin protein expression in the proximal tibia of OVX-operated rats. Activation of Wnt signaling by inhibiting the activity of glycogen synthase kinase-3 β re-

duces the phosphorylation of β -catenin, which results in increased β -catenin in the cytosol and its translocation into the nucleus. Subsequently, the interaction of β -catenin with T cell factor/lymphoid enhancer binding factor enhances gene transcription to trigger osteoblast proliferation [26]. *In vivo* and *in vitro* studies have confirmed that β -catenin is a critical regulator of chondrocyte and osteoblast differentiation, the generation of ossification centers and perichondrial bone formation during the process of cartilage and bone development [33, 35]. In a previous study, icariin protected against glucocorticoid-induced osteoporosis by increasing β -catenin expression [33]. Moreover, improved bone mass and strength are correlated with activated Wnt/ β -catenin signaling in OVX mice [36]. Our results indicated that SchB could inhibit OVX-induced osteoporosis by enhancing β -catenin signaling. Interestingly, as a downstream target, β -catenin can be activated by phospho-PI3K and phospho-Akt, reflecting that the PI3K/Akt/ β -catenin signaling cascade is involved in osteogenic proliferation and differentiation and inhibits the development of osteoporosis [20, 33]. Our findings also provided a possible mechanism by which PI3K/Akt/ β -catenin signaling might participate in SchB-facilitated anti-osteoporotic activity in OVX-operated rats.

Usually, the majority of estrogen-like hormones, phytoestrogens and active components of Chinese traditional herbs are used as accessory therapeutic medicines for alleviating estrogen-related side effects in postmenopausal women undergoing anti-osteoporotic therapy [37]. Theoretically, the anti-osteoporotic activity of estradiol or estradiol valerate is obviously better than phytoestrogens or active pharmaceutical ingredients at equivalent dosage [38]. For example, subcutaneous administrations of genistein (10 mg/kg/day) shows similar effectiveness of 17 β -estradiol (2 μ g/kg/day) in bone protection [38]. Soy isoflavone (60 mg/kg/day) has positive effects on bone metabolism in OVX rats; the effective concentration of soy isoflavone is significantly higher than 17 β -estradiol in OVX mice [39]. Puerarin (≥ 2 mg/kg/day), a major isoflavonoid isolated from *Puerariae radix*, prevents bone loss in OVX mice, which is consistent with 17 β -estradiol (0.03 μ g/day) [40]. In the present study, our results demonstrated that high-dosage SchB treatment (50 mg/kg/day) had an equivalent anti-osteoporotic effect of EV treatment (1 mg/kg/day) in OVX rats.

Abundant Ca supplementation, at least 1000–1200 mg daily, has long been recommended for the prevention and treatment of osteoporosis [23]. Unfortunately, some side effects have been observed, including cardiovascular events, kidney stones and acute gastrointestinal symptoms [41–

43]. Therefore, there has been a growing number of publications focusing on the natural regulator of Ca metabolism. For example, ursolic acid, icariin and quercetin effectively improve calcium metabolism and bone mineral density [33, 44, 45]. Our results showed that SchB had a positive effect on Ca balance by modulating the activity of calcitropic hormones, including $25(\text{OH})_2\text{D}_3$, PTH and calcitonin.

However, there are some limitations to our study. The quantitative biological parameters of trabecular bone were not measured by micro-CT analysis. Moreover, the underlying molecular mechanisms of the effect of SchB on Ca metabolism are still not clear in our study.

In conclusion, our results demonstrated that SchB exerted anti-osteoporotic activity in OVX-operated rats, as shown by the improvement of Ca homeostasis and trabecular bone and cartilage metabolism, and the inhibition of osteoclast activity. The studies of the underlying molecular mechanism revealed that SchB accelerated the phosphorylation of PI3K and Akt and subsequently up-regulated the expression of β -catenin in OVX-operated rats, suggesting that the activated PI3K/Akt/ β -catenin signaling pathway might have a beneficial effect on OVX-induced osteoporosis.

Acknowledgments

Yimin Liang and Wei Li contributed equally to our work.

This study was supported by the Key Project of Scientific Research Foundation of the Education Department of Sichuan Province, China (Grant. No: 14ZA0185).

Conflict of interest

The authors declare no conflict of interest.

References

- Burge R, Dawson-Hughes B, Solomon DH, Wong JB, King A, Tosteson A. Incidence and economic burden of osteoporosis-related fractures in the United States, 2005-2025. *J Bone Miner Res* 2007; 22: 465-75.
- Lane NE, Haupt D, Kimmel DB, Modin G, Kinney JH. Early estrogen replacement therapy reverses the rapid loss of trabecular bone volume and prevents further deterioration of connectivity in the rat. *J Bone Miner Res* 1999; 14: 206-14.
- Ma X, Liu J, Yang L, Zhang B, Dong Y, Zhao Q. *Cynomorium songaricum* prevents bone resorption in ovariectomized rats through RANKL/RANK/TRAF6 mediated suppression of PI3K/AKT and NF-kappaB pathways. *Life Sci* 2018; 209: 140-8.
- Kanis JA, McCloskey EV, Harvey NC, Johansson H, Leslie WD. Intervention thresholds and the diagnosis of osteoporosis. *J Bone Miner Res* 2015; 30: 1747-53.
- Szeliga A, Maciejewska-Jeske M, Meczekalski B. Bone health and evaluation of bone mineral density in patients with premature ovarian insufficiency. *Prz Menopauz* 2018; 17: 112-6.
- Rossouw JE, Anderson GL, Prentice RL, et al. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results From the Women's Health Initiative randomized controlled trial. *JAMA* 2002; 288: 321-33.
- Dong XL, Cao SS, Gao QG, Feng HT, Wong MS, Denney L. Combination treatment with *Fructus Ligustri Lucidi* and *Puerariae radix* offsets their independent actions on bone and mineral metabolism in ovariectomized rats. *Menopause* 2014; 21: 286-94.
- Zhang Y, Li Q, Wan HY, et al. Study of the mechanisms by which *Sambucus williamsii* HANCE extract exert protective effects against ovariectomy-induced osteoporosis in vivo. *Osteoporos Int* 2011; 22: 703-9.
- Liu H, Xiong Y, Wang H, et al. Effects of water extract from epimedium on neuropeptide signaling in an ovariectomized osteoporosis rat model. *J Ethnopharmacol* 2018; 221: 126-36.
- Hwang YH, Kang KY, Kim JJ, Lee SJ, Son YJ, Paik SH. Effects of hot water extracts from polygonum multiflorum on ovariectomy induced osteopenia in mice. *Evid Based Complement Alternat Med* 2016; 2016: 8970585.
- Lai Q, Luo Z, Wu C, et al. Attenuation of cyclosporine A induced nephrotoxicity by schisandrin B through suppression of oxidative stress, apoptosis and autophagy. *Int Immunopharmacol* 2017; 52: 15-23.
- Xu J, Lu C, Liu Z, Zhang P, Guo H, Wang T. Schizandrin B protects LPS-induced sepsis via TLR4/NF-kappaB/MyD88 signaling pathway. *Am J Transl Res* 2018; 10: 1155-63.
- Thandavarayan RA, Giridharan VV, Arumugam S, et al. Schisandrin B prevents doxorubicin induced cardiac dysfunction by modulation of DNA damage, oxidative stress and inflammation through inhibition of MAPK/p53 signaling. *PLoS One* 2015; 10: e0119214.
- Ran J, Ma C, Xu K, et al. Schisandrin B ameliorated chondrocytes inflammation and osteoarthritis via suppression of NF-kappaB and MAPK signal pathways. *Drug Des Devel Ther* 2018; 12: 1195-204.
- Caichompoo W, Zhang QY, Hou TT, Gao HJ, Qin LP, Zhou XJ. Optimization of extraction and purification of active fractions from *Schisandra chinensis* (Turcz.) and its osteoblastic proliferation stimulating activity. *Phytother Res* 2009; 23: 289-92.
- Xi JC, Zang HY, Guo LX, et al. The PI3K/AKT cell signaling pathway is involved in regulation of osteoporosis. *J Recept Signal Transduct Res* 2015; 35: 640-5.
- Wang Y, Li B, Zhang W, et al. Impaired PI3 K Akt expression in liver and skeletal muscle of ovariectomized rats. *Endocrine* 2013; 44: 659-65.
- Wu CM, Chen PC, Li TM, Fong YC, Tang CH. Si-Wu-tang extract stimulates bone formation through PI3K/Akt/NF-kappaB signaling pathways in osteoblasts. *BMC Complement Altern Med* 2013; 13: 277.
- Jin X, Sun J, Yu B, et al. Daidzein stimulates osteogenesis facilitating proliferation, differentiation, and antiapoptosis in human osteoblast-like MG-63 cells via estrogen receptor-dependent MEK/ERK and PI3K/Akt activation. *Nutr Res* 2017; 42: 20-30.
- Wu X, Li S, Xue P, Li Y. Liraglutide, a glucagon-like peptide-1 receptor agonist, facilitates osteogenic proliferation and differentiation in MC3T3-E1 cells through phosphoinositide 3-kinase (PI3K)/protein kinase B (AKT), extracellular signal-related kinase (ERK)1/2, and cAMP/protein kinase A (PKA) signaling pathways involving beta-catenin. *Exp Cell Res* 2017; 360: 281-91.

21. Yu FY, Xie CQ, Sun JT, Peng W, Huang XW. Overexpressed miR-145 inhibits osteoclastogenesis in RANKL-induced bone marrow-derived macrophages and ovariectomized mice by regulation of Smad3. *Life Sci* 2018; 202: 11-20.
22. Li XL, Wang L, Bi XL, Chen BB, Zhang Y. Gushukang exerts osteoprotective effects by regulating vitamin D and calcium metabolism in ovariectomized mice. *J Bone Miner Metab* 2019; 37: 224-34.
23. Tai V, Leung W, Grey A, Reid IR, Bolland MJ. Calcium intake and bone mineral density: systematic review and meta-analysis. *BMJ* 2015; 351: h4183.
24. Morris HA, O'Loughlin PD, Anderson PH. Experimental evidence for the effects of calcium and vitamin D on bone: a review. *Nutrients* 2010; 2: 1026-35.
25. Liu J, Zhang Z, Guo Q, Dong Y, Zhao Q, Ma X. Syringin prevents bone loss in ovariectomized mice via TRAF6 mediated inhibition of NF-kappaB and stimulation of PI3K/AKT. *Phytomedicine* 2018; 42: 43-50.
26. Li L, Peng X, Qin Y, et al. Acceleration of bone regeneration by activating Wnt/beta-catenin signalling pathway via lithium released from lithium chloride/calcium phosphate cement in osteoporosis. *Sci Rep* 2017; 7: 45204.
27. Chen L, Yang L, Yao M, et al. Biomechanical characteristics of osteoporotic fracture healing in ovariectomized rats: a systematic review. *PLoS One* 2016; 11: e0153120.
28. Boyce BF, Xing L. Functions of RANKL/RANK/OPG in bone modeling and remodeling. *Arch Biochem Biophys* 2008; 473: 139-46.
29. Muthusami S, Ramachandran I, Muthusamy B, et al. Ovariectomy induces oxidative stress and impairs bone antioxidant system in adult rats. *Clin Chim Acta* 2005; 360: 81-6.
30. Weitzmann MN, Pacifici R. Estrogen deficiency and bone loss: an inflammatory tale. *J Clin Invest* 2006; 116: 1186-94.
31. Zhang Y, Zeng X, Zhang L, Zheng X. Stimulatory effect of puerarin on bone formation through activation of PI3K/Akt pathway in rat calvaria osteoblasts. *Planta Med* 2007; 73: 341-7.
32. Lu SY, Wang CY, Jin Y, et al. The osteogenesis-promoting effects of alpha-lipoic acid against glucocorticoid-induced osteoporosis through the NOX4, NF-kappaB, JNK and PI3K/AKT pathways. *Sci Rep* 2017; 7: 3331.
33. Hu J, Mao Z, He S, et al. Icariin protects against glucocorticoid induced osteoporosis, increases the expression of the bone enhancer DEC1 and modulates the PI3K/Akt/GSK3beta/beta-catenin integrated signaling pathway. *Biochem Pharmacol* 2017; 136: 109-21.
34. Xu X, Zhang Z, Wang W, Yao H, Ma X. Therapeutic effect of cistanoside a on bone metabolism of ovariectomized mice. *Molecules* 2017; 22: 197.
35. Dao DY, Jonason JH, Zhang Y, et al. Cartilage-specific beta-catenin signaling regulates chondrocyte maturation, generation of ossification centers, and perichondrial bone formation during skeletal development. *J Bone Miner Res* 2012; 27: 1680-94.
36. Zahoor M, Cha PH, Min do S, Choi KY. Indirubin-3'-oxime reverses bone loss in ovariectomized and hindlimb-unloaded mice via activation of the Wnt/beta-catenin signaling. *J Bone Miner Res* 2014; 29: 1196-205.
37. Wang T, Liu Q, Tjhi W, et al. Therapeutic potential and outlook of alternative medicine for osteoporosis. *Curr Drug Targets* 2017; 18: 1051-68.
38. Hertrampf T, Schleipen B, Offermanns C, Velders M, Laudenbach U, Diel P. Comparison of the bone protective effects of an isoflavone-rich diet with dietary and subcutaneous administrations of genistein in ovariectomized rats. *Toxicol Lett* 2009; 184: 198-203.
39. Kim DW, Yoo KY, Lee YB, et al. Soy isoflavones mitigate long-term femoral and lumbar vertebral bone loss in middle-aged ovariectomized mice. *J Med Food* 2009; 12: 536-41.
40. Yuan SY, Sheng T, Liu LQ, et al. Puerarin prevents bone loss in ovariectomized mice and inhibits osteoclast formation in vitro. *Chin J Nat Med* 2016; 14: 265-9.
41. Bolland MJ, Grey A, Avenell A, Gamble GD, Reid IR. Calcium supplements with or without vitamin D and risk of cardiovascular events: reanalysis of the Women's Health Initiative limited access dataset and meta-analysis. *BMJ* 2011; 342: d2040.
42. Jackson RD, LaCroix AZ, Gass M, et al. Calcium plus vitamin D supplementation and the risk of fractures. *N Engl J Med* 2006; 354: 669-83.
43. Lewis JR, Zhu K, Prince RL. Adverse events from calcium supplementation: relationship to errors in myocardial infarction self-reporting in randomized controlled trials of calcium supplementation. *J Bone Miner Res* 2012; 27: 719-22.
44. Lee SU, Park SJ, Kwak HB, Oh J, Min YK, Kim SH. Anabolic activity of ursolic acid in bone: Stimulating osteoblast differentiation in vitro and inducing new bone formation in vivo. *Pharmacol Res* 2008; 58: 290-6.
45. Derakhshanian H, Djalali M, Djazayeri A, et al. Quercetin prevents experimental glucocorticoid-induced osteoporosis: a comparative study with alendronate. *Can J Physiol Pharmacol* 2013; 91: 380-5.