

# An experimental study of buckwheat polysaccharide in adjuvant therapy for S180 sarcoma mice

Duo-Jiao Fan<sup>1</sup>, Peng Sun<sup>2</sup>, Shu-Ying Han<sup>3</sup>

<sup>1</sup>Department of Rehabilitation Medicine, Baoding Second Hospital, Baoding, China

<sup>2</sup>Department of Neurosurgery, Baoding Second Hospital, Baoding, China

<sup>3</sup>Department of Pharmacology, North China University of Science and Technology, Tangshan, Hebei, China

**Submitted:** 23 May 2019

**Accepted:** 24 December 2019

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms.2020.96882>

Copyright © 2020 Termedia & Banach

**Corresponding author:**

Shu-Ying Han

Department of Pharmacology

North China University of

Science and Technology

Tangshan, Hebei, China

E-mail: shuying\_

drhan2143@163.com

## Abstract

**Introduction:** To discuss the auxiliary therapeutic effect of buckwheat polysaccharide (BP) on S180 sarcoma.

**Material and methods:** Buckwheat polysaccharide was extracted with water and precipitated with ethanol. Solid tumor and ascites tumor mice models were established. The mice in the high, medium and low dosing groups ( $n = 24$ , each group) had their stomachs filled with different doses of BP. The cyclophosphamide (CTX) group and the model group ( $n = 24$ , each group) were used as control groups. The influence on the life cycle, the rate of suppressing the tumor, the thymus index, and the spleen index were evaluated. Tumor cells were cultured in vitro and intervened with drugs; flow cytometry was used to detect the changes in the cell cycle.

**Results:** Buckwheat polysaccharide significantly improved the lifespan and survival rate of the mice. The group of mice treated with the medium dose showed the best survival rate when compared to the ones that received high and low doses of BP ( $p < 0.01$ ). The tumor cells cultured in vitro were arrested in the G0/G1 phase to some extent ( $p > 0.05$ ). The cyclophosphamide arrested the cycle of the tumor cells in the G2/M period ( $p < 0.01$ ). Buckwheat polysaccharide could increase the thymus index, spleen index and the rate of suppressing the tumor, but the differences were not statistically significant.

**Conclusions:** Buckwheat polysaccharide had no obvious effect in inhibiting the growth of tumors, but it significantly extended the lifespan, increased the survival rate and reduced the toxic effect of CTX.

**Key words:** buckwheat polysaccharide, lifespan, thymus index, spleen index, cell cycle.

## Introduction

Malignant tumors have recently become one of the main diseases threatening the survival of human beings. In China, there are more than 1.6 million new cancer patients each year and it has become a major cause of death [1]. The main therapeutic function of antineoplastic drugs is to prolong and improve the quality of life of cancer patients; in this respect, traditional Chinese medicine has advantages. Buckwheat has the benefit of reducing blood pressure, reducing blood glucose and reducing lipids [2–5], and also has antineoplastic effects [6–9]. Buckwheat polysaccharide (BP) is a kind of polysaccharide extracted from the flowers

and leaves of buckwheat. An earlier study indicated that it was able to increase the immunologic function of experimental animals. The current study evaluated the adjuvant therapeutic effect of BP on S180 sarcoma mice.

## Material and methods

### Cell culture and animal model

S180 tumor cells were provided by the School of Preventive Medicine, North China Coal Medical University and maintained in a RPMI-1640 medium supplemented with 10% fetal bovine serum. The culture transfer cycle is 6 days. The S180 tumor-bearing model was established as previously described [10]. S180 mouse sarcoma cells were briefly diluted to a final concentration of  $1 \times 10^6$ /ml and inoculated subcutaneously into the right armpit or the abdominal cavity of healthy Kunming mice. The male-to-female ratio was 1 : 1. The weight was  $19 \pm 3$  g. The mice were provided by the Institute of Laboratory Animals Science, CAMS & PUMC. The qualified certificate number is No. SCXK (J) 2005-0013. The mice were fed in the barrier environment of the animal center of North China Coal Medical University. All animal experimental procedures were approved by the Animal Care and Use Committee of North China Coal Medical University and performed in accordance with the Guide for the Care and Use of Laboratory Animals [11].

### Main instruments

Electronic balance (Shanghai Second Balance Instrument Plant), automatic dual pure water distiller SZ-93 (Sichuan Yaxi Machinery Co., Ltd.), carbon dioxide incubator (Forma Scientific Company), inverted phase-contrast microscope UFX-ILA (OLYMPUS), clean bench (Forma Scientific Company), TGL-16G desk centrifuge (Shanghai Anting Scientific Instrument Plant) and flow cytometer (Becton-Dickinson Company).

### Drugs and reagents

Buckwheat polysaccharide: extracted by the pharmacologic teaching and research office of the North China Coal Medical University. Cyclophosphamide (CTX, Shang Hualian Pharmaceutical Co., Ltd., batch No.: 080612), PI, RNA enzyme and cell permeability agent (Beijing Solarbio Life Sciences Co., Ltd.).

### Extraction method of buckwheat polysaccharide

Distilled water (5000 ml) was poured into the flowers and leaves of buckwheat (500 g) (volume/weight = 10 : 1). Then, it was boiled at 95°C twice

(2 h and 1 h, respectively), and then filtered. After that, the filtrate was concentrated to 1/3 of the original solution. 95% ethanol was added and the crude polysaccharide was produced after sedimentation. Distilled water was added to the crude polysaccharide. 3% trichloroacetic acid was added at a ratio of 1 : 1 to remove protein. It was then left to stand for 24 h to obtain the liquid supernatant by centrifugation. Again, 95% ethanol was added for precipitation. Holosaccharide was then obtained after centrifugation and sediment drying.

### Suppression of buckwheat polysaccharide on S180 sarcoma mice

Ascitic fluid was taken from S180 tumor-bearing mice that received tumor cell transplantation and culture for 6 days. The ascitic fluid was washed with normal saline twice, counted, and the number of cells was adjusted to  $1 \times 10^8$ /ml. The fluid was subcutaneously injected into the mice's right axilla in accordance with the dose of 0.2 ml per mouse. The mice were randomly divided into six groups after inoculation: a high-dose BP group (BP-H, 400 mg/kg), a medium-concentration group (BP-M, 200 mg/kg), a low-concentration group (BP-L, 100 mg/kg), a CTX (150 µg/ml) positive control group, a BP-M+CTX (200 mg/kg + 150 µg/ml) group and a model control group. The number of mice in each group was 24. There was no significant difference in the baseline body mass among the groups. After inoculation, the BP was re-dissolved in 0.9% saline to the designated doses and intragastrically administered to the mice. Intraperitoneal injections were carried out for the mice in the CTX group. The mice in the model group received a *p.o.* administration of 0.9% saline (25 ml/kg). After successive administration for 14 days, the mice were weighed on the second day of the final administration, and executed by cervical vertebra dislocation. The tumor of each mouse was removed. The tumors were weighed, after the blood had been blotted up by filter paper, to calculate the anti-tumor rate.

Anti-tumor rate/% =  $(1 - \text{average tumor mass of the experimental groups} / \text{average tumor mass of the control groups}) \times 100\%$ .

### Effects of buckwheat polysaccharide on the thymus index and spleen index of S180 tumor-bearing mice

The thymus and spleen of the above mice were taken and weighed, after the blood on the surface was blotted up by filter paper, to calculate the thymus index and spleen index.

Thymus (spleen) index =  $\text{thymus (spleen) mass (mg)} / \text{weight (g)} \times 100\%$ .

### Effects of BP on the lifespan and survival rate of S180 mice with solid tumor(s)

Mouse tumor inoculation, administration and grouping were the same as above. There were 24 mice in each group. The first day was the same day of administration. The experimental time was 25 days. The general conditions and time of death of the tumor-bearing mice were recorded to calculate the average lifetime and survival rate after 25 days.

Lifetime = (the sum of the survival time of the animals in the experimental group/the number of the animals in the experimental group) × 100%.

Survival rate after 25 days = (the number of animals that survived after 25 days/the number of the animals in the experimental group) × 100%.

### Effects on lifespan and survival rate of S180 ascitic tumor mice

Mouse tumor inoculation, administration and grouping are the same as above. There were 24 mice in each group. The first day was the same day of administration. The experimental time was 25 days. The general conditions and time of death of the tumor-bearing mice were recorded to calculate the average lifetime and survival rate after 25 days.

Average lifespan = (the sum of the survival time of the animals in the experimental group/the number of the animals in the experimental group) × 100%.

Survival rate after 25 days = (the number of animals that survived after 25 days/the number of the animals in the experimental group) × 100%.

### Cell apoptosis detection via flow cytometry

Ascitic fluid from the tumor cells of the S180 mice that had been inoculated for three days was drawn, and the fluid was washed with Hanks solution twice to adjust the cell concentration to 10<sup>6</sup>/ml. The culture medium with cells was placed on a 24-well culture plate. These cells were divided into 6 groups with 6 duplicate wells, each well containing 0.6 mL of solution. Different drugs were then added to the culture wells in each group: the model group (with appropriate culture medium added); the high, medium and lower concentration groups (the BP solution concentration in the culture solution was 150 µg/ml and 75 µg/ml, respectively); the CTX group (the final concentration of CTX in the culture solution was 150 µg/ml), the CTX (150 µg/ml) + BP-M (150 µg/ml) group. It was ensured that the final volume of the groups was the same. After 24 h, the cells were collected into sterile centrifuge tubes. The liquid supernatant was removed by centrifugation twice. Then, 70% cold ethanol was added and the solution was placed in the refrigerator at -20°C. During analysis, the stored cell sap was taken to be washed with phosphate buffer saline (PBS) twice. After that, the liquid supernatant

was removed by centrifugation, and the BPS solution was re-suspended. The prepared propidium iodide (PI) dying liquor and RNA enzyme were added to produce a reaction by keeping it in the dark for 15 min. The solution was filtered by a 400-mesh net and then analyzed by computer.

### Statistical analysis

The database of the experimental data was established in Excel. Statistical analysis was carried out using SPSS 13.0 software, and the pairwise comparison of the multiple samples conducted using one-way analysis of variance (ANOVA).

### Results

#### Effects of buckwheat polysaccharide on lifespan and survival rate of S180 mice with solid tumor(s)

After the mice were inoculated with S180 sarcoma cells, the right fore armpits of the mice were enlarged after about 6 days, and the right fore armpit of the mice grew larger with time. In the model control group and the CTX group, the mice gradually reduced their activity and diet. The luster of the mice's fur became dull and the mice were slow to respond. All the above conditions gradually became more serious. These conditions were rather more severe in the CTX group. The activity, diet, fur luster and mental conditions of the mice in the group with different BP doses were better than those of the mice in the CTX group and the model control group. From the 9<sup>th</sup> day, the mice in the model group and the CTX group began to die. The results during the 25-day observation period are shown in Figure 1 and Table I.

#### Effects of buckwheat polysaccharide on lifespan and survival rate of S180 ascitic tumor mice

For the mice with S180 ascitic fluid sarcoma strains inoculated in the abdominal cavity, their




Figure 1. Effect of buckwheat polysaccharide (BP) on the lifespan of S180 solid tumor mice (n = 24)

**Table I.** Effect of buckwheat polysaccharide (BP) on lifespan of  $S_{180}$  solid tumor mice ( $x \pm s, n = 24$ )


Group	Dose ( $\text{mg} \cdot \text{kg}^{-1}$ )	Lifespan (d)	Survival rate (%)
Model	–	25.29 $\pm$ 8.22 <sup>d</sup>	25.00
BP-H	400	29.29 $\pm$ 8.14	54.17
BP-M	200	33.42 $\pm$ 4.61 <sup>b</sup>	62.50
BP-L	100	29.38 $\pm$ 7.80	45.83
CTX	3	32.46 $\pm$ 5.76 <sup>b</sup>	66.67
CTX + BP-M	3 + 200	33.29 $\pm$ 8.53 <sup>b</sup>	70.83

<sup>a</sup> $p < 0.05$ , <sup>b</sup> $p < 0.01$  vs. model group; <sup>c</sup> $p < 0.05$ , <sup>d</sup> $p < 0.01$  vs CTX group.

abdomen began to enlarge on the 5<sup>th</sup> day, and the abdomen girth and size increased with time. The activity, diet and mental conditions of the mice in the group with different BP doses were better than those of the mice in the CTX group and the model group. From the 6<sup>th</sup> day, the mice in the CTX group began to die. The results during the 25-day observation period are shown in Figure 2 and Table II.

#### Effects of buckwheat polysaccharide on the thymus and spleen indexes of S180 sarcoma mice

The thymus and spleen indexes of the mice in the group with different doses were higher than those of the mice in the model group; however, the differences were not statistically significant.



**Figure 2.** Effect of buckwheat polysaccharide (BP) on the lifespan of S180 ascites tumor mice ( $n = 24$ )

**Table II.** Effect of buckwheat polysaccharide (BP) on lifespan of  $S_{180}$  ascites tumor mice ( $x \pm s, n = 24$ )


Group	Dose ( $\text{mg} \cdot \text{kg}^{-1}$ )	Lifespan (d)	Survival rate (%)
Model	–	18.87 $\pm$ 6.10 <sup>d</sup>	20.83
BP-H	400	22.00 $\pm$ 5.54	50.00
BP-M	200	23.25 $\pm$ 4.67 <sup>b</sup>	58.33
BP-L	100	20.46 $\pm$ 6.41 <sup>c</sup>	45.83
CTX	3	23.67 $\pm$ 4.66 <sup>b</sup>	62.50
CTX+BP-M	3 + 200	24.08 $\pm$ 5.58 <sup>b</sup>	66.67

<sup>a</sup> $p < 0.05$ , <sup>b</sup> $p < 0.01$  vs. model group; <sup>c</sup> $p < 0.05$ , <sup>d</sup> $p < 0.01$  vs. CTX group.

The thymus index and spleen index of the mice in the CTX group were significantly reduced, while BP-M successfully controlled the lowering of the thymus index and spleen index caused by CTX (Figure 3).

#### Observation of BP for the inhibition rate of mice S180 sarcoma

As for the weights of the tumors in each group, the tumors in the model group were the most severe. The weight of the tumors in the CTX and CTX+BP groups was significantly lower than that in the model group ( $p < 0.01$ ). High, medium and low doses of BP had the ability to suppress the growth of mouse S180 sarcoma. However, the dif-



**Figure 3.** The effect of EBFL on thymus and spleen index

<sup>a</sup> $p < 0.05$ , <sup>b</sup> $p < 0.01$  vs. model group; <sup>c</sup> $p < 0.05$ , <sup>d</sup> $p < 0.01$  vs. CTX group

**Table III.** Effect of buckwheat polysaccharide (BP) on inhibiting S<sub>180</sub> tumor in SA mice ( $\bar{x} \pm s$ ,  $n = 24$ )

Group	Dose (mg · kg <sup>-1</sup> )	Tumor weight (g)	Inhibition rate (%)
Model		3.28 ± 0.46 <sup>d</sup>	
BP-H	400	3.00 ± 0.40 <sup>d</sup>	9.04 ± 1.55
BP-M	200	2.98 ± 0.45 <sup>d</sup>	9.27 ± 2.21
BP-L	100	3.04 ± 0.36 <sup>d</sup>	7.37 ± 1.71
CTX	3	1.91 ± 0.31 <sup>b</sup>	42.67 ± 8.61
CTX + BP-M	3 + 200	1.69 ± 0.38 <sup>b</sup>	46.67 ± 9.71

<sup>a</sup> $p < 0.05$ , <sup>b</sup> $p < 0.01$  vs. model group; <sup>c</sup> $p < 0.05$ , <sup>d</sup> $p < 0.01$  vs. CTX group.

ference compared with the model group was not statistically significant ( $p > 0.05$ ). The difference in weight of the tumors in the CTX and CTX + BP-M groups was not statistically significant. The anti-tumor rate was significantly higher than in the BP-H, BP-M and BP-L groups. The detailed results are shown in Table III.

#### Cell cycle detection via flow cytometry

In the G0/G1 phase, the cells of the tumor cells in the group with different doses of BP increased, with the BP-M group having the most obvious increase. However, the difference compared with the model group was not statistically significant ( $p > 0.05$ ). It indicated that BP was able to arrest the cycle of the tumor cells in the G0/G1 phase. The cell content of the tumor cells in the CTX group greatly increased in the G2/M phase compared with the model group ( $p < 0.01$ ), while the content in the G0/G1 phase was significantly lower than the model group ( $p < 0.05$ ), which indicated that the drug mainly arrested the tumor cells in the G2/M phase. Details are shown in Figure 4 and Table IV.

#### Discussion


The types of drugs available for the treatment of malignant tumors are varied [12, 13]. Chemotherapeutic drugs such as CTX, etc., can suppress the growth of tumors, but the toxic side effects remarkably lower the quality of the patients' lives. Traditional Chinese medicine has gradually become a research hotspot for its advantages of high efficiency, low toxicity, and overall adjustment [14, 15]. The study of immunocompetence polysaccharide in the aspects of promoting and adjusting immunologic function has progressed greatly, and the study of its mechanism of action has been deepened at the molecular and genetic levels [16–22]. At present, the mechanism for polysaccharide antineoplastic study has two main aspects: one is that the polysaccharide directly kills tumor cells; the second is that the polysaccharide achieves the indirect antineoplastic effect by activating or

enhancing immunologic function [23]. The documents related to this study indicated that, at the same time as enhancing the immunity, this polysaccharide has no direct toxic side effects on normal cells. Therefore, this polysaccharide has been recognized as a very important biological effect regulator [24–26].


The results of this experiment indicate that BP had a certain suppressing effect on mouse S180 sarcoma; however, the difference was not statistically significant. BP was able to improve the lifespan, survival rate and survival quality of S180 sarcoma mice. Its application alone was able to improve the thymus and spleen index of tumor mice, while its combined application with CTX was able to improve the suppressing effects of CTX on the thymus index and spleen index of tumor mice. A flow cytometer was used to observe the effects of BP on tumor cell cycle. It was found that BP was able to arrest the tumor cell cycle in the G0/G1 phase to prevent cell transformation. The experimental results indicated that BP did not have obvious antineoplastic effects, but it could prolong the lifespan of the tumor mice and improve the survival quality. In addition, it was able to prevent the suppressing and toxic reaction of CTX on immunologic function, which functions as adjuvant therapy for tumors.

There are some limitations to the present study. Firstly, the effect of BP in adjuvant therapy was solely investigated in S180 xenografts. Further studies on different tumor-bearing models will be needed to confirm the beneficial effect of BP in tumor progression. Secondly, we did not explore the molecular mechanisms of BP in extending the lifetime of tumor-bearing mice. Further investigation will be required to identify the potential regulatory targets of BP in tumor progression.

In conclusion, our results showed that BP significantly extended the lifespan, increased the survival rate and reduced the toxic effect of CTX in S180 tumor-bearing mice. The adjuvant therapeutic function of BP deserves further investigation to determine its role in the treatment of cancer.



**Figure 4.** The effect of buckwheat polysaccharide (BP) on the cell cycle of S180 ascites tumor mice ( $x \pm s$ )  
 6.001 = Control, 1.001 = BP-H, 2.001 = BP-M, 3.001 = BP-L



**Figure 4.** Cont. The effect of buckwheat polysaccharide (BP) on the cell cycle of S180 ascites tumor mice ( $x \pm s$ )  
 6.001 = Control, 1.001 = BP-H, 2.001 = BP-M, 3.001 = BP-L

**Table IV.** Effect of buckwheat polysaccharide (BP) on cell cycle of  $S_{180}$  ascites tumor mice ( $x \pm s$ )

Group	Dose ( $\mu\text{g/ml}$ )	G0/G1 (%)	S (%)	G2/M (%)
Model	–	44.16 $\pm$ 1.03	44.18 $\pm$ 2.70	11.66 $\pm$ 3.06
BP-H	300	45.19 $\pm$ 2.13 <sup>d</sup>	42.11 $\pm$ 3.27	12.59 $\pm$ 1.16 <sup>d</sup>
BP-M	150	47.30 $\pm$ 4.70 <sup>d</sup>	40.88 $\pm$ 4.54	11.84 $\pm$ 0.68 <sup>d</sup>
BP-L	75	46.26 $\pm$ 4.67 <sup>d</sup>	42.47 $\pm$ 3.98	11.27 $\pm$ 1.36 <sup>d</sup>
CTX	150	39.71 $\pm$ 2.49 <sup>a</sup>	42.72 $\pm$ 1.12	17.57 $\pm$ 2.48 <sup>b</sup>
CTX + BP-M	150 + 150	43.71 $\pm$ 3.18 <sup>c</sup>	42.93 $\pm$ 1.01	13.37 $\pm$ 2.85 <sup>d</sup>

<sup>a</sup> $p < 0.05$ , <sup>b</sup> $p < 0.01$  vs. model group; <sup>c</sup> $p < 0.05$ , <sup>d</sup> $p < 0.01$  vs. CTX group.

### Acknowledgments

Duo-Jiao Fan and Peng Sun contributed equally to this study as first authors.

### Conflict of interest

The authors declare no conflict of interest.

### References

- Wang ZL, Dong BW. [Research on the application of immune cells in immunotherapy of malignant tumors] [Article in Chinese]. Medical Recapitulate 2004; 11: 705-7.
- Hou Z, Hu Y, Yang X, Chen W. Antihypertensive effects of Tartary buckwheat flavonoids by improvement of vascular insulin sensitivity in spontaneously hypertensive rats. Food Funct 2017; 8: 4217-28.
- Mišan A, Petelin A, Stubelj M, et al. Buckwheat – enriched instant porridge improves lipid profile and reduces inflammation in participants with mild to moderate hypercholesterolemia. J Funct Foods 2017; 36: 186-94.
- Li J, Gong FY, Li FL. Hypoglycemic and hypolipidemic effects of flavonoids from tartary buckwheat in type 2 diabetic rats. Biomed Res 2016; 27: 132-7.
- Ji X, Han L, Liu F, Yin S, Peng Q, Wang M. A mini-review of isolation, chemical properties and bioactivities of polysaccharides from buckwheat (Fagopyrum Mill). Int J Biol Macromol 2019; 127: 204-9.
- Chen H, He LJ, Lin LF. [Advances in anti-tumor mechanism of buckwheat] [Article in Chinese]. Strait Pharmaceutical Journal 2014; 26: 41-2.
- Bai C, Feng M, Hao X, Zhao Z, Li Y, Wang Z. Anti-tumoral effects of a trypsin inhibitor derived from buckwheat in vitro and in vivo. Mol Med Rep 2015; 12: 1777-82.
- Zhou XL, Meng XX, Wang Q, Zhou YM, Li ZJ. Comparative anti-tumor activity study of tartary buckwheat flavonoids and amphibian peptides. Adv Mat Res 2013; 781-784: 1270-4.
- Guo X, Zhu K, Zhang H, Yao H. Anti-tumor activity of a novel protein obtained from tartary buckwheat. Int J Mol Sci 2010; 11: 5201-11.
- Liu F, Wang JG, Wang SY, Li Y, Wu YP, Xi SM. Antitumor effect and mechanism of Gecko on human esophageal carcinoma cell lines in vitro and xenografted sarcoma 180 in Kunming mice. World J Gastroenterol 2008; 14: 3990-6.
- National Research Council (US) Committee for the Update of the Guide for the Care and Use of Laboratory Animals. Guide for the Care and Use of Laboratory Animals. 8th Ed. Washington (DC): National Academies Press (US); 2011. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK54050/>; doi: 10.17226/12910.
- Long HY, Huang QX, Yu YY, et al. Dehydrocostus lactone inhibits in vitro gastrinoma cancer cell growth through

- apoptosis induction, sub-G1 cell cycle arrest, DNA damage and loss of mitochondrial membrane potential. *Arch Med Sci* 2019;15:765-773.
13. Zhang Z, Feng Y, Li Z, Cao X. Antiproliferative and apoptotic activity of glycyrrhizic acid in MCF-7 human breast cancer cells and evaluation of its effect on cell cycle, cell migration and m-TOR/PI3K/Akt signalling pathway. *Arch Med Sci* 2019; 15: 174-82.
  14. Nie J, Zhao C, Deng LI, et al. Efficacy of traditional Chinese medicine in treating cancer. *Biomed Rep* 2016; 4: 3-14.
  15. Chen Y, Yao F, Ming K, Wang D, Hu Y, Liu J. Polysaccharides from traditional chinese medicines: extraction, purification, modification, and biological activity. *Molecules* 2016; 21: 1705.
  16. Ruijun W, Shi W, Yijun X, Mengwuliji T, Lijuan Z, Yumin W. Antitumor effects and immune regulation activities of a purified polysaccharide extracted from *Juglan regia*. *Int J Biol Macromol* 2015; 72: 771-5.
  17. Wu GH, Lu CL, Jiang JG, Li ZY, Huang ZL. Regulation effect of polysaccharides from *Pleurotus tuber-regium* (Fr.) on the immune activity of mice macrophages. *Food Funct* 2014; 5: 337-44.
  18. Gu P, Liu Z, Sun Y, et al. *Angelica sinensis* polysaccharide encapsulated into PLGA nanoparticles as a vaccine delivery and adjuvant system for ovalbumin to promote immune responses. *Int J Pharm* 2019; 554: 72-80.
  19. Yin M, Zhang Y, Li H. Advances in research on immunoregulation of macrophages by plant polysaccharides. *Front Immunol* 2019; 10: 145.
  20. Sun J, Xiong J, Yao L, et al. The effect of dietary ginseng polysaccharide supplementation on porcine milk-derived esRNAs involved in the host immune responses. *J Anim Physiol Anim Nutr (Berl)* 2019; 103: 276-82.
  21. Wang YQ, Mao JB, Zhou MQ, et al. Polysaccharide from *Phellinus igniarius* activates TLR4-mediated signaling pathways in macrophages and shows immune adjuvant activity in mice. *Int J Biol Macromol* 2019; 123: 157-66.
  22. Bo R, Liu Z, Zhang J, et al. Mechanism of *Lycium barbarum* polysaccharides liposomes on activating murine dendritic cells. *Carbohydr Polym* 2019; 205: 540-9.
  23. Du JW, Yang MF, Su YY, Xia ZL. [Experimental study on anti-S180 sarcoma of *Grifola frondosa* polysaccharide in vivo] [Article in Chinese]. *Shandong Journal of Traditional Chinese Medicine* 2006; 26: 193-5.
  24. Furusawa E, Hirazumi A, Story S, Jensen J. Antitumour potential of a polysaccharide-rich substance from the fruit juice of *Morinda citrifolia* (Noni) on sarcoma 180 ascites tumour in mice. *Phytother Res* 2003; 17: 1158-64.
  25. Zheng M. [Advances in pharmacology of Polysaccharides from traditional Chinese medicine against tumors] [Article in Chinese]. *Foreign Medical Sciences* 2000; 23: 259-63.
  26. Bo R, Sun Y, Zhou S, et al. Simple nanoliposomes encapsulating *Lycium barbarum* polysaccharides as adjuvants improve humoral and cellular immunity in mice. *Int J Nanomedicine* 2017; 12: 6289-301.