

The use of prognostic scales in upper gastrointestinal tract bleeding.

Type

Research paper

Keywords

gastrointestinal bleeding, prognostic scales, AIMS 65, Glasgow-Blatchford, Rockall

Abstract

Introduction

In order to select high-risk patients, many prognostic scales have been invented. Among them, Rockall, Glasgow-Blatchford and AIMS 65 scales were considered the most useful.

Material and methods

Patients with upper GI tract bleeding, treated between 2017 and 2018 were retrospectively enrolled to the study. Every patient had a Rockall, Glasgow-Blatchford and AIMS 65 score calculated retrospectively. Data on hospitalization as blood transfusions, length of hospital stay, rebleeding, intensive care unit (ICU) admission, mortality was included into data-base.

Results

Univariate logistic regression revealed that only the AIMS65 scale was a prognostic factor for in-hospital mortality (OR 11.028; 95% CI: [2.271, 53.563], $p=0.001$). The AIMS 65 score >2 was the only factor predicting the need of >4 blood units transfusion during hospitalization (OR 3.977; 95% CI: [1.305, 12.122], $p=0.015$), whereas Glasgow-Blatchford scale >5 was the only risk factor for the need of fresh frozen plasma transfusion (OR 3.657; 95% CI: [1.010-13.242], $p=0.048$). The area under a curve (AUC) in the ROC analysis revealed that the AIMS 65 scale was the most accurate in mortality prediction (AUC=0.859, $p=0.002$), whereas Rockall score and Glasgow-Blatchford were not significant (AUC=0.614, $p=0.093$ and AUC=0.504, $p=0.97$, respectively).

Conclusions

Based on our results, we recommend using AIMS 65 scoring system. It's simple and requires few of parameters to be counted. Also, it proved to be the most efficient in predicting in-hospital mortality.

Abstract:

In order to select high-risk patients, many prognostic scales have been invented. Among them, Rockall, Glasgow-Blatchford and AIMS 65 scales were considered the most useful.

Materials and methods: Patients with non-variceal upper GI tract bleeding, treated between 2017 and 2018 were retrospectively enrolled to the study. Every patient had a Rockall, Glasgow-Blatchford and AIMS 65 score calculated retrospectively. Data on hospitalization as blood transfusions, length of hospital stay, rebleeding, intensive care unit (ICU) admission, mortality was included into data-base.

Results: Univariate logistic regression revealed that only the AIMS65 scale was a prognostic factor for in-hospital mortality (OR 11.028; 95% CI: [2.271, 53.563], $p=0.001$). The AIMS 65 score >2 was the only factor predicting the need of >4 blood units transfusion during hospitalization (OR 3.977; 95% CI: [1.305, 12.122], $p=0.015$), whereas Glasgow-Blatchford scale >5 was the only risk factor for the need of fresh frozen plasma transfusion (OR 3.657; 95% CI: [1.010-13.242], $p=0.048$). The area under a curve (AUC) in the ROC analysis revealed that the AIMS 65 scale was the most accurate in mortality prediction (AUC=0.859, $p=0.002$), whereas Rockall score and Glasgow-Blatchford were not significant (AUC=0.614, $p=0.093$ and AUC=0.504, $p=0.97$, respectively).

Conclusions: Based on our results, we recommend using AIMS 65 scoring system. It's simple and requires few of parameters to be counted. Also, it proved to be the most efficient in predicting in-hospital mortality.

Keywords: gastrointestinal bleeding, prognostic scales, AIMS 65, Glasgow-Blatchford, Rockall

1. Introduction

Despite improvement in the medical treatment of peptic ulcer disease through the use of proton pump inhibitors (PPIs), the incidence of upper gastrointestinal bleeding remains high. Estimates of the incidence of upper GI bleeding range between 50 to 150 cases per 100,000 citizens annually, depending on geographic localization [1]. Moreover, the incidence of patients requiring anticoagulant and anti-platelet medications due to cardiac or vascular comorbidity continues to rise [2,3]. These patients are also more commonly treated for concomitant diseases such as diabetes or chronic renal insufficiency, which may worsen the overall prognosis. The overall mortality related to upper GI bleeding varies between 2 - 2.5% [4,5].

In order to select high-risk patients, many prognostic scales have been invented. Among them, Rockall, Glasgow-Blatchford and AIMS 65 scales were considered the most useful. A Rockall score assesses patient age, symptoms of shock, concomitant diseases and endoscopic findings[4]. Whereas the Glasgow-Blatchford scale is based on hemodynamic parameters, hemoglobin count, symptoms of GI tract bleeding and presence of cardiac and hepatic diseases [6]. A recently introduced score called AIMS 65 takes into consideration albumin level, INR, blood pressure, mental status and age [7]. However, despite their value, none of these scales are widely used in clinical practice. There is a lack of data on which scale offers the most accurate risk stratification in patients with acute upper GI bleeding. Therefore, we designed a study comparing all three scales.

2. Aim of the study:

The aim of the study was to compare different scoring systems (Rockall score, Glasgow-Blatchford, AIMS 65) as a tool for mortality prediction in patients with non-variceal upper GI bleeding.

3. Materials and methods

Patients with non-variceal upper GI tract bleeding, treated between January 2017 and December 2018 in 2nd Department of General Surgery of the Jagiellonian University, were retrospectively enrolled to the study.

Patients presenting with symptoms of GI bleeding, such as hematemesis, melena, coffee ground vomiting or blood in the nasogastric tube, were considered for endoscopic examination. The Forrest scale was used to classify severity of bleeding [8]. In case of active bleeding, injection of adrenaline with 10% saline solution was used to achieve hemostasis. Forrest Ia bleeding and Forrest IIa ulcers were additionally secured by an endoclip. Every patient has had constant 8 mg/hour PPI infusion. All patients had basic biochemistry, CBC and vital parameters assessed. Blood transfusion was indicated when hemoglobin count dropped below 7g/dL, or the hematocrit level was below 21%, or signs of hemodynamic instability were present, despite fluid injection. Fresh Frozen Plasma was transfused in case of coagulation disorders (INR>1,3, APTT> 36s). Every patient had a clinical Rockall, Glasgow-Blatchford and AIMS 65 score calculated retrospectively. Data on hospitalization as blood transfusions, length of hospital stay, rebleeding, intensive care unit (ICU) admission, mortality was included into data-base. Primary endpoint was in-hospital mortality. Secondary endpoints were bleeding recurrence, need for blood and fresh frozen plasma transfusion.

Statistical analysis and ethical consideration

All data was analyzed with Statistica version 13.0 PL (StatSoft Inc., Tulsa, OK, USA). The results are presented as mean standard deviation (mean \pm SD), median, and interquartile range (IQR). The study of categorical variables used the chi-square test of independence. The Shapiro-Wilk test was used to check for normal distribution of data and the Student *t* test was used for normally distributed quantitative data. For non-normally distributed quantitative variables, the Mann-Whitney *U* test was used. A *p*-value < 0.05 was considered statistically

significant. All considerable patient- and treatment-related factors were analyzed in univariate logistic regression models, then significant factors were analyzed in the multiple logistic regression model in search of independent risk factors for blood and fresh frozen plasma transfusion and rebleeding or in-hospital mortality. The results are presented as Odds Ratio (OR) and Confidence Interval (CI). Receiver operating characteristic curves with area under curve (AUC) calculation were used to compare the scoring systems in terms of sensitivity and specificity. Youden index was used to select cut-off points.

All procedures were performed in accordance with the ethical standards of the 1964 Declaration of Helsinki and its later amendments (Fortaleza 2013). The study was approved by the Local Ethics Committee of the Jagiellonian University Medical College (KBN no 122.6120.36.2016).

3. Results

137 patients were included in the analysis. Mean age was 64.13 ± 16.04 .

The median AIMS65 score was 1 (IQR: 1-2). There were 15 patients (10.95%) with high risk of in-hospital mortality. 118 (86.13%) patients had Glasgow-Blatchford score >6 , which stratifies the risk of needing an intervention as $>50\%$. Median Glasgow-Blatchford score was 10 (IQR: 8-13). Median Rockall score was 5 (IQR: 3-6). There were 6 (4.38%) patients with high risk of mortality. The summary is included in Table 2. Six (4.38%) patients died during hospitalization. Univariate logistic regression revealed that only the AIMS65 scale was a prognostic factor for in-hospital mortality (OR 11.028; 95% CI: [2.271, 53.563], $p=0.001$). The results of univariate logistic regression are shown in Table 3. The area under a curve (AUC) in the ROC analysis revealed that the AIMS 65 scale was the most accurate in mortality prediction (AUC=0.859, $p=0.002$), whereas Rockall score and Glasgow-Blatchford were not significant (AUC=0.614, $p=0.093$ and AUC=0.504, $p=0.97$, respectively).

There were 11 patients with bleeding recurrence during their hospitalization. ROC analysis revealed that the AIMS 65 and Glasgow-Blatchford score were comparable in rebleeding prediction (AUC=0.735; p=0.004 and AUC=0.737; p=0.001 respectively). The Rockall score had lower accuracy (AUC=0.56; p=0.451).

102 patients required blood transfusions during hospitalization and 29 patients required more than 4 units of packed red blood cells. 51 patients required fresh frozen plasma transfusions. The data are summarized in table 1.

The AIMS 65 score >2 was the only factor predicting the need of >4 blood units transfusion during hospitalization (OR 3.977; 95% CI: [1.305, 12.122], p=0.015), whereas Glasgow-Blatchford scale >5 was the only risk factor for the need of fresh frozen plasma transfusion (OR 3.657; 95% CI: [1.010-13.242], p=0.048). The results are summarized in Table 4.

4. Discussion

This study has demonstrated the value of the AIMS65 scoring system, which proved to be most accurate in mortality prediction and need of blood transfusion in patients with upper GI tract bleeding.

The AIMS65 scoring system's main objective is to stratify patients with a high risk of in-hospital mortality [7]. In our study, it was the only scoring system that helped to accurately predict in-hospital death. It stands in line with study by Robertson et al. who showed superiority of this scoring system in comparison to Glasgow-Blatchford and Rockall scale [9]. Japanese study also revealed superiority of the AIMS65 score [10]. In addition to its high accuracy, AIMS 65 system is simple and based on basic parameters [11]. On the contrary, Choe et al. suggested that both Rockall and Glasgow-Blatchford scales are more accurate in predicting the need of blood transfusion and endoscopic intervention [12]. Stanley et al., using data from six hospitals in Europe, North America, Asia and Oceania showed superiority of AIMS 65 for mortality prediction, however authors still recommended Glasgow-Blatchford scale as most accurate in general risk stratification [13]. Shafagi et al. proposed modified Glasgow-Blatchford scoring system, taking albumin level into account, as more accurate in Central European population[14].

AIMS 65 and Glasgow-Blatchford scales were comparable in prediction of rebleeding. The same findings were revealed in a study by Rout [15]. However, as the author pointed out, the AIMS 65 scale was originally invented for risk stratification of non-variceal upper GI bleedings [16].

In our study, the AIMS 65 score was also superior to the others in predicting the need for blood transfusion. Conversely, Hyett et al. underlined the superiority of the Glasgow-Blatchford scale [11]. One possible explanation for this discrepancy is the frequent use of oral anticoagulants in central Europe, which elevates INR level, one of the components of

AIMS65 scale. However, in our study the Glasgow-Blatchford scale was the only one that helped with predicting the need for plasma transfusion.

Our study has some limitations, which may affect final results. At first, the study is designed as a retrospective analysis of data, therefore some of the important data might have been missed in the medical records. Secondly, the study group is middle-sized, which may have an impact on the final results. Lastly, we did not perform a multivariate-analysis of logistic regression, because in all cases there was only one significant factor in univariate analysis.

5. Conclusion

Based on our results, we recommend using AIMS 65 scoring system. It's simple and requires few of parameters to be counted. Also, it proved to be the most efficient in predicting in-hospital mortality.

Acknowledgements: Authors would like to thank Miranda Pursley for manuscript editing and proofreading.

Conflict of interests: The authors declare no conflict of interests.

References:

- [1] Beyda R, Johari D. Tranexamic Acid for Upper Gastrointestinal Bleeding. Acad Emerg Med 2019;26:1181–2. doi:10.1111/acem.13835.
- [2] Schjerning Olsen A-M, McGettigan P, Gerds TA, et al. Risk of gastrointestinal bleeding associated with oral anticoagulation and non-steroidal anti-inflammatory drugs in patients with atrial fibrillation: a nationwide study. Eur Hear J - Cardiovasc Pharmacother 2019. doi:10.1093/ehjcvp/pvz069.

- [3] Werdan K, Braun-Dullaeus R, Presek P. Antikoagulation bei Vorhofflimmern: Zauberwort NOAK. *Dtsch Arztebl Int* 2013;110:523–4. doi:10.3238/arztebl.2013.0523.
- [4] Rockall TA, Logan RFA, Devlin HB, Northfield TC. Risk assessment after acute upper gastrointestinal haemorrhage. *Gut* 1996;38:316–21. doi:10.1136/gut.38.3.316.
- [5] Committee BS of GE. Non-variceal upper gastrointestinal haemorrhage: guidelines. *Gut* 2002;51 Suppl 4:iv1-6.
- [6] Blatchford O, Murray WR, Blatchford M. A risk score to predict need for treatment for upper-gastrointestinal haemorrhage. *Lancet* 2000;356:1318–21. doi:10.1016/S0140-6736(00)02816-6.
- [7] Saltzman JR, Tabak YP, Hyett BH, et al. A simple risk score accurately predicts in-hospital mortality, length of stay, and cost in acute upper GI bleeding. *Gastrointest Endosc* 2011;74:1215–24. doi:10.1016/j.gie.2011.06.024.
- [8] Forrest JAH, Finlayson NDC, Shearman DJC. ENDOSCOPY IN GASTROINTESTINAL BLEEDING. *Lancet* 1974;304:394–7. doi:10.1016/S0140-6736(74)91770-X.
- [9] M. R, A. M, R. B, et al. Risk stratification in acute upper GI bleeding: Comparison of the AIMS65 score with the Glasgow-Blatchford and Rockall scoring systems. *Gastrointest Endosc* 2016;83:1151–60. doi:10.1016/j.gie.2015.10.021 LK -
- [10] Nakamura S, Matsumoto T, Sugimori H, et al. Emergency endoscopy for acute gastrointestinal bleeding: prognostic value of endoscopic hemostasis and the AIMS65 score in Japanese patients. *Dig Endosc* 2014;26:369–76. doi:10.1111/den.12187.
- [11] Hyett BH, Abougergi MS, Charpentier JP, et al. The AIMS65 score compared with the Glasgow-Blatchford score in predicting outcomes in upper GI bleeding. *Gastrointest*

Endosc 2013;77:551–7. doi:10.1016/j.gie.2012.11.022.

- [12] Choe JW, Kim SY, Hyun JJ, et al. Is the AIMS 65 score useful in predicting clinical outcomes in Korean patients with variceal and nonvariceal upper gastrointestinal bleeding? Gut Liver 2017;11:813–20. doi:10.5009/gnl16607.
- [13] Stanley AJ, Laine L, Dalton HR, et al. Comparison of risk scoring systems for patients presenting with upper gastrointestinal bleeding: International multicentre prospective study. BMJ 2017;356. doi:10.1136/bmj.i6432.
- [14] Shafaghi A, Gharibpoor F, Mahdipour Z, Samadani AA. Comparison of three risk Scores to predict outcomes in upper gastrointestinal bleeding; modifying Glasgow Blatchford with Albumin. Rom J Intern Med 2019;0. doi:10.2478/rjim-2019-0016.
- [15] Rout G, Sharma S, Gunjan D, et al. Comparison of various prognostic scores in variceal and non-variceal upper gastrointestinal bleeding: A prospective cohort study. Indian J Gastroenterol 2019;38:158–66. doi:10.1007/s12664-018-0928-8.
- [16] Chandra S. AIMS65 score predicts short-term mortality but not the need for intervention in acute upper GI bleeding. Gastrointest Endosc 2013;78:381–2. doi:10.1016/j.gie.2013.02.034.

Figure Captions:

1. Receiver operative curves for mortality rate
2. Receiver operative curves for bleeding recurrence

Preprint

Table 1.

Baseline characteristics of study group	
Parameter	Data
Number of patients	137
Number of females	40(29.20%)
Mean patients' age \pm SD [years]	64.13 \pm 16.04
Cause of the bleeding	
Duodenal ulcer	85(62%)
Gastric ulcer	43(31%)
Mallory-Weiss syndrome	7(5%)
Dielaufoy malformation	2(2%)
Forrest classification of the bleeding from ulcer	
Ia	25 (18%)
Ib	85 (62%)
IIa	14 (10%)
IIb	11 (8%)
IIc	2 (2%)
III	0 (0%)
Median HR at admission (IQR) [bpm]	109 (95-119)
Median systolic blood pressure at admission (IQR) [mmHg]	92 (86-97)
Number of patients needing PRBC transfusion	102 (74.45%)
Number of patients needing > 4 units of PRBC transfusion	29 (21.67%)
Median number of PRBC units transfused (IQR)	2 (0-4)
Number of patients needing FFP transfusion	51 (37,23%)
Mean Hemoglobin level at admission \pm SD [g/dL]	8.62 \pm 2.52
Median INR value at admission (IQR)	1.33 (1.14-1.89)

Table 2.


Analysis of gastrointestinal bleeding assesment scales in study group	
Parameter	Data
Rockall Scale:	
Median score in Rockall Scale (IQR)	5 (3-6)
Number of patients with intermediate risk of mortality	111 (81.02%)
Number of patients with high risk of mortality	6 (4.38%)
Number of patients with good prognosis	20 (14.60%)
Glasgow-Batchford Score:	
Median score in Glasgow-Batchford (IQR)	10 (8-13)
Number of patients with >50% risk of needing of intervention	118 (86.13%)
AIMS65 Score:	
Median AIMS65 score (IQR)	1 (1-2)
Number of patients with 1.2% in-hospital mortality risk	56 (40.88%)
Number of patients with 5.3% in-hospital mortality risk	48 (35.04%)
Number of patients with 10.3% in-hospital mortality risk	15 (10.95%)

Table 3.


Risk factors for need of transfusion > 4 units of PRBC v1.0			
Parameter	OR	95% CI	P value
Univariate analysis:			
Male	0.895	0.367-2.179	0.806
Every year of age	0.988	0.963-1.014	0.356
Cardiovascular disease	0.678	0.251-1.830	0.443
Pulmonary disease	1.665	0.403-6.885	0.482
Metabolic disease	3.296	0.825-13.174	0.092
Liver disease	1.759	0.704-4.394	0.227
Alcohol abuse	3.006	0.878-10.294	0.080
Kidney failure	1.676	0.649-4.326	0.286
Rockall Score > 8 points(high risk of mortality)	1.926	0.335-11.0076	0.463
Glasgow-Blatchford Score > 5 points (>50% risk of needing of intervention)	5.600	0.715-43.845	0.101
AIMS65 Score > 2 points (10.3% In-hospital risk of mortality)	3.977	1.305-12.122	0.015

Table 4.

Risk factors for need of transfusion FFP v1.0			
Parameter	OR	95% CI	P value
Univariate analysis:			
Male	0.984	0.459-2.106	0.966
Every year of age	1.007	0.985-1.029	0.549
Cardiovascular disease	1.509	0.695-3.273	0.298
Pulmonary disease	0.705	0.174-2.858	0.625
Metabolic disease	2.228	0.570-8.712	0.249
Liver disease	1.823	0.810-4.102	0.147
Alcohol abuse	1.227	0.368-4.089	0.739
Kidney failure	1.346	0.578-3.133	0.490
Rockall Score > 8 points(high risk of mortality)	0.837	0.148-4.738	0.830
Glasgow-Blatchford Score > 5 points (>50% risk of needing of intervention)	3.657	1.010-13.242	0.048
AMIDS65 Score > 2 points (10.3% In-hospital risk of mortality)	2.857	0.953-8.570	0.061



Receiver operative curves for mortality rate



Receiver operative curves for bleeding recurrence