

The eccentric mechanotransduction, neuro-muscular transmission, and structural reversibility of muscle fatty infiltration. An experimental advanced disuse muscle-wasting model of rabbit supraspinatus

Jarostaw Fabiś¹, Marian Danilewicz², Kryspin R. Niedzielski³, Michał Waszczykowski¹, Anna Fabiś-Strobin³, Andrzej Bogucki⁴

¹Department of Arthroscopy, Minimally Invasive Surgery and Sports Traumatology
Medical University of Lodz, Lodz, Poland

²Morphometry Division, Department of Pathology, Medical University of Lodz, Lodz,
Poland

³Clinic of Orthopaedic and Traumatology Polish Mother's Memorial Hospital Research
Institute, Lodz, Poland

⁴Department of Extrapyrimal Diseases, Medical University of Lodz, Lodz, Poland

Submitted: 13 September 2020; **Accepted:** 26 December 2020

Online publication: 5 March 2021

Arch Med Sci 2021; 17 (5): 1400–1407

DOI: DOI: <https://doi.org/10.5114/aoms/131956>

Copyright © 2021 Termedia & Banach

Corresponding author:

Prof. Jarostaw Fabiś
Department of Arthroscopy,
Minimally Invasive
Surgery and Sports
Traumatology
Medical University of Lodz
4 Kosciuszki St
90-419 Lodz, Poland
E-mail: fabis@onet.eu

Abstract

Introduction: Full-thickness rotator cuff tear is present in almost 50% of patients over age 65 years, and its degree is known to be a good predictor of the severity of muscle-wasting (MW) sarcopaenia, also known as fatty degeneration (FD). A FD CT grade > 2° is recognized as a borderline of its reversibility. A disuse model of supraspinatus FD (grade 2) in rabbits provides clinically relevant data. Therefore, the present study evaluates the correlation between eccentric mechanotransduction, neuromuscular transmission (NT), and reversibility of muscle fatty infiltration (MFI) in rabbit supraspinatus FD > 2°.

Material and methods: The supraspinatus tendon was detached from the greater tubercle, infraspinatus, and subscapularis in 16 rabbits. The tendon was reinserted after 12 weeks, and the animals were euthanized 24 weeks after reconstruction. MFI was measured in the middle part of the supraspinatus. Single-fibre EMG (SFEMG) examination of the supraspinatus NT was performed on 4 animals.

Results: The power of analysis was 99%. Significant differences in MFI volume were found between the operated (4.6 ± 1.1%) and the opposite control sides (2.91 ± 0.61%) ($p < 0.001$). SFEMG revealed no significant differences between the disuse and the control supraspinatus muscles ($p > 0.05$); however, 6.5% of the examined muscle fibres exhibited NT disorders combined with blockade of conduction in 2.5% of muscle fibres.

Conclusions: Critical MFI in a disuse model of rabbit supraspinatus FD, CT grade > 2°, is substantially reversible by eccentric training despite subclinical impairment of neuromuscular transmission. In addition, 0.63% reversal of MFI is correlated with 1% hypertrophy of type I and II muscle fibre diameter.

Key words: muscle wasting, eccentric training, fatty degeneration, rabbit supraspinatus, neuromuscular transmission.

Introduction

To improve the quality of life of the aging population, effective treatment of muscle-wasting (MW) diseases [1, 2] is needed [3–5]. One common approach is based on the use of exercise [3–11]. Many studies emphasize the importance of eccentric training [6–11]. As muscles regularly undergo eccentric contraction [6, 8], eccentric mechanotransduction appears to be a natural strategy for training muscle strength [6, 10] and the best activator of muscle mechanosensation [12], and one that may allow the reversal of MW changes [11, 13]. However, this reversal also depends on neuromuscular transmission (NT) [14], a self-contained factor known to stimulate neurogenic muscle atrophy and muscle fatty infiltration (MFI) [15].

Disuse MW changes to the shoulder caused by rotator cuff tear result in dysfunction of the upper limb, and hence a weakening of the second key pillar of life, independence, with such disability being particularly burdensome for the elderly [16]. The prevalence of rotator cuff tear is almost 50% in patients aged over 65 years [17], and its size is known to be a good predictor of the severity of MW or sarcopaenia [18]. Changes to the rotator cuff associated with MW are characterized by atrophy, muscle fibre loss, and muscle fatty infiltration (MFI) [19–21] also known as fatty degeneration (FD) [22].

Although various biomarkers have been proposed to measure the advancement of such MW changes [23], one of the best basic approaches to muscular study for this purpose is believed to be CT [24]. Therefore, CT images are commonly used to classify the degree of FD (MFI) in rotator cuff muscles according to the Goutallier *et al.* classification [22]; a score of grade 2 FD, i.e. tissue containing more muscle than fat, indicates borderline reversibility [22, 25].

A significant part of our knowledge of the development of changes to the rotator cuff muscles associated with disuse MW caused by tendon damage is derived from experimental research. One of the most widely used experimental models is the rabbit model [13, 26–34].

However, to the best of our knowledge, no human or animal data exist regarding the complex relationship between eccentric mechanotransduction, NT, and histomorphometric reversibility of MFI, in the case of advanced MW changes associated with muscle disuse.

The present study continues our previous work [13] by examining the reversibility of type I and II supraspinatus muscle fibre atrophy combined with MFI greater than second degree FD (MFI = 2.28) [29] according to Goutallier *et al.* [22], based on the rabbit model of disuse MW. The present work assumes the following hypotheses: 1) the

presence of critical MFI changes to the supraspinatus corresponding to greater than second degree FD according to Goutallier *et al.* [22] is associated with an impairment of neuromuscular transmission highlighted by SFEMG; 2) in spite of FD greater than second degree, muscle mechanosensation is preserved and the implementation of eccentric mechanotransduction after successful supraspinatus tendon reconstruction may influence the reversibility of MFI; 3) the reversibility of MFI is correlated with muscle fibre type I and II hypertrophy.

Material and methods

The study was approved by the local Animal Bioethics Committee of the Medical University of Lodz (research protocol q/BO/207). The study used a previously described model of FD in the supraspinatus muscle of rabbits achieved 12 weeks after cutting the supraspinatus muscle tendon; the FD was evaluated as being greater than second degree (2.28) according to Goutallier *et al.* following CT examination [22]. Sixteen adult male rabbits of mixed breed (weight 3.8–4.6 kg) were anaesthetized with intravenous Vetbutal injection (30 mg/kg) (Biowet, Pulawy, Poland) supplemented by local infiltration with 1% lidocaine. The left shoulder was exposed, and after shaving and disinfecting the skin the lateral surgical approach was applied. The deltoid muscle was split and retracted, and about 3 mm of the supraspinatus tendon and its fibrocartilaginous part was resected and detached from the greater tubercle, as well as from the subscapularis and infraspinatus tendons (cut interdigitation-split length: 1 cm) [13, 30]. The wounds were closed with layers, and the animals were allowed to move freely within a standard cage. The water was supplemented with metamizole sodium (Polpharma, Poland) to reduce postoperative pain for the first 3 postoperative days.

Twelve weeks after the procedure, the tendon was reinserted under the same anaesthetic conditions. According to our previous work on a similar rabbit supraspinatus model, 12 weeks was a sufficient period for development of grade > 2 FD (2.28) [29] according to Goutallier *et al.* [22]. Thus, the thin scar tissue and the edge of the supraspinatus tendon were resected, and the tendon was reattached to the abraded groove between the humeral head and the greater tubercle with the help of a modified Mason-Allen stitch [35] with a number 2 AMIFIL polyamide non-absorbable suture (Simpo, Poznan, Poland). The reattachment was performed under tension due to the shortening of the tendon. The suture was passed through the groove with the help of a needle and tied over the external cortex. The connections between the

supraspinatus and infraspinatus, as well as the subscapularis, were reconstructed using an AMIL-FIL 3/0 polyamide non-resorbable suture (Simplo, Poznań, Poland). Very strong fixation was observed, and the tendon stump adapted well to the bone bed. The wound was closed in layers. The shoulder was not immobilized post-operatively.

After 6 weeks, the animals were moved from the standard cage to an open one 4 times as large every second week to allow more space for activity: daily changes of water and food were provided in opposite corners. Although two animals were euthanized 8 and 9 weeks after reattachment of the tendon due to an unknown general infection, they were not included in the histomorphometric study due to excessive loss of body weight. Twenty-four weeks after reattachment of the tendon, 10 animals were euthanized by intravenous application of a lethal dose of Vetbutal (100 mg/kg), and full thickness muscle samples were taken from the middle part of the supraspinatus, both from the operated and the non-operated limbs.

Histomorphometry

The muscle samples had previously been used to measure type I and type II muscle fibre diameter in a recent study [13]. In the present study, they were used to determine the cross-sectional areas of MFI regions within the muscle. For this purpose, the muscle samples were stained with haematoxylin and eosin. Care was taken to ensure that the cuts were perpendicular to the long axis of the muscle fibres. Histological morphometry was performed with an image analysis system consisting of a computer running MultiScan image analysis software (CSS-Poland) equipped with an Aver 2000 card (frame grabber, true colour, real time) and a colour TV camera linked to a microscope. This system was programmed to calculate the distance between 2 points and a regulated area of a structure using a stereological net with a regulated number of points. The cross-sectional areas of the MFI regions were measured with the help of a point counting method based on Weibel [36] using a stereological net. The point spacing was 16 µm. The total number of points in each net was 169, and the total area 0.37 mm². Ten fields of the middle part of the muscle were studied under these nets.

SFEMG study

The SFEMG was performed on 4 randomly selected rabbits from a group of 14 animals 12 weeks after the supraspinatus tendon detachment. The animals were anaesthetized under identical anaesthetic conditions, and the supraspinatus muscle and the suprascapular nerve were exposed.

SFEMG stimulation studies were performed with Keypoint (Medtronic) apparatus using extra-muscular axonal stimulation techniques, which allowed direct muscle fibre stimulation to be avoided [16, 37–40]. Bipolar hook electrodes were placed under the suprascapular nerve. The axons were stimulated at 10 Hz (pulses of 50 µs duration). The strength of the stimulus used was 0.7–4 mA. Artifacts related to threshold stimulation were avoided by using supramaximal stimulation. A ring electrode around the ear provided grounding. Recordings were made from the middle part of the supraspinatus muscle with a 26 G × 40 mm SFEMG needle electrode. The high-pass filter was set at 1 kHz and the low-pass filter at 20 kHz. After introduction of the needle electrode into the muscle belly, its position had to be precisely adjusted to find the potentials.

When muscle fibre action potentials are recorded during repetitive nerve stimulation, the latency from the stimulus to the response is subject to some fluctuations. This variation is called neuromuscular jitter, and its value (in µs) is calculated as the mean consecutive difference (MCD) of successive stimulus-response intervals (SRI) using the following formula: $MCD = [(SRI_2 - SRI_1) + (SRI_3 - SRI_2) + \dots + (SRI_{n-1} - SRI_n)] / (n-1)$ [16, 37].

The MCD was calculated automatically by the equipment software as the mean of the absolute differences of 100 consecutive inter-potential intervals between stimuli and single-fibre potential. Twenty potentials were recorded, and the mean MCD for 20 muscle fibres was then calculated during the examination of each muscle. To eliminate the possibility of the direct muscle stimulation only potentials with MCD values ≥ 3.0 µs were accepted for analysis. The number of abnormal potentials and potentials with blocking were recorded. The examination was performed on the operated side and then on the control side.

The upper limit of normal jitter is defined as the mean MCD of the control muscles plus 3 standard deviations [38]. The upper value of the MCD range of the control muscles is considered the individual normative value of a single muscle fibre [37].

Statistical analysis

A power analysis was performed ($\alpha = 0.05$, 80% power). The arithmetic mean and standard deviation were calculated from the basic position measurements. The Shapiro-Wilk test was used to confirm whether the tested values were normally distributed, and the *t*-test and Wilcoxon test were used for further analyses depending on the distribution. Differences with *p*-values < 0.05 were considered significant. All calculations were performed using Statistica ver. 10 software (StatSoft, Inc., 2011, DASS, www.statsoft.com).

Results

The result of the power analysis was 99%. Significant differences were found between the operated and control sides with regard to MFI volume (Student's *t*-test; $p < 0.001$) (Table I). No significant differences were found between the mean jitter values of the disuse and the control supraspinatus muscles (Wilcoxon test; $p = 0.051$) in the SFEMG examination.

The mean MCD value of the unoperated control muscles was $11.79 \pm 80 \mu\text{s}$ (4–28), and the normative MCD value of control animals was $14.19 \mu\text{s}$ (11.79 ± 2.4) (Table II). The normative MCD value of an individual muscle fibre in controls was $28 \mu\text{s}$ (Table II). Only 5 abnormal potentials (MCD) (6.5%) associated with the presence of blocking were observed within 2 fibres (2.5%) of the entire pool of 80 muscle fibres evaluated in the 4 operated muscles (Table II).

Discussion

To our knowledge, this study is the first to document the correlation between eccentric mechanotransduction, NT and the histomorphometric reversibility of MFI in an animal model of changes resulting from critical disuse MW in the supraspinatus, i.e. grade > 2 FD according to Goutallier *et al.* based on CT classification [22].

Our present findings and those of previous studies [13] corroborate those of Gerber *et al.* [41], who reported that passive mechanotransduction of sheep infraspinatus can lead to partial reversal of

muscle atrophy and the inhibition of fatty infiltration, as well as the study by Park *et al.* [42], who proved the reversibility of supraspinatus muscle atrophy after arthroscopic repair of its tendon. However, in contrast, Matsumoto *et al.* [31] did not report hypertrophy or the regression of fatty tissue of rabbit supraspinatus 3 months after reattachment of its tendon. These differences can be accounted for by differences in follow-up time, one study being twice the length of the other, the stitching technique, and the choice of rehabilitation program, as detailed previously [13]. In addition, Rubino *et al.* [32] report the progression of fatty degeneration and muscle atrophy halting 6 months after the rabbit supraspinatus tendon reconstruction. However, their study used a larger cage to stimulate animal activity, with water and food being placed in the same position within the cage.

The rabbit supraspinatus is responsible for the eccentric control of the shoulder when landing on the upper limbs after jumping. In the procedure described by Rubino *et al.* [32], the tendon was reinserted in its primary attachment side, i.e. the greater tubercle, resulting in the restoration of normal eccentric load, leading ultimately to a halt in the progression of MW changes. Our biomechanical modification of supraspinatus tendon reinsertion combined with its adequate resection resulted in greater tendon tension and eccentric load, thus fostering muscle fibre hypertrophy [13] and MIF reversibility. This hypothesis coincides with previous findings by Goldspink *et al.* [43]. The authors proved on the rabbit model that the

Table I. The muscle fatty infiltration volume of rabbit supraspinatus muscle 6 months after the reconstruction of its tendon and its eccentric mechanotransduction

Supraspinatus	Mean	Min.	Max.	SD
Operated	4.6	3.2	6.2	1.1
Control	2.91	2.3	4.1	0.61
Operated			Control	
4.2			2.3	
3.5			2.5	
3.2			2.9	
4.1			2.7	
3.4			2.6	
5.2			4.1	
4.6			3.3	
6.1			2.8	
5.5			3.4	
6.2			2.5	

Table II. The results of SFEMG stimulation of the supraspinatus muscle 3 months after the tendon detachment. MCD - mean calculated difference between jitters in μ s

Muscle	Control muscles				Operated muscles			
	1	2	3	4	1	2	3	4
Animal	1	2	3	4	1	2	3	4
MCD range [μ s]	5–28	2–24	6–25	4–20	6–77	4–173	5–24	4–55
Mean MCD [μ s]	12.95	11.55	11.50	11.15	17.35	25.8	10.65	13.30
Potentials with abnormal MCD/ with blockage	0/0	0/0	0/0	0/0	2/0	2/2	0/0	1/0

stretch by itself prevents collagen accumulation and combined with electrical stimulation is also an anabolic signal inducing rapid and marked growth of skeletal muscle [43].

The development of MFI in cases of disuse MW follows a complex pathomechanism. Itoigawa *et al.* [44] proposed that a decrease in muscle tension caused by a tear in the tendon induces a reduction in the expression of *Wnt10b* mRNA, which is responsible for adipocytogenesis and fatty degeneration. Other studies [20, 45, 46] indicate that chronic unloading of muscle induced by a rotator cuff tear disrupts muscle homeostasis, leading to decreased myogenesis (*Myf5*), lowered metabolism (*PPAR α*), and increased adipogenesis (*CEBP α* , *Lep*, *Wnt10b*). Liu *et al.* [47] postulate that the cellular origin of rotator cuff muscle MFI after its tendon rupture lies with muscle mesenchymal progenitor cells. Earlier works documented the influence of mechanotransduction on the inhibition of mesenchymal stem cell differentiation in adipocytes [48], and adipocyte proliferation [49, 50]. Hence, these findings and those of our present and past studies [13] confirm the positive impact of exercise on preventing increased in MFI [51].

Our current and previous findings [13], together with those of Rubino *et al.* [32], Gerber *et al.* [41], and Park *et al.* [42], suggest that regardless of the severity of disuse MW changes, mechanosensation is preserved in the muscle, and it is capable of responding to mechanotransduction. Eccentric mechanotransduction relates to the simultaneous generation of tensile, contractive, and compressive stresses [11]. Therefore, eccentric mechanotransduction has unique potential to counter the negative impact of MFI on muscle mechanosensation associated with stiffness and the number of adipocytes [52]. In addition, eccentric training also offers advantages of a greater impact on type II muscle fibre mechanosensation and less energetic demands than concentric training [11]. Collectively, these findings indicate that the eccentric load gradually activates the process of transduction of mechanical stimuli to the molecular cascade, thus stimulating muscle fibre hypertrophy and decreas-

ing MFI, as confirmed by our previous [13] and current studies.

Moreover, these studies support data about the skeletal muscle memory related to previously encountered environmental stimuli and the retention of epigenetic information in skeletal muscle cells [53]. Furthermore, this skeletal muscle memory [53] seems to explain why, despite experiencing age-related progressive decline in muscle strength, seniors appear to retain the ability to produce eccentric torque [9].

The present study uses an experimental model described in a previous study [30]. It was found that the presence of grade 2 FD is connected with an increase of MFI by 18.34% (from 2.9% to 21.21%) over a 12-week period following tendon detachment [30]. During the same period, 18.2% atrophy was observed for type I muscle fibres and 19.7% for type II fibres from the same supraspinatus sample [30]. Thus, a 1% decrease in type I and type II muscle fibre size was related to a 1.01% growth in MFI (18.34 : 18.2%) for the former and 0.93% (18.34 : 19.7%) for the latter [30].

The previous [13] and current studies allow a unique assessment of the change in the diameters of type I and II muscle fibres regarding MFI regression by morphometric testing of the same muscle. Their results indicate that following eccentric training of the rabbit supraspinatus muscle after tendon stitching, 26.5% hypertrophy was observed for type I muscle fibres and 26.9% for type II, while MFI was reduced by 16.61%, i.e. from 21.21% to 4.6% (Table I) during the same period [13, 30]. Hence, a 1% increase of type I and type II muscle fibre diameter was found to be connected with a 0.63% reversal of MFI (16.61 : 26.5; 16.61 : 26.9) [13, 30]. Moreover, these results indicate for the first time that type I muscle fibres possessed the same quantitative impact on MFI reversibility as type II muscle fibres, which are mainly responsible for the loss of muscle mass with age [54, 55]); however, the final impact depends on the proportions of particular muscle fibre types [30, 54, 55]. Therefore, although most research studies regarding MW disorders have focused attention on type II muscle fibres, type I fibres possess underestimat-

ed potential and should be considered a separate physiotherapeutic target. This is especially important in light of the recent findings that the loss of satellite cells is more significant in the case of type II fibres than type I fibres [56]; however, muscle fibre type II hypertrophy is connected with an increase of satellite cell content [57].

Neuro-muscular junction dysfunction is currently recognized as an important co-mechanism in the development and progression of changes associated with disuse MW [14, 58–60]. Therefore, the present study examines whether the presence of critical MW changes to the supraspinatus, i.e. grade 2 FD according to the Goutallier *et al.* [22] CT scale, is associated with an impairment of neuromuscular transmission. The most sensitive test for the evaluation of neuromuscular transmission is SFEMG. SFEMG reveals neuromuscular transmission abnormalities caused by both pre- and postsynaptic conditions, and the usefulness of SFEMG has been confirmed also in animal studies [38–40].

Our SFEMG results corroborate those of a recent histopathological study on a very similar supraspinatus rabbit model, which also used 4 rabbits and the same follow-up time [33]. Gayton *et al.* [33] reported the occurrence of 19% MFI hyperplasia; however, this did not correlate with the histopathological state of the neuromuscular junction and was not influenced by the motor endplate. However, in the present study, a few potential relationships were recorded with pathological MCD and blocking (Table II), indicating that rabbits with advanced MW changes, i.e. FD > Grade 2 [29], demonstrated some electrophysiological symptoms of subclinical neuromuscular transmission disorders. The study also suggests that subclinical NT disorder depends more on the muscular side than the potential traction of the suprascapular nerve caused by supraspinatus muscle contraction [61] due to its tendon detachment. In our opinion, this is because the physiological activation of the suprascapular nerve is not blocked in the brain and occurs during every shoulder movement in which the supraspinatus muscle innervated by it is involved. It should also be mentioned that recent experimental studies [62] provide evidence that exercise training attenuates skeletal muscle atrophy through the activation of BDNF pathway. Moreover, the conducted studies also indicate that increasing the expression level of the FTO gene has an influence on the increase of skeletal muscle mass [63].

There are some limitations to our study. Firstly, as in previous studies, adult animals were used as test subjects; this may have influenced the muscle regenerative potential because the process is known to depend on age [55, 56, 64, 65]. What is more, as we know, the commonly used therapy

for lipid disorders in old age can exacerbate MW [23, 66]. Secondly, the SFEMG study was limited to 4 rabbits; however, our findings are in line with those given by Gayton *et al.* [33]. However, our present SFEMG results, indicating MAC deviations and blockage of individual muscle fibres (Table II), indicate that dysfunction of subclinical neuromuscular transmission may be a signal of MW changes [66]. Thirdly, a limitation of our work is the inability to assess whether and to what extent the connective tissue hypertrophy has been reduced. Despite these limitations, our findings, as well as those from our previous study [13], provide new insight into the regenerative potential of muscle. It appears to depend on 3 types of fundamental structural change associated with MW, these being related to muscle fibres, MFI, and NT. Furthermore, it should be emphasised that the chosen model fulfils the strict requirements placed on animal models for MW evaluation [64]: the time points used for the study, i.e. 12 and 24 weeks, conform to human recommendations related to intervention with exercise formulated by the International Sarcopaenia Initiative [3].

Although it is difficult to directly transfer the results of animal tests to humans, close similarities exist between the response of the supraspinatus in animals following tendon detachment and that of rotator cuff rupture in seniors [67–69]. Furthermore, the results of our recent [13] and present studies, for the first time, provide extremely important quantitative histomorphometric data supporting the rationale for incorporating routine eccentric training in a program aimed at preventing the development and progression of MW changes or their reversal in ageing adults [11, 13].

In conclusion, the critical increase of muscle fatty infiltration associated with disuse changes in rabbit supraspinatus muscle, i.e. those greater than CT Grade 2 Fatty Degeneration, is substantially reversible by eccentric training. The SFEMG revealed no significant differences between the disuse and the control supraspinatus muscles; however, 6.5% of the examined muscle fibres exhibited NT disorders combined with blockade of conduction in 2.5% of fibres. A 0.63% decrease of MFI is correlated with 1% hypertrophy of type I and II muscle fibre diameter. The obtained results strongly support the rationale for incorporating routine eccentric training in a program aimed at preventing the development and progression of MW changes or their reversal in ageing adults.

Acknowledgments

The study was supported by internal funds of the Medical University of Lodz, Lodz, Poland (grant number: 503/1-040-02/503-51-001-1900 and 503/1-040-02/503-51-001-17).

Conflict of interest

The authors declare no conflict of interest.

References

1. Anker SD, Coats AJ, Morley JE et al. Muscle wasting disease: a proposal for a new disease classification. *J Cachexia Sarcopenia Muscle* 2014; 5: 1-3.
2. Lewandowicz A, Sławiński P, Kądalska E, Targowski T. Some clarifications of terminology may facilitate sarcopenia assessment. *Arch Med Sci* 2019; 16: 225-32.
3. Cruz-Jentoft AJ, Landi F, Schneider SM, et al. Prevalence of and interventions for sarcopenia in ageing adults: a systematic review. Report of the International Sarcopenia Initiative (EWG SOP and IWGS). *Age Ageing* 2014; 43: 748-59.
4. Aagaard P, Suetta C, Caserotti P, et al. Role of the nervous system in sarcopenia and muscle atrophy with aging: strength training as a countermeasure. *Scand J Med Sci Sports* 2010; 20: 49-64.
5. Liu CJ, Latham NK. Progressive resistance strength training for improving physical function in older adults. *Cochrane Database Syst Rev* 2009; 3: CD002759.
6. Roig M, Macintyre DL, Eng JJ, et al. Preservation of eccentric strength in older adults: evidence, mechanisms and implications for training and rehabilitation. *Exp Gerontol* 2010; 45: 400-9.
7. Hedayatpour N, Falla D. Physiological and neural adaptations to eccentric exercise: mechanisms and considerations for training. *Biomed Res Int* 2015; 2015: 193741.
8. LaStayo PC, Ewy GA, Pierotti DD, et al. The positive effects of negative work: increased muscle strength and decreased fall risk in a frail elderly population. *J Gerontol A Biol Sci Med Sci* 2003; 58: M419-24.
9. Hortobágyi T, Zheng D, Weidner M, et al. The influence of aging on muscle strength and muscle fiber characteristics with special reference to eccentric strength. *J Gerontol A Biol Sci Med Sci* 1995; 50: B399-406.
10. Gault ML, Willems ME. Aging, functional capacity and eccentric exercise training. *Aging Dis* 2013; 4: 351-63.
11. Franchi MV, Reeves ND, Narici MV. Skeletal muscle remodeling in response to eccentric vs. concentric loading: morphological, molecular, and metabolic adaptations. *Front Physiol* 2017; 8: 447.
12. Jahed Z, Shams H, Mehrbod M, et al. Mechanotransduction pathways linking the extracellular matrix to the nucleus. *Int Rev Cell Mol Biol* 2014; 310: 171-220.
13. Fabis J, Danilewicz M, Zwierzchowski JT, Niedzielski K. Atrophy of type I and II muscle fibers is reversible in the case of grade >2 fatty degeneration of the supraspinatus muscle: an experimental study in rabbits. *J Shoulder Elbow Surg* 2016; 25: 487-92.
14. Russ DW, Gregg-Cornell K, Conaway MJ, Clark BC. Evolving concepts on the age-related changes in "muscle quality". *J Cachexia Sarcopenia Muscle* 2012; 3: 95-109.
15. Mallon WJ, Wilson RJ, Basamania CJ. The association of suprascapular neuropathy with massive rotator cuff tears: a preliminary report. *J Shoulder Elbow Surg* 2006; 15: 395-8.
16. Stålberg E, Trontelj JV. The study of normal and abnormal neuromuscular transmission with single fibre electromyography. *J Neurosci Methods* 1997; 74: 145-54.
17. Mather RC, III, Koenig L, Acevedo D, et al. The societal and economic value of rotator cuff repair. *J Bone Joint Surg Am* 2013; 95: 1993-2000.
18. Tashjian RZ. Epidemiology, natural history, and indications for treatment of rotator cuff tears. *Clin Sports Med* 2012; 31: 589-604.
19. Chung SW, Yoon JP, Oh KS, et al. Rotator cuff tear and sarcopenia: are these related? *J Shoulder Elbow Surg* 2016; 25: 249-55.
20. Choo A, McCarthy M, Pichika R, et al. Muscle gene expression patterns in human rotator cuff pathology. *J Bone Joint Surg Am* 2014; 96: 1558-65.
21. Davies MR, Liu X, Lee L, et al. TGF-beta small molecule inhibitor SB431542 reduces rotator cuff muscle fibrosis and fatty infiltration by promoting fibro/adipogenic progenitor apoptosis. *PLoS One* 2016; 11: 0155486.
22. Goutallier D, Postel JM, Bernageau J, et al. Fatty muscle degeneration in cuff ruptures. Pre- and postoperative evaluation by CT scan. *Clin Orthop Relat Res* 1994; 304: 78-83.
23. Muntean DM, Thompson PD, Catapano AL, et al. Statin-associated myopathy and the quest for biomarkers: can we effectively predict statin-associated muscle symptoms? *Drug Discov Today* 2017; 22: 85-96.
24. Heymsfield SB, Adamek M, Gonzalez MC, et al. Assessing skeletal muscle mass: historical overview and state of the art. *J Cachexia Sarcopenia Muscle* 2014; 5: 9-18.
25. Melis B, DeFranco MJ, Chuinard C, et al. Natural history of fatty infiltration and atrophy of the supraspinatus muscle in rotator cuff tears. *Clin Orthop Relat Res* 2010; 468: 1498-505.
26. Chen X, Giambini H, Ben-Abraham E, et al. Effect of bone mineral density on rotator cuff tear: an osteoporotic rabbit model. *PLoS One* 2015; 10: e0139384.
27. Valencia AP, Iyer SR, Pratt SJ, et al. A method to test contractility of the supraspinatus muscle in mouse, rat, and rabbit. *J Appl Physiol* 2016; 120: 310-7.
28. Fabiś J, Kordek P, Bogucki A, et al. Function of the rabbit supraspinatus muscle after detachment of its tendon from the greater tubercle. Observations up to 6 months. *Acta Orthop Scand* 1998; 69: 570-4.
29. Fabiś J, Kordek P, Bogucki A, et al. Function of the rabbit supraspinatus muscle after large detachment of its tendon: 6-week, 3-month, and 6-month observation. *J Shoulder Elbow Surg* 2000; 9: 211-6.
30. Fabis J, Danilewicz M, Omulecka A. Rabbit supraspinatus tendon detachment: effects of size and time after tenotomy on morphometric changes in the muscle. *Acta Orthop Scand* 2001; 72: 282-6.
31. Matsumoto F, Uthoff HK, Trudel G, et al. Delayed tendon reattachment does not reverse atrophy and fat accumulation of the supraspinatus: an experimental study in rabbits. *J Orthop Res* 2002; 20: 357-63.
32. Rubino LJ, Sprott DC, Stills HF Jr, et al. Fatty infiltration does not progress after rotator cuff repair in a rabbit model. *Arthroscopy* 2008; 24: 936-40.
33. Gayton JC, Rubino LJ, Rich MM, et al. Rabbit supraspinatus motor endplates are unaffected by a rotator cuff tear. *J Orthop Res* 2013; 31: 99-104.
34. Abdou MA, Kim GE, Kim J, et al. How long should we wait to create the goutallier stage 2 fatty infiltrations in the rabbit shoulder for repairable rotator cuff tear model? *Biomed Res Int* 2019; 2019: 7387131.
35. Gerber C, Fuchs B, Hodler J. The results of repair of massive tears of the rotator cuff. *J Bone Joint Surg Am* 2000; 82: 505-15.
36. Point Counting Methods. Vol 1. Weibl ER (ed.). Academic Press, London 1979; 100-61.
37. Trontelj JV, Stålberg E. Jitter measurement by axonal micro-stimulation. Guidelines and technical notes. *Electroencephalogr Clin Neurophysiol* 1992; 85: 30-7.

38. Añor S, Lipsitz D, Williams DC, et al. Evaluation of jitter by stimulated single-fiber electromyography in normal dogs. *J Vet Intern Med* 2003; 17: 545-50.
39. Gooch CL, Mosier DR. Stimulated single fiber electromyography in the mouse: techniques and normative data. *Muscle Nerve* 2001; 24: 941-5.
40. Finley DB, Wang X, Graff JE, et al. Single fiber electromyographic jitter and detection of acute changes in neuromuscular function in young and adult rats. *J Pharmacol Toxicol Methods* 2009; 59: 108-19.
41. Gerber C, Meyer DC, Frey E, et al. Reversion of structural muscle changes caused by chronic rotator cuff tears using continuous musculotendinous traction. An experimental study in sheep. *J Shoulder Elbow Surg* 2009; 18: 163-71.
42. Park YB, Ryu HY, Hong JH, et al. Reversibility of supraspinatus muscle atrophy in tendon-bone healing after arthroscopic rotator cuff repair. *Am J Sports Med* 2016; 44: 981-8.
43. Goldspink DF, Easton J, Winterburn SK, et al. The role of passive stretch and repetitive electrical stimulation in preventing skeletal muscle atrophy while reprogramming gene expression to improve fatigue resistance. *J Card Surg* 1991; 6 (1 Suppl): 218-24.
44. Itoigawa Y, Kishimoto KN, Sano H, et al. Molecular mechanism of fatty degeneration in rotator cuff muscle with tendon rupture. *J Orthop Res* 2011; 29: 861-6.
45. Shah SA, Korpakakis I, Cavinatto L, et al. Rotator cuff muscle degeneration and tear severity related to myogenic, adipogenic, and atrophy genes in human muscle. *J Orthop Res* 2017; 35: 2808-14.
46. Frey E, Regenfelder F, Sussmann P, et al. Adipogenic and myogenic gene expression in rotator cuff muscle of the sheep after tendon tear. *J Orthop Res* 2009; 27: 504-9.
47. Liu X, Ning AY, Chang NC, et al. Investigating the cellular origin of rotator cuff muscle fatty infiltration and fibrosis after injury. *Muscles Ligaments Tendons* 2016; 6: 6-15.
48. Li R, Liang L, Dou Y, et al. Mechanical stretch inhibits mesenchymal stem cell adipogenic differentiation through TGFβ1/Smad2 signaling. *J Biomech* 2015; 48: 3665-71.
49. Shoham N, Gefen A. Mechanotransduction in adipocytes. *J Biomech* 2012; 45: 1-8.
50. Tanabe Y, Koga M, Saito M, et al. Inhibition of adipocyte differentiation by mechanical stretching through ERK-mediated downregulation of PPARγ. *J Cell Sci* 2004; 117: 3605-14.
51. Goodpaster BH, Chomentowski P, Ward BK, et al. Effects of physical activity on strength and skeletal muscle fat infiltration in older adults: a randomized controlled trial. *J Appl Physiol* 2008; 105: 1498-503.
52. Shoham N, Girshovitz P, Katzung R, et al. Adipocyte stiffness increases with accumulation of lipid droplets. *Biophys J* 2014; 106: 1421-31.
53. Sharples AP, Stewart CE, Seaborne RA. Does skeletal muscle have an 'epi'-memory? The role of epigenetics in nutritional programming, metabolic disease, aging and exercise. *Aging Cell* 2016; 15: 603-16.
54. Nilwik R, Snijders T, Leenders M, et al. The decline in skeletal muscle mass with aging is mainly attributed to a reduction in type II muscle fiber size. *Exp Gerontol* 2013; 48: 492-8.
55. Lexell J. Human aging, muscle mass, and fiber type composition. *J Gerontol A Biol Sci Med Sci* 1995; 50: 11-6.
56. Frontera WR, Zayas AR, Rodriguez N. Aging of human muscle: understanding sarcopenia at the single muscle cell level. *Phys Med Rehabil Clin N Am* 2012; 23: 201-7.
57. Verdijk LB, Snijders T, Drost M, et al. Satellite cells in human skeletal muscle; from birth to old age. *Age (Dordr)* 2014; 36: 545-7.
58. Jang YC, Van Remmen H. Age-associated alterations of neuromuscular junction. *Exp Gerontol* 2011; 46: 193-8.
59. Tintignac LA, Brenner HR, Rüegg MA. Mechanisms regulating neuromuscular junction development and function and causes of muscle wasting. *Physiol Rev* 2015; 95: 809-52.
60. Gonzalez-Freire M, de Cabo R, Studenski SA, et al. The neuromuscular junction: aging at the crossroad between nerves and muscle. *Front Aging Neurosci* 2014; 6: 208.
61. Fabis-Strobin A, Topol T, Fabis J, et al. A new anatomical insight into the aetiology of lateral trunk of suprascapular nerve neuropathy: isolated infraspinatus atrophy. *Surg Radiol Anat* 2018; 40: 333-41.
62. Zhang Z, Wang B, Fei A. BDNF contributes to the skeletal muscle anti-atrophic effect of exercise training through AMPK-PGC1α signaling in heart failure mice. *Arch Med Sci* 2019; 15: 214-22.
63. Doaei S, Kalantari N, Mohammadi NK, et al. Up-regulation of FTO gene expression was associated with increase in skeletal muscle mass in overweight male adolescents. *Arch Med Sci* 2019; 15: 1133-7.
64. Ishida J, Saitoh M, Doehner W, et al. Animal models of cachexia and sarcopenia in chronic illness: cardiac function, body composition changes and therapeutic results. *Int J Cardiol* 2017; 238: 12-8.
65. Kragstrup TW, Kjaer M, Mackey AL. Structural, biochemical, and functional changes in skeletal muscle extracellular matrix with aging. *Scand J Med Sci Sports* 2011; 21: 749-57.
66. Bielecka-Dabrowa A, Fabis J, Mikhailidis DP, et al. Pro-sarcopenic effects of statins may limit their effectiveness in patients with heart failure. *Trends Pharmacol Sci* 2018; 39: 327-436.
67. Deschenes MR, Roby MA, Eason MK, et al. Remodeling of the neuromuscular junction precedes sarcopenia related alterations in myofibers. *Exp Gerontol* 2010; 45: 389-93.
68. Edelstein L, Thomas SJ, Soslowky LJ. Rotator cuff tears: what have we learned from animal models? *J Musculoskelet Neuronal Interact* 2011; 11: 150-62.
69. Mannava S, Plate JF, Tuohy CJ, et al. The science of rotator cuff tears: translating animal models to clinical recommendations using simulation analysis. *Knee Surg Sports Traumatol Arthrosc* 2013; 21: 1610-9.