

# Contracting COVID-19 in the first and second trimester of pregnancy: what we know – a concise qualitative systematic review

Jasmine Abu-Amara<sup>1</sup>, Dawid Szpecht<sup>2</sup>, Salwan R. Al-Saad<sup>1</sup>, Lukasz M. Karbowski<sup>1</sup>

<sup>1</sup>Poznan University of Medical Sciences, Poznan, Poland

<sup>2</sup>Department of Neonatology, Poznan University of Medical Sciences, Poznan, Poland

**Submitted:** 27 October 2020; **Accepted:** 7 February 2021

**Online publication:** 19 March 2021

Arch Med Sci 2021; 17 (6): 1548–1557

DOI: <https://doi.org/10.5114/aoms/133121>

Copyright © 2021 Termedia & Banach

**Corresponding author:**

Dawid Szpecht MD, PhD,

Assist. Prof.

Department of Neonatology

Poznan University

of Medical Sciences

33 Polna St

60-535 Poznan, Poland

E-mail: [dawid.szpecht@poczta.fm](mailto:dawid.szpecht@poczta.fm)

## Abstract

**Introduction:** Recent medical literature has drawn attention to the possible influence of COVID-19 on the course of pregnancies. As the coherence of results seems to vary, especially in relation to first and second trimester pregnancies, a concise qualitative systematic review can shed light on the most recent data.

**Material and methods:** A structured systematic search was performed to collect all COVID-19 pregnancy-related articles published between January 1 and September 16, 2020. Two independent reviewers evaluated studies using the STROBE statement in combination with the CERQual quality assessment of findings.

**Results:** In total 1387 articles were screened and finally 22 studies were evaluated (179 1<sup>st</sup>/2<sup>nd</sup> trimesters of pregnant women with COVID-19). The majority of reported mothers who contracted COVID-19 during 1<sup>st</sup>/2<sup>nd</sup> trimesters are yet to complete their pregnancy.

**Conclusions:** Evidently a limited amount of data is available. Usually, mothers and newborns are discharged from the hospital without any serious complications. Further observations are imperative.

**Key words:** COVID-19, pregnancy, SARS-CoV-2, maternal, gestation, first trimester, second trimester.

## Introduction

The disease COVID-19, which is caused by SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2), has had large scale ramifications as a global pandemic [1, 2]. The virus was first detected in Wuhan, China in December 2019, and has since progressively spread worldwide and infected more than 91 million people from numerous countries [3–5]. Emerging data have suggested a generally comparable disease course in pregnant and non-pregnant women [6–8]. However, the physiological changes accompanying pregnancy (such as decreased functional residual volume, increased stroke volume, and edema of the respiratory tract mucosa) [9, 10] and the possibility of adverse neonatal complications necessitate vigilant efforts to thoroughly assess possible risk factors for poor SARS-CoV-2 outcome in the pregnant population.

Previous reports have associated an increased risk of perinatal mortality and preterm birth in pregnant women diagnosed with viral pneumonia [11–13]. In 2004, SARS-CoV-1 was linked to an increase rate of maternal morbidity and mortality, intrauterine growth restriction, mis-

carriage and preterm delivery amongst infected mothers [14, 15]. However, it is not yet clear how SARS-CoV-2 may influence pregnancy and the risk factors it presents to the mother and unborn child [16]. More precisely, due to the novelty of the virus and relatively recent spread, there are extremely scarce data on maternal and perinatal outcomes when the infection is contracted in the first or second trimester of pregnancy. The most recent report by the CDC placed emphasis on COVID-19 adverse effects on pregnancy and urged the need for more extensive investigation in this field [17]. Assessing the relationship of COVID-19 with the early gestational trimesters of pregnancy is key to effective counseling, and offers invaluable insight on how the pandemic may influence pregnancy at large in the foreseeable future.

## Material and methods

### Search strategy and selection

The present review paper was guided by established protocols for healthcare systematic reviews [18]. A systematic search was conducted by two independent reviewers, using the primary databases PubMed and ScienceDirect. The search included articles published from January 1 to December 27, 2020. A secondary database, Google Scholar, was also used to provide broader coverage of published literature. The keywords “COVID-19” OR “SARS-CoV-2” were followed by each of “pregnancy” OR “gestation” OR “maternal” during the search. Reference lists of high value studies were also screened for potential relevant studies.

Reports included in this review consisted of case-report, cross-sectional, case-control, cohort (both prospective and retrospective) and case series. Studies that reported pregnant women, who contracted COVID-19 in either their first (1–12 weeks of gestation) or second trimester (13–28 weeks gestation), were of main interest. Duplicate studies, unrelated topics, reviews, guidelines, surveys, non-English language studies, and studies that included COVID-19 positive mothers contracting the virus in the third trimester were all excluded based on view of the abstract and paper. Any studies with unclear distinction on time of COVID-19 contraction during pregnancy were, after thorough assessment, not considered for results.

### Retrieving relevant studies and data extraction

Following the initial search to identify possible relevant studies, the two reviewers excluded all duplicates. Next, the studies were briefly screened for their overall relatedness to the topic and excluded

if deemed not valuable to the systematic review. A list was compiled of the studies with potential relevant data, and the two reviewers independently and thoroughly assessed them for eligibility. Following this process, a final list of relevant studies was compiled. Consensus between reviewers was reached concerning the final studies included.

Having established the group of studies that comprise the systematic review, data were extracted by the two reviewers independently into a standard form. The form included structured subheadings of the clinical data of interest, such as: symptoms, pregnancy-related complications, laboratory and radiological findings, maternal and neonatal outcomes (stillbirth (death after 20 weeks of gestation), miscarriage (death before 20 weeks of gestation), terminated, delivered/alive, or ongoing pregnancy), and others (drugs administered, placental changes). The data were cross-checked for any inaccuracies by a third reviewer. A high level of scrutiny was applied as the data extracted strictly concerned 1<sup>st</sup> and 2<sup>nd</sup> trimester contraction of COVID-19 in pregnant women. Any data that were merged with 3<sup>rd</sup> trimester mothers or did not clearly attribute the data to the trimesters of interest were excluded. There was a discussion and a consensus was reached between all reviewers as to which variables to include in the qualitative synthesis.

To give an estimate of overall incidence, a combined incidence rate of certain clinical phenomena (symptoms, comorbidities, complications, etc.) was calculated by adding the individual incidences in the respective studies and constructing a total percentage. Only studies with contributing data were included in the calculation. The constructed values should be interpreted with caution.

### Quality assessment

A combination of the STROBE statement (Strengthening the Reporting of Observational Studies in Epidemiology) [19] and CERQual approach [20] was implemented in quality assessment of studies and findings. Studies included in the systematic review were individually evaluated using the STROBE statement, with careful appraisal of methodological limitations, studied population, epidemiological information (such as mode of transmission), follow-up information, selection and other bias, as well as incomplete reporting of data.

Providing a transparent explanation, and following the CERQual assessment outline, the review findings were then allocated a confidence score. The confidence score is based on the combined quality of studies contributing to the finding, with particular emphasis on adequacy of data, methodological limitations, consistency in findings or coherence, and relevance of article to




Figure 1. Study selection diagram

review topic. Confidence was assigned using one of the four statements: very low, low, moderate, and high (“unclear”, “it is possible”, “likely”, and “very likely” – that the review finding is a reasonable representation of phenomena, respectively). The constructed review follows the standardized PRISMA guidelines to allow for elaborate presentation [21].

## Results

The search revealed 4585 findings, of which 3196 were noted as duplicates. Of the total 1389 articles, 1262 studies were excluded based on screening of the title, abstract, and brief view of the paper. The majority of exclusions were based on unrelatedness to subject matter or study type (reviews, guidelines). Full text assessment of the remaining 127 articles yielded a total of 24 studies that met the inclusion criteria and were considered relevant to the review (Figure 1). One hundred and three records were removed largely due to an improper separation of data between 1<sup>st</sup>/2<sup>nd</sup> and 3<sup>rd</sup> trimester COVID-19 patients, with the majority reporting combined results. A combined total of 220 first and second trimester pregnant women with COVID-19 were included.

### Study characteristics

The majority of the studies were published between March and December 2020 and they

originated from 11 different countries. Maternal age range was 21–49 years. The range of gestation when the mothers contracted the virus was 5–27 weeks. Most patients had COVID-19 diagnosis confirmed by positive RT-PCR. At the time of publication, 10 studies reported ongoing pregnancies; of the remaining 14 studies, 3 reported neonatal survival while rest had unclear status (Table I). In one particular study reported through the CDC, stillborn or fetal death (at < 20 weeks) was observed in 4 infants; the outcomes of the remaining neonates remained unspecified [17]. Study characteristics and clinical features can be found in Table I.

### Symptoms

All studies [17, 22–44] disclosed symptom presentation, pertaining to 220 pregnant women. Despite a majority of women not reporting any symptoms, the most frequently observed symptoms and clinical signs were fever (86/220 = 39%) and cough (73/220 = 33%). To a smaller extent, dyspnea (52/220 = 24%), nausea and vomiting (27/220 = 12%), and loss of taste or smell (57/220 = 26%) were observed. Out of the 220 first and second trimester pregnant women included in this review, 132 were symptomatic (60%).

### Laboratory and radiological findings

Radiological findings of 24 pregnant women in 12 studies were reported [22, 26, 28–30, 32–34, 36, 37, 40, 41]. Pneumonia and ground glass appearance were the first ( $n = 12/24$ : 50%) and second ( $n = 5/24$ : 21%) most common observed radiological findings respectively. More than half of the reported findings revealed bilateral distribution of the disease. Normal radiological findings were also noted in 2/24 patients. The laboratory results of 21 pregnant women were reported. Elevated C-reactive protein as well as lymphocytopenia were observed in 4/21 women (19%), and high levels of hepatic markers (AST/ALT) were recorded in 9 patients (43%) (Table I).

### Pregnancy-related complications

Only a portion of the studies (9) reported some sort of complication during the span of pregnancy which pertained to 15 patients [22–27, 37, 39, 40]. Disseminated intravascular coagulation (DIC) (20%; 3/15), preeclampsia (13%; 2/15), as well as liver cytolysis, chorioamnionitis, ARDS, hyperemesis gravidarum, pneumothorax and gestational diabetes (each observed in 1/15 of the reported patients) were the most commonly observed complications.

Table 1. Symptoms, complications, laboratory and radiological findings, neonatal outcomes

Authors	Country	Pub. date	No. of patients (trimester)	Symptoms	Pregnancy-related complications	Laboratory findings	Radiological findings	Neonatal outcomes
Delahoy <i>et al.</i> [17]	USA	Sep 2020	14 (1 <sup>st</sup> trim) 61 (2 <sup>nd</sup> trim)	Fever, cough, myalgia, SOB	N/A	N/A	N/A	4: miscarriage (1 <sup>st</sup> trim)
Hantoushadeh <i>et al.</i> [22]	Iran	July 2020	3 (2 <sup>nd</sup> trim)	Fever, cough, dyspnea, tachycardia	ARDS, pneumothorax, ARF, cardiopulmonary collapse, septic shock, DIC, left HF	N/A	Patchy ground-glass feature on CT	2 (twins): IUFD 2 (twins): premature labor; death at day 3 1: stillbirth
Browne <i>et al.</i> [23]	USA	June 2020	1 (2 <sup>nd</sup> trim)	Myalgia, fever, cough	Hyperemesis gravidarum, acid reflux	Elevated WBC, low HCT	N/A	Ongoing twin preg
Baud <i>et al.</i> [24]	Switzerland	June 2020	1 (2 <sup>nd</sup> trim)	Fever, cough, myalgia, fatigue	Funifinitis	N/A	N/A	Stillbirth
Richtmann <i>et al.</i> [25]	Brazil	July 2020	3 (2 <sup>nd</sup> trim)	Fever, cough, myalgia	Chorioamnionitis	Normal	N/A	3: stillbirth
Hosier <i>et al.</i> [26]	USA	June 2020	1 (2 <sup>nd</sup> trim)	Fever, cough, myalgia, fatigue, diarrhea	Preeclampsia, DIC	Elevated ALT, AST, urine protein, prolonged PTT, decreased fibrinogen	Hazy opacity	Opted for termination of preg
Garcia-Manau <i>et al.</i> [27]	Spain	July 2020	2 (2 <sup>nd</sup> trim)	"Mild symptoms"	DIC, preeclampsia, Fetal skin edema on ultrasound	N/A	N/A	Ongoing preg
Lokken <i>et al.</i> [28]	USA	Aug 2020	3 (2 <sup>nd</sup> trim)	Dyspnea	N/A	Normal	Consolidation	Ongoing preg
Wu <i>et al.</i> [29]	China	Aug 2020	5 (1 <sup>st</sup> trim) 2 (2 <sup>nd</sup> trim)	Fever, cough, myalgia	N/A	Elevated ALT, AST and CRP, lymphocytopenia	Pneumonia on CT	1: chemical preg (1 <sup>st</sup> trim) 6: ongoing preg
Yu <i>et al.</i> [30]	China	April 2020	2 (1 <sup>st</sup> trim)	Fever, diarrhea, cough, dyspnea	N/A	N/A	Pneumonia on CT	Ongoing preg
Liu <i>et al.</i> [31]	China	March 2020	2 (2 <sup>nd</sup> trim)	Fever, fatigue, dyspnea	N/A	N/A	N/A	Both neonates survived
Wang <i>et al.</i> [32]	China	July 2020	1 (2 <sup>nd</sup> trim)	Myalgia, fever, cough, dyspnea	N/A	Decreased Hb content and serum albumin, elevated neutrophil ratio and ALT, AST	Consolidation	Ongoing preg

Table I. Cont.

Authors	Country	Pub. date	No. of patients (trimester)	Symptoms	Pregnancy-related complications	Laboratory findings	Radiological findings	Neonatal outcomes
Panichaya <i>et al.</i> [33]	Thailand	July 2020	1 (2 <sup>nd</sup> trim)	Fever, chest discomfort	N/A	Normal	Normal	Opted for termination of preg
Inchingolo <i>et al.</i> [34]	Italy	July 2020	1 (2 <sup>nd</sup> trim)	Fever, cough	N/A	N/A	Pneumonia	N/A
Crovetto <i>et al.</i> [35]	Spain	Aug 2020	54 (1 <sup>st</sup> trim)	Fever, cough, diarrhea, loss of taste or smell	N/A	N/A	N/A	N/A
Hachem <i>et al.</i> [36]	France	July 2020	1 (2 <sup>nd</sup> trim)	Normal	N/A	Elevated CRP, lymphocytopenia	Pneumonia on CTPA	Miscarriage
Piersigilli <i>et al.</i> [37]	Belgium	June 2020	1 (2 <sup>nd</sup> trim)	Normal	Cholecystitis, HELLP syndrome	Elevated CRP	Pneumonia on X-ray	Neonate survived
Cosma <i>et al.</i> [38]	Italy	July 2020	14 (1 <sup>st</sup> trim)	Fever, diarrhea, cough, loss taste/smell, arthralgia	N/A	N/A	N/A	N/A
Lamazou <i>et al.</i> [39]	France	May 2020	1 (1 <sup>st</sup> trim)	Dyspnea, nausea, asthenia	Liver (hepatic) cytolysis	N/A	N/A	Ongoing preg
Tang <i>et al.</i> [40]	China	July 2020	2 (2 <sup>nd</sup> trim)	Fever, dyspnea	Gestational diabetes, neonatal jaundice	Elevated CRP, lymphocytopenia	Ground glass opacity on CT	Neonates survived
Hong <i>et al.</i> [41]	USA	July 2020	1 (2 <sup>nd</sup> trim)	Fever, cough, dyspnea, myalgia	N/A	N/A	Normal	Ongoing preg
Rubio Lorente <i>et al.</i> [42]	Spain	Aug 2020	1 (1 <sup>st</sup> trim) 1 (2 <sup>nd</sup> trim)	Fever, rhinorrhea, cough	N/A	N/A	N/A	1: ongoing preg 1: abortion (1 <sup>st</sup> trim)
la Cour Freiesleben <i>et al.</i> [43]	Denmark	Dec 2020	18 (1 <sup>st</sup> trim)	Loss of taste or smell, dry cough, fatigue, fever	N/A	N/A	N/A	1 fetal death 17 ongoing preg
Cosma <i>et al.</i> [44]	Italy	Oct 2020	23 (1 <sup>st</sup> trim)	Fever, cough, arthralgia	N/A	N/A	N/A	N/A

Trim – trimester, DIC – disseminated intravascular coagulation, Hb – hemoglobin, ALT – alanine aminotransferase, AST – aspartate aminotransferase, CRP – C-reactive protein, PTT – partial thromboplastin time, Preg – pregnancy, N/A – not available, HCT – hematocrit, CTPA – CT pulmonary angiogram, ARF – acute renal failure, HF – heart failure, IUFD – intrauterine fetal demise, SOB – shortness of breath.

### Maternal outcomes

The overwhelming majority of mothers included in this review were successfully discharged from the hospital or were alive as of the date of publication (220 women). However, Hantoushza-deh *et al.* reported 3 cases of which COVID-19 maternal death was the outcome [22]. No other significant maternal outcomes were reported.

### Neonatal outcomes

Out of 223 neonates, the outcomes of 60 infants (31/60: maternal infection during 1<sup>st</sup> trimester and 29/60: 2<sup>nd</sup> trimester infections) from 20 studies were specified as deceased/terminated, miscarriage, alive/delivered, or still ongoing pregnancy [17, 22–33, 36, 37, 39–43]. While the majority of 1<sup>st</sup> trimester infection neonates had a reported status of ongoing pregnancy ( $n = 24/31$ : 77.4%), 4/31 (12.9%) reported miscarriages, 1/31 (3.2%) abortion, 1 (3.2%) stillbirth and 1 (3.2%) chemical pregnancy were also noted.

In the studies with documented maternal COVID-19 infection during the second trimester of pregnancy, neonatal outcomes were as follows: 5/29 (17.2%) completed term and were delivered healthy, 14/29 (48.2%) had ongoing pregnancy, 5/29 (17.2%) stillbirths, 1/29 (3.4%) miscarriage, 2/29 (6.9%) intrauterine fetal demise (IUFD), and 2/29 (6.9%) opted for termination of pregnancy. The rest of the neonates ( $n = 163/223$  (73.1%)) had unreported status concerning the outcome of pregnancy (unclear whether delivered, deceased, or ongoing pregnancy).

### Others

A minority of studies [24–26, 33, 42] documented the condition of 7 placentas upon delivery, of which 2/7 were normal. Histopathological findings documented were: acute villitis and intervillitis (2/7 (28.6%)), acute deciduitis (1/7 (14.3%)), diffuse perivillous fibrin (5/7 (71.4%)) and an inflammatory infiltrate composed of either neutrophils, T-lymphocytes or macrophages (2/7 (28.6%)). One of the placental histopathological investigations determined no membrane inflammation or viral inclusion [33]. A case of decidual inflammation was also reported, but was eventually declared to be caused by abortion of the first trimester rather than COVID-19 disease [42].

Drug treatment of 12 patients in 9 studies was reported [22–27, 31, 36, 41]. Azithromycin (5/12 = 42%) and hydroxychloroquine (5/12 = 42%) were the most commonly administered medications. Other medications such as lopinavir-ritonavir (4/12 = 33%), oseltamivir (3/12 = 33%), acetaminophen (2/12 = 17%) and ceftriaxone (2/12 = 17%) were also reported.

### Qualitative synthesis

The majority of included studies were retrospective. Epidemiological information and mode of contraction were only recorded in 7 of the studies [23, 29, 30, 32, 33, 36, 40]. The virus was usually transmitted through direct contact with an exposed spouse. Mainly, studies included only hospitalized or severe cases, which was noted as a minority of cases in the respective article, and as such was considered as selection bias [22, 25, 26, 28, 30–33, 35, 36]. Other methodological limitations include short follow-up, with a majority being less than 1 month. Nine studies were of women with still ongoing pregnancy [23, 27–30, 32, 39, 41, 42], and only 3 studies had recorded the complete term of pregnancy [31, 37, 40]. Pregnancy-related complications were under-reported, with 9/24 studies mentioning incidence of any complications [22–27, 37, 39, 40]. Overall limited data were available, as 15 studies reported  $\leq 2$  patients [23, 24, 26, 27, 30–34, 36, 37, 39–42]. Lack of extensive data, short follow-up, and a large proportion of ongoing pregnancies resulted in no findings receiving a “high” confidence score. CERQual confidence in review findings and their respective explanation can be found in Table II.

### Discussion

Despite the majority of symptoms presenting as mild, pregnant women and newborns are a worrisome population considering that there are very limited data in regards to the outcome and complications in COVID-19 infected pregnant women [45]. During the SARS-CoV-1 outbreak in 2002–2003, a high maternal mortality rate was observed [14, 46]. Infected pregnant mothers who survived reported experiencing 1<sup>st</sup> trimester miscarriages and intrauterine growth restriction (IUGR) during their 2<sup>nd</sup> and 3<sup>rd</sup> trimesters [14]. Considering the similarities, including genetic makeup and mode of cell entry, between SARS-CoV-1 and the recent SARS-CoV-2 [47], particular attention should be placed on the pregnant population in order to identify and attend to any possible detrimental complications.

To the best of the author’s knowledge, this is the first qualitative systematic review focused on the clinical characteristics of 1<sup>st</sup>/2<sup>nd</sup> trimester infections of SARS-CoV-2 during pregnancies. Primary data showed high coherence in relation to symptoms, reporting fever (39%) and cough (33%) to be the most prevalent in the population group of 1<sup>st</sup> and 2<sup>nd</sup> trimester pregnant women. A few other studies have reported similar results [48, 49]. Compared to 3<sup>rd</sup> trimester infections with SARS-CoV-2, comparable results have been reported in terms of symptoms. In one study that included 108 SARS-CoV-2 positive pregnant women during

**Table II.** CERQual assessment of review findings

Review findings	CERQual assessment of confidence in the evidence	Explanation of CERQual assessment	Studies contributing to the review finding
<b>Symptoms:</b>			
<p>Majority of pregnant women who contract COVID-19 during the 1<sup>st</sup> or 2<sup>nd</sup> trimester are symptomatic with minor symptoms on admission. Most common: fever, cough, fatigue/myalgia, dyspnea, diarrhea, shortness of breath</p> <hr/> <p>Least common: sore throat, loss of taste or smell, nausea, asthenia, tachycardia</p>	Moderate	7 studies with minor to moderate methodological limitations, of which selection bias was noted due to the studies pertaining to hospitalized patients only. Patient exposure history prior to onset of symptoms was largely under-reported. Relatively adequate data pertaining to 177 1 <sup>st</sup> /2 <sup>nd</sup> trimester pregnant COVID-19 patients. High coherence	17, 22–44
<b>Pregnancy-related complications:</b>			
A wide range of multisystem complications such as preeclampsia, DIC, hyperemesis gravidarum, hypothyroidism, gestational diabetes, neonatal jaundice are noted	Low	5 studies with minor to moderate methodological limitations. 4 studies were admitted with severe cases of COVID-19, raising concern for potential selection bias. Selective reporting bias due to gaps and inconsistency in reporting of pregnancy-related complications. Short follow-up of patients (< 1 month) and incomplete/ongoing course of pregnancy (as of date of publication) largely limit the reflectiveness of findings. Inadequate data pertaining to only 19 pregnant COVID-19 patients. Low to moderate coherence	22–27, 36, 37, 39–41
<b>Outcome:</b>			
<p>Maternal outcome: The short term survival rate amongst mothers is estimated to be very high with complete discharge from the hospital</p> <hr/> <p>Neonatal outcome: A risk for miscarriage or stillbirth is present. Majority of pregnancies remain uneventful for neonatal health upon mother hospitalization. Successful delivery of healthy neonates has been recorded</p>	Low	11 studies reported no further follow-up in regards to the mother or neonates' wellbeing after discharge. Such limitations make findings incomplete for the whole course of pregnancy. 10 studies admitted severe cases, hence concerns for selection bias. Neonatal status was clarified in only 60 out of the 223 neonates, raising concerns for inadequacy of data. Moderate to high coherence in maternal outcome and moderate coherence in neonatal outcome	17, 22–33, 36, 37, 39–42
<b>Laboratory and radiological findings:</b>			
<p>Normal laboratory or radiological findings may occur.</p> <p>Laboratory Findings: Elevated CRP, liver enzymes (ALT/AST) and lymphocytopenia are the most prominent abnormalities. Findings such as prolonged PTT, decreased fibrinogen, Hb content and serum albumin are observed to a lesser extent.</p> <p>Radiological findings: pneumonia features and ground glass opacity are the most commonly occurring. Findings such as consolidation and thickened pleural line can also be observed</p>	Low	7 studies with minor to moderate methodological limitations. Selection bias due to admission of strictly severe cases was the most prominent methodological concern. Majority of data stemming from two geographic regions (China and USA), raising concern for lack of geographic diversity. Inadequate data pertaining to only 28 1 <sup>st</sup> /2 <sup>nd</sup> trimester COVID-19 patients. Moderate coherence	22, 23, 25, 26, 28–30, 32–34, 36, 37, 40

their third trimester, the majority of the patients experienced fever (68%), while cough was observed in only 34% of the patients [50]. Elevated body temperature during pregnancy is associated with disruption of cell migration, damage to the vascular system and death of neuroblast cells [51]. Hence, this is a potential concern during the organogenesis period that takes place during the 1st trimester of pregnancy [52]. Nevertheless, further research is required to analyze the association of certain symptoms with maternal and neonatal outcomes.

According to our systematic review, 1<sup>st</sup> and 2<sup>nd</sup> trimester pregnant women could potentially be at risk of experiencing placental inflammation such as villitis/intervillitis and perivillous fibrin deposition. The literature defines villitis as a damaging inflammation that is characterized by maternal T-cell invasion into chorionic villi [53]. Furthermore, an increase in proinflammatory markers such as tumor necrosis factor (TNF- $\alpha$ ) and interleukin (IL)-6 is characteristic of COVID-19 infection and has been associated with endothelial activation and apoptosis of the trophoblastic cells [54]. It is unclear whether these placental findings can be attributed to the severity of the infection when contracted during the 1<sup>st</sup> and 2<sup>nd</sup> trimester or perhaps be linked to the overall outcome of the neonates' well-being. It is worth noting that 4 of the 17 (23%) recorded neonatal deaths were of pregnancies with developed inflammation such as intervillitis/villitis and perivillous fibrin deposition of the placenta. In some third trimesters delivered placentas, highly abnormal findings such as extensive intervillitis as well as perivillous fibrin were recorded on histopathological examination [55, 56]. Similarly, in a study regarding cases of SARS-CoV-1, three placentas were delivered from pregnancies in which the mothers had acute infection – these were abnormal and demonstrated increased subchorionic and intervillous fibrin [57]. Such findings can be associated with abnormal maternal blood flow to the placenta. Overall, placental histological findings were under-reported in the studies included in this review. This raises concerns for inadequacy of the data and highlights the need for larger samples.

The radiological results of the studies assessed were largely definitive for CT diagnosed pneumonia. Pneumonia features, such as consolidations and ground glass opacities, were the most commonly reported changes observed in roughly half of the patients and one-fifth of the patients, respectively. The majority of the reports indicate a bilateral spread of the disease. In pregnant women infected with COVID-19 during their 3<sup>rd</sup> trimester, similar patterns of consolidations and patchy ground glass opacities were observed on the CT scans [58]. These radiological changes ap-

pear to be the pattern for COVID-19 cases in general, including that of older aged individuals [59]. Chest CT plays a vital role in the initial assessment, treatment and follow-up of COVID-19 patients, and some studies have even shown that chest CT sensitivity could be preferable to RT-PCR for early identification of COVID-19 [60, 61].

In the small portion of studies that detailed therapeutic treatments for the COVID-19 infected pregnant women, azithromycin (42%) and hydroxychloroquine (42%) were the most common, followed by lopinavir-ritonavir (33%) and oseltamivir (33%). The inconsistency in the administration of drugs underlies the lack of sufficient data for therapeutic efficacy against the novel coronavirus. A study concerning the safety of hydroxychloroquine in 1<sup>st</sup> trimester pregnancies reported no notable changes in the nervous/cardiac system formation, prematurity or birth weight of the infants [62]. Despite some data indicating the efficacy of hydroxychloroquine against SARS-CoV-2 [63], consistent clinical results are yet to be presented. Furthermore, lopinavir and ritonavir have also shown an inconsequential effect on pregnancy, and in their study of 955 pregnant women, Roberts *et al.* observed that the incidence of birth defects in children with prenatal exposure to lopinavir/ritonavir did not differ substantially from the control [64]. Hence, similar to hydroxychloroquine, lopinavir and ritonavir are being used as possible deterrents for COVID-19 exacerbation in pregnant women. Further studies are necessary to examine the curative role of pharmaceutical drugs in the context of COVID-19 during pregnancy, and may allow for more consistent and effective treatment plans.

The reported complications associated with COVID-19 pregnancy may influence the adverse outcomes of first and second trimester pregnant women and their newborns. 10 of our neonatal deaths concerned mothers who reported complications during pregnancy. On the other hand, one fetal death was attributed to a miscarriage in a healthy mother with no reported pregnancy-related complication [36]. Moreover, it was noted that preeclampsia and DIC were the most prevalent complications. Narang *et al.* explained that infection with SARS-CoV-2 throughout pregnancy holds the possibility of initiating microvascular dysfunction by causing endotheliitis [65]. Furthermore, ischemia and vasoconstriction may ensue as a result of the microcirculatory dysfunction and systemic inflammation, which can further contribute to a pro-coagulopathic state [65]. Given that complications in pregnant women act as detrimental prognostic factors, it is crucial to pay close attention to the development of any laboratory abnormalities throughout pregnancy.

Due to the novelty of the subject, follow-up of COVID-19 positive first and second trimester

mothers and their newborns is insufficient. The majority of 1<sup>st</sup> and 2<sup>nd</sup> trimester women who got pregnant near the start of the pandemic are yet to give birth or complete pregnancy. Such limitations restrict the assessment of complication development, outcomes (both maternal and neonatal), and course of pregnancies. The presented review provides a comprehensive take on patterns concerning COVID-19 contracted in 1<sup>st</sup> and 2<sup>nd</sup> trimesters, and allows for the construction of appropriate research on the topic.

In conclusion, considering that the majority of mothers who contracted COVID-19 during the 1<sup>st</sup> and 2<sup>nd</sup> trimesters are yet to complete their pregnancy, an evidently scarce amount of data is available concerning neonatal and maternal outcomes. Mothers were largely discharged from the hospital without any serious complications; however, placental inflammatory changes and neonatal death have been reported in a number of cases. Further research is required to determine whether or not contracting SARS-CoV-2 early on in pregnancy poses a consequential risk to either the mother or neonate.

### Conflict of interest

The authors declare no conflict of interest.

### References

- World Health Organization Coronavirus disease 2019 (COVID-19): situation report. Vol. 72. WHO; 2020.
- UpToDate. [www.uptodate.com/contents/coronavirus-disease-2019-covid-19-epidemiology-virology-and-prevention](https://www.uptodate.com/contents/coronavirus-disease-2019-covid-19-epidemiology-virology-and-prevention).
- Abuelgasim E, Saw L, Shirke M, Zeinah M, Harky A. COVID-19: unique public health issues facing Black, Asian and minority ethnic communities. *Curr Probl Cardiol* 2020; 45: 100621.
- Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; 395: 497-506.
- Worldometer. Coronavirus Toll Update: Cases & Deaths by Country of Wuhan, China Virus – Worldometer. [Worldometers.info](https://www.worldometers.info/coronavirus/); 2020. Available from: <https://www.worldometers.info/coronavirus/>.
- Vega M. From the trenches: inpatient management of coronavirus disease 2019 in pregnancy. *Am J Obstet Gynecol* 2020; 2: 100154.
- Gao Y, Ye L, Zhang J, et al. Clinical features and outcomes of pregnant women with COVID-19: a systematic review and meta-analysis. *BMC Infect Dis* 2020; 20: 564.
- Matar R, Alrahmani L, Monzer N, et al. Clinical presentation and outcomes of pregnant women with coronavirus disease 2019: a systematic review and meta-analysis. *Clin Infect Dis* 2021; 72: 521-33.
- Chen YH, Keller J, Wang IT, Lin CC, Lin HC. Pneumonia and pregnancy outcomes: a nationwide population-based study. *Am J Obstet Gynecol* 2012; 207: 288.e1-7.
- Rasmussen SA, Jamieson DJ, Uyeki TM. Effects of influenza on pregnant women and infants. *Am J Obstet Gynecol* 2012; 207 (3 Suppl): S3-8.
- Madinger NE, Greenspoon JS, Ellrodt AG. Pneumonia during pregnancy: has modern technology improved maternal and fetal outcome? *Am J Obstet Gynecol* 1989; 161: 657-62.
- Kourtis AP, Read JS, Jamieson DJ. Pregnancy and infection. *N Engl J Med* 2014; 370: 2211-8.
- Kwon JY, Romero R, Mor G. New insights into the relationship between viral infection and pregnancy complications. *Am J Reprod Immunol* 2014; 71: 387-90.
- Wong SF, Chow KM, Leung TN, et al. Pregnancy and perinatal outcomes of women with severe acute respiratory syndrome. *Am J Obstet Gynecol* 2004; 191: 292-7.
- Schwartz DA, Graham AL. Potential maternal and infant outcomes from (Wuhan) coronavirus 2019-nCoV infecting pregnant women: lessons from SARS, MERS, and other human coronavirus infections. *Viruses* 2020; 12: 194.
- Akhtar H, Patel C, Abuelgasim E, Harky A. COVID-19 (SARS-CoV-2) infection in pregnancy: a systematic review. *Gynecol Obstet Invest* 2020; 85: 295-306.
- Delahoy MJ, Whitaker M, O'Halloran A, et al. Characteristics and maternal and birth outcomes of hospitalized pregnant women with laboratory-confirmed COVID-19 – COVID-NET, 13 States, March 1–August 22, 2020. *MMWR Morb Mortal Wkly Rep* 2020; 69: 1347-54.
- Khan KS, Kunz R, Kleijnen J, Antes G. Five steps to conducting a systematic review. *J R Soc Med* 2003; 96: 118-21.
- Von Elm E, Altman DG, Egger M, et al. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007; 370: 1453-7.
- Lewin S, Glenton C, Munthe-Kaas H, et al. Using qualitative evidence in decision making for health and social interventions: an approach to assess confidence in findings from qualitative evidence syntheses (GRADE-CERQual). *PLoS Med* 2015; 12: e1001895.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009; 62: 1006-12.
- Hantoushzadeh S, Shamshirsaz AA, Aleyasin A, et al. Maternal death due to COVID-19 disease. *Am J Obstet Gynecol* 2020; 223: 109.e1-109.e16.
- Browne PC, Linfert JB, Perez-Jorge E. Successful treatment of preterm labor in association with acute COVID-19 infection. *Am J Perinatol* 2020; 37: 866-8.
- Baud D, Greub G, Favre G, et al. Second-trimester miscarriage in a pregnant woman with SARS-CoV-2 infection. *JAMA* 2020; 323: 2198-200.
- Richtmann R, Torloni MR, Oyamada Otani AR, et al. Fetal deaths in pregnancies with SARS-CoV-2 infection in Brazil: a case series. *Case Rep Womens Health* 2020; 27: e00243.
- Hosier H, Farhadian SF, Morotti RA, et al. SARS-CoV-2 infection of the placenta. *J Clin Invest* 2020; 130: 4947-53.
- García-Manau P, García-Ruiz I, Rodo C, et al. Fetal transient skin edema in two pregnant women with coronavirus disease 2019 (COVID-19). *Obstet Gynecol* 2020; 136: 1016-20.
- Lokken EM, Walker CL, Delaney S, et al. Clinical characteristics of 46 pregnant women with a severe acute respiratory syndrome coronavirus 2 infection in Washington State. *Am J Obstet Gynecol* 2020; 223: 911.e1-4.
- Wu Y, Liu C, Dong L, et al. Coronavirus disease 2019 among pregnant Chinese women: case series data on the safety of vaginal birth and breastfeeding. *BJOG* 2020; 127: 1109-15.

30. Yu N, Li W, Kang Q, Zeng W, Feng L, Wu J. No SARS-CoV-2 detected in amniotic fluid in mid-pregnancy. *Lancet Infect Dis* 2020; 20: 1364.
31. Liu Y, Chen H, Tang K, Guo Y. Clinical manifestations and outcome of SARS-CoV-2 infection during pregnancy. *J Infect* 2020. DOI: 10.1016/j.jinf.2020.02.028.
32. Wang J, Shu S, Zhang T, Zheng C. Chest CT findings in a pregnant woman in the second trimester with COVID-19 pneumonia. *Clin Imaging* 2021; 69: 266-8.
33. Panichaya P, Thaweerat W, Uthaisan J. Prolonged viral persistence in COVID-19 second trimester pregnant patient. *Eur J Obstet Gynecol Reprod Biol* 2020; 250: 263.
34. Inchingolo R, Smargiassi A, Moro F, et al. The diagnosis of pneumonia in a pregnant woman with coronavirus disease 2019 using maternal lung ultrasound. *Am J Obstet Gynecol* 2020; 223: 9-11.
35. Crovetto F, Crispi F, Lllurba E, Figueras F, Gómez-Roig MD, Gratacós E. Seroprevalence and presentation of SARS-CoV-2 in pregnancy. *Lancet* 2020; 396: 530-1.
36. Hachem R, Markou GA, Velupillai C, Poncelet C. Late miscarriage as a presenting manifestation of COVID-19. *Eur J Obstet Gynecol Reprod Biol* 2020; 252: 614.
37. Piersigilli F, Carkeek K, Hocq C, et al. COVID-19 in a 26-week preterm neonate. *Lancet Child Adolesc Health* 2020; 4: 476-8.
38. Cosma S, Borella F, Carosso A, et al. The “scar” of a pandemic: cumulative incidence of COVID-19 during the first trimester of pregnancy. *J Med Virol* 2021; 93: 537-40.
39. Lamazou F, Oger P, Dieli-Crimi R, et al. COVID-19 infection in first trimester of pregnancy marked by a liver cytolysis in a woman previously treated by hydroxychloroquine for repeated implantation failure: a case report. *BMC Infect Dis* 2020; 20: 845.
40. Tang J, Song W, Xu H, Wang N. No evidence for vertical transmission of SARS-CoV-2 in two neonates with mothers infected in the second trimester. *Infect Dis* 2020; 52: 913-6.
41. Hong L, Smith N, Keerthy M, et al. Severe COVID-19 infection in pregnancy requiring intubation without preterm delivery: a case report. *Case Rep Womens Health* 2020; 27: e00217.
42. Rubio Lorente AM, Pola Guillén M, López Jimenez N, et al. Study of amniotic fluid in pregnant women infected with SARS-CoV-2 in first and second trimester. Is there evidence of vertical transmission? *J Matern Fetal Neonatal Med* 2020. DOI: 10.1080/14767058.2020.1811669.
43. la Cour Freiesleben N, Egerup P, Hviid KVR, et al. SARS-CoV-2 in first trimester pregnancy: a cohort study. *Hum Rep* 2021; 36: 40-7.
44. Cosma S, Carosso AR, Cusato J, et al. Coronavirus disease 2019 and first-trimester spontaneous abortion: a case-control study of 225 pregnant patients. *Am J Obstet Gynecol* 2021; 224: 391.e1-391.e7.
45. Rasmussen SA, Smulian JC, Lednický JA, Wen TS, Jamieson DJ. Coronavirus disease 2019 (COVID-19) and pregnancy: what obstetricians need to know. *Am J Obstet Gynecol* 2020; 222: 415-26.
46. Alfaraj SH, Al-Tawfiq JA, Memish ZA. Middle East respiratory syndrome coronavirus (MERS-CoV) infection during pregnancy: report of two cases and review of the literature. *J Microbiol Immunol Infect* 2019; 52: 501-3.
47. Wang H, Li X, Li T, et al. The genetic sequence, origin, and diagnosis of SARS-CoV-2. *Eur J Clin Microbiol Infect Dis* 2020; 39: 1629-35.
48. Zhang L, Dong L, Ming L, et al. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection during late pregnancy: a report of 18 patients from Wuhan, China. *BMC Pregnancy Childbirth* 2020; 20: 394.
49. San-Juan R, Barbero P, Fernández-Ruiz M, et al. Incidence and clinical profiles of COVID-19 pneumonia in pregnant women: a single-centre cohort study from Spain. *EClinicalMedicine* 2020; 23: 100407.
50. Zaigham M, Andersson O. Maternal and perinatal outcomes with COVID-19: A systematic review of 108 pregnancies. *Acta Obstet Gynecol Scand* 2020; 99: 823-9.
51. Edwards MJ. Hyperthermia in utero due to maternal influenza is an environmental risk factor for schizophrenia. *Congenit Anom* 2007; 47: 84-9.
52. Ovali F. SARS-CoV-2 Infection and the Newborn. *Front Pediatr* 2020; 8: 294.
53. Kim CJ, Romero R, Chaemsaitong P, Kim JS. Chronic inflammation of the placenta: definition, classification, pathogenesis, and clinical significance. *Am J Obstet Gynecol* 2015; 213 (4 Suppl): S53-69.
54. Wu C, Yang W, Wu X, et al. Clinical manifestation and laboratory characteristics of SARS-CoV-2 infection in pregnant women. *Virol Sin* 2020; 35: 305-10.
55. Mongula JE, Frenken MWE, van Lijnschoten G, et al. COVID-19 during pregnancy: non-reassuring fetal heart rate, placental pathology and coagulopathy. *Ultrasound Obstet Gynecol* 2020; 56: 773-6.
56. Cribiù FM, Croci GA, Del Gobbo A, et al. Histological characterization of placenta in COVID19 pregnant women. *Eur J Obstet Gynecol Reprod Biol* 2020; 252: 619-21.
57. Ng WF, Wong SF, Lam A, et al. The placentas of patients with severe acute respiratory syndrome: a pathophysiological evaluation. *Pathology* 2006; 38: 210-8.
58. Gong X, Song L, Li H, et al. CT characteristics and diagnostic value of COVID-19 in pregnancy. *PLoS One* 2020; 15: e0235134.
59. Neumann-Podczaska A, Al-Saad SR, Karbowski LM, Chojnicki M, Tobis S, Wiczorowska-Tobis K. COVID 19 – clinical picture in the elderly population: a qualitative systematic review. *Aging Dis* 2020; 11: 988-1008.
60. Liu Q, Chen H, Zeng Q. Clinical characteristics of COVID-19 patients with complication of cardiac arrhythmia. *J Infect* 2020; 81: e6-e8.
61. Ai T, Yang Z, Hou H, et al. Correlation of chest CT and RT-PCR testing in coronavirus disease 2019 (COVID-19) in China: a report of 1014 cases. *Radiology* 2020; 296: E32-E40.
62. Kaplan YC, Ozsarfati J, Nickel C, Koren G. Reproductive outcomes following hydroxychloroquine use for autoimmune diseases: a systematic review and meta-analysis. *Br J Clin Pharmacol* 2016; 81: 835-48.
63. Favilli A, Mattei Gentili M, Raspa F, et al. Effectiveness and safety of available treatments for COVID-19 during pregnancy: a critical review. *J Matern Fetal Neonatal Med* 2020. DOI: 10.1080/14767058.2020.1774875.
64. Roberts SS, Martinez M, Covington DL, Rode RA, Pasley MV, Woodward WC. Lopinavir/ritonavir in pregnancy. *J Acquir Immune Defic Syndr* (1999) 2009; 51: 456-61.
65. Narang K, Enninga EAL, Gunaratne MDSK, et al. SARS-CoV-2 infection and COVID-19 during pregnancy: a multidisciplinary review. *Mayo Clin Proc* 2020; 95: 1750-65.