

# The role of the latency period in the preterm premature rupture of membranes: implications for treatment

Lin Lu, Jianhua Li, Bei Gan, Shan Zheng, Lihong Chen

Department of Obstetrics and Gynecology, The First Affiliated Hospital of Fujian Medical University, Fuzhou, China

Submitted: 3 May 2021, accepted: 24 June 2021

Online publication: 2 July 2021

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms/139314>

Copyright © 2021 Termedia & Banach

## Corresponding author:

Lihong Chen  
Department of  
Obstetrics and  
Gynecology  
The First Affiliated  
Hospital of Fujian  
Medical University  
20 Chazhong Road  
Taijiang District  
Fuzhou City  
Fujian Province  
Fuzhou 350005  
China  
Phone: 13645072616  
Fax: 0024 07898 1023  
E-mail:  
ucookyq0640226@163.com

## Abstract

**Introduction:** Conservative treatments with a latency period have been used for the treatment of preterm premature rupture of membranes (PPROM) in clinical practice. We aimed to evaluate the role and potential influencing factors of the latency period, to provide insights for the clinical treatment of PPRM.

**Material and methods:** PPRM pregnant women treated in our hospital from January 1, 2015 to September 30, 2020 were included. PPRM patients were divided into a 48–168 h group and a > 168 h latency group; the characteristics and prognosis of these two groups were compared and analyzed. Logistic regression analyses were conducted to analyze the relevant influencing factors of the latency period.

**Results:** A total of 131 PPRM patients were included. There were significant differences in the age, BMI, gestational age on admission, amniotic fluid volume before childbirth, and positive rate of cervical secretion culture between the two groups (all  $p < 0.05$ ). Logistic regression analyses indicated that the latency period was shorter in the PPRM patients with age  $\geq 30$  years (OR = 0.048, 95% CI: 0.121–0.863) and gestational age  $\geq 32$  weeks on admission (OR = 0.463, 95% CI: 0.069–0.811), and the latency period was prolonged in the PPRM patients with BMI  $\geq 23$  kg/m<sup>2</sup> (OR = 1.591, 95% CI: 1.134–1.944) and amniotic fluid volume  $\geq 6$  cm (OR = 2.129, 95% CI: 1.093–3.042) (all  $p < 0.05$ ). There were significant differences in the incidence of low birth weight and neonatal respiratory distress syndrome (NRDS) between the 48–168 h group and > 168 h group (all  $p < 0.05$ ).

**Conclusions:** The latency period plays an important role in PPRM, which is associated with the pregnant women's age, BMI, gestational week of rupture and amniotic fluid index.

**Key words:** premature rupture of membranes, preterm, latency period, care, treatment.

## Introduction

Premature rupture of membrane (PROM) refers to the spontaneous rupture of fetal membranes before labor [1]. According to the gestational age at the time of occurrence, PROM is divided into preterm premature rupture of membranes (PPROM) and full-term premature rupture of membranes. PPRM occurs when the pregnancy is more than 20 weeks and less than 37 weeks. According to statistics [2, 3], the incidence of PPRM is 2.0–3.5%, and 30–40% of preterm births are related to PPRM. PPRM can cause premature birth, placental abruption, oligohydran-

nios, umbilical cord prolapse, fetal distress and neonatal respiratory distress syndrome (NRDS), intraventricular hemorrhage (IVH), and necrotizing enterocolitis (NEC), leading to higher perinatal morbidity and mortality [4–6]. Therefore, taking correct treatment measures for PPRM patients is crucial to reduce the perinatal complications and improve the prognosis of newborns.

It has been reported that for PPRM patients, the lower the gestational age is, the higher are the infection rates of pregnant women and fetuses and perinatal mortality [7]. Previous studies [8, 9] have shown that prolonging the latency period for pregnant women with PPRM can reduce the infection rate of newborns, but it is difficult to attain a balance between prolonging the latency period and the timing of termination of pregnancy.

Therefore, we aimed to retrospectively evaluate the influencing factors of the latency period in pregnant women with PPRM, to provide insights for the clinical treatment and management of PPRM.

## Material and methods

### Ethical concerns

In this study, all methods were performed in accordance with the relevant guidelines and regulations. This study was checked and approved by the ethical committee of our hospital ([2016]005), all the included patients were well informed and written informed consent was obtained from patients.

### Patients

We selected PPRM pregnant women admitted to our hospital from January 1, 2015 to September 30, 2020 as the research population. The selection criteria of the patients in this study were as follows: 1) patients were admitted to our hospital within 12 h after the occurrence of PPRM; 2) the gestational age of patients was 28 to 33 weeks<sup>+</sup>6 on admission; 3) pathological examination of the placenta, fetal membranes, and umbilical cord after delivery was performed in our hospital. Exclusion criteria were as follows: 1) The latency period after admission was less than 48 h; 2) pregnant women with congenital abnormalities of the fetus; 3) patients who were unwilling to participate in this study. Patients with PPRM were divided into two groups according to the latency period after the occurrence of PPRM: the 48–168 h group and the > 168 h group.

### Treatment of PPRM

We routinely used dexamethasone 5 mg intramuscular injection, 1 time/12 h, 2 days to promote

fetal lung maturity. At the same time, second-generation cephalosporins or azithromycin was used to prevent infection. For patients with  $\geq 8$  uterine contractions within 60 min, uterine pressure  $\geq 60$  mm Hg, and uterine orifice dilation  $< 3$  cm, magnesium sulfate, ritodrine hydrochloride, or nifedipine would be used with the patient's consent. All patients were monitored for leukocytes, C-reactive protein (CRP), and fetal heart rate every 3 days, and B-ultrasound monitoring and cervical secretion culture were performed once a week.

### Diagnostic criteria

1) Oligo and oligohydramnios: B-mode ultrasound measurement of amniotic fluid index (AFI)  $\leq 8$  cm was considered as oligohydramnios; 2) intrauterine infection: maternal fever  $\geq 37.8^\circ\text{C}$ , accompanied by two or more of the following conditions: maternal pulse  $> 100$  beats/min, white blood cell count  $> 15 \times 10^9/\text{l}$ , fetal heart rate  $> 160$  beats/min, uterine tension and tenderness, smelly amniotic fluid, or positive amniotic cavity culture; 3) histology chorioamnionitis (HCA): inflammatory cell infiltration occurred in any tissue of chorionic membrane, amniotic membrane, and umbilical cord; 4) cervical secretion culture was positive. After PPRM occurred, we took a sterile swab to take the secretions from the cervix, and culture the bacteria, myochlamydia, fungi, anaerobic bacteria, and gonococcus in our laboratory. For any culture with a positive result it was considered that the cervical secretion culture was positive. All the included patients received ceftriaxone sodium 2 g *i.v.* before delivery.

### Data collection

Two investigators collected the following patients' information: age, body mass index (BMI), first pregnancy or not, the gestational age, white blood cells, amniotic fluid volume and CRP at the timepoint of admission and before childbirth. Additionally, we recorded the uterine contraction inhibitor use after 48 h, results of cervical secretion culture, the incidence of intrauterine infection and HCA.

### Statistical analysis

We used the SPSS 25.0 software package for statistical analysis. The measurement data were described using mean  $\pm$  standard deviation. The comparison between groups was performed by the two independent samples *t*-test or rank-sum test. The count data were represented by rate, and the comparison between groups was performed by the  $\chi^2$  test. With 0.05 as the level of selected variables and 0.10 as the level of excluded variables, forward step-by-step logistic

regression analysis was used to analyze relevant influencing factors.  $P < 0.05$  indicated that the difference between groups was statistically significant.

## Results

### Characteristics of included patients

A total of 131 PPROM patients were included in this study. As presented in Table I, compared with the  $> 168$  h group, there were significant differences in the age, BMI, gestational age on admission, amniotic fluid volume before childbirth, and positive rate of cervical secretion culture in the 48–168 h group (all  $p < 0.05$ ). There were no statistical differences in the first pregnancy, gestational age before childbirth, white blood cell, amniotic fluid volume on admission, CRP, uterine contraction inhibitor use after 48 h, incidence of intrauterine infection and HCA between the 48–168 h group and  $> 168$  h group (all  $p > 0.05$ ).

### Logistic regression analysis of the influencing factors of latency period

We used global optimization logistic regression, took the latency period time as the dependent variable, and all other observed indicators as independent variables, and we used the Hosmer-Lemeshow test to detect the goodness of fit of the model. It was found that the age of pregnant women, the gestational age on admission and the latency period were negatively correlated. Also BMI and amniotic fluid volume were positively correlated with the latency period. As presented in Table II, we used forward stepwise logistic regression to analyze the influencing factors. It was found that the latency period was shorter in the PPROM patients with age  $\geq 30$  years (OR = 0.048, 95% CI: 0.012–0.863) and gestational age  $\geq 32$  weeks on admission (OR = 0.463, 95% CI: 0.069–0.811), and the latency period was prolonged in the PPROM patients with BMI  $\geq 23$  kg/m<sup>2</sup> (OR =

**Table I.** Characteristics of included patients

Items	48–168 h group (n = 68)	> 168 h group (n = 63)	t/ $\chi^2$	P-value
Age [years]	30.81 $\pm$ 8.09	25.53 $\pm$ 7.16	1.024	0.027
BMI [kg/m <sup>2</sup> ]	24.72 $\pm$ 2.55	22.47 $\pm$ 1.94	1.217	0.014
First pregnancy	38 (55.88%)	35 (55.56%)	1.108	0.092
Gestational age [weeks]:				
On admission	31.13 $\pm$ 1.32	31.56 $\pm$ 1.60	1.133	0.031
Before childbirth	31.74 $\pm$ 1.24	31.89 $\pm$ 1.39	1.106	0.072
White blood cells [ $\times 10^9/l$ ]:				
On admission	12.38 $\pm$ 4.91	11.95 $\pm$ 4.14	1.118	0.097
Before childbirth	14.72 $\pm$ 5.13	14.68 $\pm$ 6.01	1.201	0.104
Amniotic fluid volume [cm]:				
On admission	7.63 $\pm$ 4.35	8.03 $\pm$ 5.23	1.186	0.092
Before childbirth	4.290 $\pm$ 2.31	5.66 $\pm$ 2.75	1.159	0.034
CRP [mg/l]:				
On admission	5.69 $\pm$ 4.11	6.04 $\pm$ 2.96	3.137	0.121
Before childbirth	24.08 $\pm$ 10.27	22.36 $\pm$ 8.05	6.223	0.076
Uterine contraction inhibitor use after 48 h	61 (89.71%)	50 (79.37%)	1.303	0.116
Positive cervical secretion culture	21 (30.88%)	24 (38.09%)	1.125	0.044
Intrauterine infection	7 (10.29%)	7 (11.11%)	1.203	0.089
HCA	56 (82.35%)	51 (80.95%)	1.167	0.105

**Table II.** Logistic regression analysis of the influencing factors of latency period in the PPROM patients

Variables	$\beta$	Wald	OR	95% CI	P-value
Age $\geq 30$ years	-0.153	2.127	0.048	0.012–0.863	0.042
Gestational age $\geq 32$ weeks on admission	-0.122	1.199	0.463	0.069–0.811	0.025
BMI $\geq 23$ kg/m <sup>2</sup>	0.113	2.037	1.591	1.134–1.944	0.038
Amniotic fluid volume $\geq 6$ cm	0.101	3.115	2.129	1.093–3.042	0.018

**Table III.** Comparison of prognosis of newborns between 48–168 h and > 168 h groups

Items	48–168 h group (n = 68)	> 168 h group (n = 63)	t/ $\chi^2$	P-value
Low birth weight	32 (47.06%)	36 (57.12%)	1.327	0.011
NRDS	12 (17.65%)	3 (4.76%)	1.105	0.043
Pneumonia	35 (51.47%)	33 (52.38%)	1.041	0.098
Sepsis	14 (20.59%)	11 (17.46%)	1.294	0.077
IVH	20 (29.41%)	20 (31.75%)	1.217	0.102
HIE	31 (45.59%)	29 (46.03%)	1.454	0.096
NEC	12 (17.65%)	11 (17.46%)	1.049	0.079

1.591, 95% CI: 1.134–1.944) and amniotic fluid volume  $\geq 6$  cm (OR = 2.129, 95% CI: 1.093–3.042) (all  $p < 0.05$ ).

### Prognosis of newborns

As indicated in Table III, there were significant differences in the incidence of low birth weight and NRDS between the 48–168 h group and > 168 h group (all  $p < 0.05$ ), and no significant differences in the incidence of pneumonia, sepsis, IVH, hypoxic-ischemic encephalopathy (HIE), or NEC were found (all  $p > 0.05$ ).

### Discussions

At present, the specific mechanisms of PROM are not fully understood. Studies [10–12] have shown that bacterial vaginitis, urinary tract infection, smoking during pregnancy, history of premature rupture of membranes, sexually transmitted diseases, abnormal cervical function, polyhydramnios, and invasive procedures are risk factors for premature rupture of membranes, among which genital tract infection has been considered to be the main cause of premature rupture of membranes. The hormone levels and immunity of pregnant women have changed, and their susceptibility to a variety of pathogens has increased, and they are susceptible to infection with these pathogens [13–15]. As an active immune site, the placenta can cause inflammation and an immune response under the action of pathogenic bacteria, and promote the production and release of a variety of inflammatory cytokines [16, 17]. The maternal immune system is activated in this environment, resulting in damage to the placenta, causing adverse pregnancy outcomes such as premature birth [18]. Studies [19, 20] have found that the detection rate of bacteria in the amniotic fluid for PROM reaches 30%, and another 70% of patients may also be infected by the living maternal tissues of amniotic fluid. Therefore, the early diagnosis and timely treatment are vital to the prognosis of PROM patients.

We have found that the earlier the gestational week of the PPROM is, the longer is the latency

period, which is consistent with the results of previous studies [21, 22], and the pregnant women whose membrane rupture occurs before 32 weeks have a greater probability of a latency period > 168 h. It has been reported [23] that expecting a time longer than 1 week does not increase the risk of HCA, so pregnant women who have ruptured membranes before 32 weeks should keep the latency period more than one week. There are still many controversies about the relationship between the number of pregnancies, age and latency period of pregnant women. The results of this study show that the patient's parity is not a risk factor that affects the latency period, which is consistent with the results of previous studies [24, 25]. It has been reported that gestational age is closely associated with preterm premature rupture of membranes [26, 27]. Pregnant women over the age of 30 have an increased probability of shortening the latency period. It is estimated that different medical institutions have biases in the selection of the latency period. Yet the specific latency period requires large-scale, multi-centered research in the future.

In our study, the amniotic fluid index of the two groups of patients is not statistically significantly different at admission, but during the latency period, when the amniotic fluid index is above 6 cm, the possibility of a latency period > 168 h increases accordingly. The possible reasons may be as follows: rupture of the membrane leads to the fetal head getting closer to the lower part of the uterus, which has a mechanical expansion effect on it, causing the start of labor [28]. Also, the fetus gets closer to the uterine wall, the umbilical cord is compressed, the incidence of fetal distress increases, and the termination of pregnancy is early [29–31]. The amniotic fluid volume of pregnant women is closely related to infection. Infection places the fetus in a state of stress, reduces blood flow in the kidneys and urine output, and reduces amniotic fluid volume [9]. Amniotic fluid contains a variety of anti-infective cytokines, and the loss of amniotic fluid weakens the capacity for resistance to infection, thereby aggravating intrauterine infections, which can easily induce contractions [32,

33]. Therefore, for pregnant women with PPROM with progressive reduction in amniotic fluid, even if there is no clinical evidence of infection, they must be highly vigilant for the occurrence of chorioamnionitis [34].

Previous studies have reported [35–37] that for PPROM patients who have fetal lung maturity and infection, simply extending the gestational period will not reduce the serious complications of preterm infants, but PPROM pregnant women have reduced amniotic fluid, higher incidence of chorioamnionitis and various inflammations in the amniotic fluid. The increase in substances can affect the growth and development of the fetus, causing the fetal body mass not to increase correspondingly with the increase of the gestational age. The more the gestational age increases, the more the fetal body mass lags behind [38]. We have found that extending the gestational age does not reduce the incidence of serious complications of preterm infants and maternal infectious diseases, and the use of uterine contraction inhibitors does not help to extend the gestational age. Therefore, for PPROM patients, the focus should be on predicting and treating infections. For patients with increased WBC, progressive increase in CRP, and progressive decrease in amniotic fluid, even if intrauterine infection cannot be diagnosed clinically, a comprehensive assessment should be combined with the specific conditions of the patient. It has been reported that intrauterine infection is thought to be one cause of PPROM, and use of antibiotic regimens is beneficial for the control of intrauterine infection and reduction of infant morbidity [39, 40]. Still, the timing and dose of antibiotic regimens in the PPROM require further investigations.

In conclusion, we have found that for PPROM patients, pregnant women's age, BMI, gestational week of rupture and amniotic fluid index are associated with the latency period. With a longer latency period, the incidence of NRDS is lower, but the incidence of low birth weight is higher. As this study is limited by sample size and study design, future studies with a larger sample size and prospective design are needed to further elucidate the potential influencing factors of the latency period, to provide evidence for the management of PPROM.

### Acknowledgments

Lin Lu and Jianhua Li – equal contributor.

This study has been funded by Industry-University-Research Project of Fujian Provincial Department of Education (JAT160189). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Conflict of interest

The authors declare no conflict of interest.

### References

1. Areia AL, Areia M, Mota-Pinto A. Procalcitonin in preterm rupture of membranes: a systematic review and meta-analysis. *Arch Gynecol Obstet* 2021; 303: 917-24.
2. Volpe N, di Pasquo E, Ferretti A, et al. Hyperechoic amniotic membranes in patients with preterm premature rupture of membranes (p-PPROM) and pregnancy outcome. *J Perinat Med* 2021; 49: 311-8.
3. Niyaty S, Moghaddam-Banaem L, Sourinejad H, Mokhlesi S. Are maternal metabolic syndrome and lipid profile associated with preterm delivery and preterm premature rupture of membranes? *Arch Gynecol Obstet* 2021; 303: 113-9.
4. Shiqiao H, Bei X, Yudi G, Lei J. Assisted reproductive technology is associated with premature rupture of membranes. *J Matern Fetal Neonatal Med* 2021; 34: 555-61.
5. Maged AM, Kamel HH, Sanad AS, et al. The value of amniopatch in pregnancies associated with spontaneous preterm premature rupture of fetal membranes: a randomized controlled trial. *J Matern Fetal Neonatal Med* 2021; 34: 267-73.
6. Pierson RC, Gordon SS, Haas DM. A retrospective comparison of antibiotic regimens for preterm premature rupture of membranes. *Obstet Gynecol* 2014; 124: 515-9.
7. Huang CC, Pan SC, Chin WS, et al. Maternal proximity to petrochemical industrial parks and risk of premature rupture of membranes. *Environ Res* 2021; 194: 110688.
8. Zhang SE, Chen XY, Chen C, Qiu XM, Lin BC, Yang CZ. Influence of premature rupture of membranes on the early prognosis of extremely premature infants. *Zhongguo Dang Dai Er Ke Za Zhi* 2021; 23: 25-30.
9. Kole-White MB, Nelson LA, Lord M, et al. Pregnancy latency after preterm premature rupture of membranes: oral versus intravenous antibiotics. *Am J Obstet Gynecol MFM* 2021; 3: 100333.
10. Rouzaire M, Corvaisier M, Roumeau V, et al. Predictors of short latency period exceeding 48 h after preterm premature rupture of membranes. *J Clin Med* 2021; 10: 150.
11. Liu Y, Wu M, Song L, et al. Association between prenatal rare earth elements exposure and premature rupture of membranes: results from a birth cohort study. *Environ Res* 2021; 193: 110534.
12. Suff N, Kunitsyna M, Shennan A, Chandiramani M. Optimal timing of cervical cerclage removal following preterm premature rupture of membranes: a retrospective analysis. *Eur J Obstet Gynecol Reprod Biol* 2021; 259: 75-80.
13. Wolfensberger A, Zimmermann R, von Mandach U. Neonatal mortality and morbidity after aggressive long-term tocolysis for preterm premature rupture of the membranes. *Fetal Diagn Ther* 2006; 21: 366-73.
14. Medina TM, Hill DA. Preterm premature rupture of membranes: diagnosis and management. *Am Fam Physician* 2006; 73: 659-64.
15. Sae-Lin P, Wanitpongpan P. Incidence and risk factors of preterm premature rupture of membranes in singleton pregnancies at Siriraj Hospital. *J Obstet Gynaecol Res* 2019; 45: 573-7.
16. Mehra S, Amon E, Hopkins S, Gavard JA, Shyken J. Transvaginal cervical length and amniotic fluid index: can it predict delivery latency following preterm premature

- rupture of membranes? *Am J Obstet Gynecol* 2015; 212: 400e1-9.
17. Tchirikov M, Schlalbritz-Loutsevitch N, Maher J, et al. Mid-trimester preterm premature rupture of membranes (PPROM): etiology, diagnosis, classification, international recommendations of treatment options and outcome. *J Perinat Med* 2018; 46: 465-88.
  18. Schmitz T, Sentilhes L, Lortie E, et al. Preterm premature rupture of the membranes: Guidelines for clinical practice from the French College of Gynaecologists and Obstetricians (CNGOF). *Eur J Obstet Gynecol Reprod Biol* 2019; 236: 1-6.
  19. Sim WH, Ng H, Sheehan P. Maternal and neonatal outcomes following expectant management of preterm prelabor rupture of membranes before viability. *J Matern Fetal Neonatal Med* 2020; 33: 533-41.
  20. Guckert M, Clouqueur E, Drumez E, et al. Is homecare management associated with longer latency in preterm premature rupture of membranes? *Arch Gynecol Obstet* 2020; 301: 61-7.
  21. Musilova I, Andrys C, Hornychova H, et al. Gastric fluid used to assess changes during the latency period in preterm prelabor rupture of membranes. *Pediatr Res* 2018; 84: 240-7.
  22. Yagur Y, Weitzner O, Ravid E, Biron-Shental T. Can we predict preterm delivery in patients with premature rupture of membranes? *Arch Gynecol Obstet* 2019; 300: 615-21.
  23. Mei-Dan E, Hutchison Z, Osmond M, Pakenham S, Ng E, Green J, Nevo O. Preterm premature rupture of membranes in twins: comparison of rupture in the presenting versus non-presenting sac. *J Obstet Gynaecol Can* 2020; 42: 163-8.
  24. Phupong V, Kulmala L. Factors associated with latency period in preterm prelabor rupture of membranes. *J Matern Fetal Neonatal Med* 2016; 29: 2650-3.
  25. Ilhan N, Aygun BK, Gungor H. The relationship between the latency period, infection markers, and oxidant and antioxidant states in women with preterm premature rupture of membranes. *Ir J Med Sci* 2017; 186: 965-70.
  26. Gibson KS, Brackney K. Perivable premature rupture of membranes. *Obstet Gynecol Clin North Am* 2020; 47: 633-51.
  27. Swiatkowska-Freund M, Traczyk-Los A, Partyka A, Obara K, Damdinsuren A, Preis K. Perinatal outcome in preterm premature rupture of membranes before 37 weeks of gestation. *Ginekol Pol* 2019; 90: 645-50.
  28. Snowise S, Mann LK, Moise KJ Jr, Johnson A, Bebbington MW, Papanna R. Preterm prelabor rupture of membranes after fetoscopic laser surgery for twin-twin transfusion syndrome. *Ultrasound Obstet Gynecol* 2017; 49: 607-11.
  29. Workneh Y, Gultie T. Latency period and early initiation of breastfeeding in term premature rupture of membrane in Southern Ethiopia, 2017. *Ital J Pediatr* 2019; 45: 70.
  30. Schreiber H, Shitrit P, Man-El G, Engel O, Berkovitz A, Biron-Shental T. Macrolide antibiotics roxithromycin vs. azithromycin for preterm premature rupture of membranes: a retrospective comparison. *Arch Gynecol Obstet* 2019; 300: 569-73.
  31. Choi EK, Kim SY, Heo JM, et al. Perinatal outcomes associated with latency in late preterm premature rupture of membranes. *Int J Environ Res Public Health* 2021; 18: 672.
  32. Di Sarno R, Raffone A, Saccone G. Effects of progestogens in women with preterm premature rupture of membranes. *Minerva Ginecol* 2019; 71: 121-4.
  33. Walker MW, Picklesimer AH, Clark RH, Spitzer AR, Garite TJ. Impact of duration of rupture of membranes on outcomes of premature infants. *J Perinatol* 2014; 34: 669-72.
  34. Sorano S, Fukuoka M, Kawakami K, Momohara Y. Prognosis of preterm premature rupture of membranes between 20 and 24 weeks of gestation: a retrospective cohort study. *Eur J Obstet Gynecol Reprod Biol X* 2020; 5: 100102.
  35. Test G, Levy A, Wiznitzer A, et al. Factors affecting the latency period in patients with preterm premature rupture of membranes. *Arch Gynecol Obstet* 2011; 283: 707-10.
  36. Galletta MAK, Bittar RE, Agra I, Guerra ECL, Francisco RPV, Zugaib M. Epidemiological profile of patients with preterm premature rupture of membranes at a tertiary hospital in Sao Paulo, Brazil. *Clinics* 2019; 74: e1231.
  37. Catt E, Chadha R, Tang S, Palmquist E, Lange I. Management of preterm premature rupture of membranes: a comparison of inpatient and outpatient care. *J Obstet Gynaecol Can* 2016; 38: 433-40.
  38. Kacerovsky M, Romero R, Stepan M, et al. Antibiotic administration reduces the rate of intraamniotic inflammation in preterm prelabor rupture of the membranes. *Am J Obstet Gynecol* 2020; 223: 114e111-20.
  39. Mercer BM, Miodovnik M, Thurnau GR, et al. Antibiotic therapy for reduction of infant morbidity after preterm premature rupture of the membranes. A randomized controlled trial. National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. *JAMA* 1997; 278: 989-95.
  40. Martingano D, Singh S, Mitrofanova A. Azithromycin in the treatment of preterm prelabor rupture of membranes demonstrates a lower risk of chorioamnionitis and postpartum endometritis with an equivalent latency period compared with erythromycin antibiotic regimens. *Infect Dis Obstet Gynecol* 2020; 2020: 2093530.