

Analysis of 622 pediatric hospitalizations due to arterial ischemic stroke in Poland – National Health Fund registry-based study from 2011 to 2020

Keywords

registry, hospitalisation, paediatrics, arterial ischaemic stroke

Abstract

Introduction

The present study aimed to evaluate the prevalence of the arterial ischaemic stroke (AIS) in Polish children, as well as to analyse the parameters related to AIS hospitalization, including age, gender, region, month and season of a year at admission, duration and costs based on data from National Health Fund (NHF) registry in 2011–2020.

Material and methods

Data from the NHF were analysed statistically. The disease was identified according to the codes I63 and I64 of the International Classification of Diseases, Tenth Revision (ICD-10), and patients included only individuals up to 18 years of age. Results We identified 622 hospitalisations due to paediatric AIS in Poland in the study period.

Results

The most frequent age subgroups were adolescents, followed by toddlers or pre-school children (34.73% and 24.12%, respectively), while the least frequent – neonates or infants (9.81%). ICD-10 procedures did significantly affect the duration and costs of hospitalisation ($p < 0.001$). The highest costs of hospitalisations concerned the I63.1 procedure (cerebral infarction due to embolism of precerebral arteries), which included thrombectomy. Duration and costs of hospitalisation were positively correlated to each other ($r = 0.525$, $p < 0.001$). Age correlated negatively with duration of hospitalisation ($r = -0.154$, $p < 0.001$) and positively with costs of hospitalisation ($r = 0.133$, $p = 0.008$).

Conclusions

Data from the NHF registry proved that AIS occurs more often in boys than girls and is more common in adolescents (15–18 years) than younger children.

Explanation letter

Dear Prof. Maciej Banach, MD, PhD, FAHA, FESC, FNLA
Editor-in-Chief,
Archives of Medical Science

I kindly inform you that a rewritten (according to the Reviewers' comments) version of the manuscript has just been uploaded to the editorial board. We also added citations from the AMS papers - one of them had been in the bibliography before, another one appeared during the revision. We hope it meets the expectations of both the reviewers and the magazine's editors.
Kindest regards,

[Analysis of 622 pediatric hospitalizations due to arterial ischemic stroke in Poland second revision 06062022.docx](#)

**Analysis of 622 pediatric hospitalizations due to arterial ischemic stroke in Poland –
National Health Fund registry-based study from 2011 to 2020**

Preprint

Abstract

Introduction: The present study aimed to evaluate the prevalence of the arterial ischaemic stroke (AIS) in Polish children, as well as to analyse the parameters related to AIS hospitalization, including age, gender, region, month and season of a year at admission, duration and costs based on data from National Health Fund (NHF) registry in 2011–2020.

Materials and methods Data from the NHF were analysed statistically. The disease was identified according to the codes I63 and I64 of the International Classification of Diseases, Tenth Revision (ICD-10), and patients included only individuals up to 18 years of age.

Results We identified 622 hospitalisations due to paediatric AIS in Poland in the study period.

Results: The most frequent age subgroups were adolescents, followed by toddlers or pre-school children (34.73% and 24.12%, respectively), while the least frequent – neonates or infants (9.81%). ICD-10 procedures did significantly affect the duration and costs of hospitalisation ($p < 0.001$). The highest costs of hospitalisations concerned the I63.1 procedure (cerebral infarction due to embolism of precerebral arteries), which included thrombectomy. Duration and costs of hospitalisation were positively correlated to each other ($r = 0.525$, $p < 0.001$). Age correlated negatively with duration of hospitalisation ($r = -0.154$, $p < 0.001$) and positively with costs of hospitalisation ($r = 0.133$, $p = 0.008$).

Conclusion Data from the NHF registry proved that AIS occurs more often in boys than girls and is more common in adolescents (15–18 years) than younger children.

Keywords: arterial ischaemic stroke; hospitalisation, paediatrics; registry

Introduction

Arterial ischemic stroke is a set of rapidly developing clinical symptoms characterized by a sudden, focal or generalized brain disorder resulting from the dysfunction of brain circulation, lasting 24 hours or longer or leading to death. The frequency of ischemic stroke in children is relatively low in contrast to adults. One of the latest data demonstrated that in developed countries prevalence of both ischemic and hemorrhagic stroke ranges from 3 to 25 cases per 100 000 children while in newborns this frequency is higher, i.e. 1 in 4000 live births [1]. The study from Taiwan reported that 2-year-period prevalence for ischemic lesions in children was 5.2 per 100 000 per year [2]. Recent data from Germany showed an overall incidence of 0.41 per 100,000 per year [3]. In Switzerland, the incidence of posterior AIS in children was established at 0.183 per 100 000 (i.e. 16% of all childhood AIS) [4].

AIS causes deleterious neurological consequences that affect the daily functioning of the patient as well as the costs of medical care and rehabilitation both in adults and in children. The most frequent consequences of AIS in children are motor disorders, mainly hemiparesis, epileptic seizures, aphasia or intellectual delay [5-7]. If AIS occurs in the perinatal period, it may have an impact on cognitive and behavioral deficiencies appearing in the future life [8]. In addition, pediatric patients are at a high risk of recurrent stroke and post-stroke mortality [5,9]. In the USA, cerebrovascular disease was found among ten of the most common causes of death in pediatric population [10].

Valuable sources of strategic information in regard to arterial ischemic stroke in adults worldwide are the databases managed by public health institutions [11-13]. In contrast, analysis of data on AIS in pediatric population coming from national registries are rather scarce [2].

Registry-based data show many advantages, i.e. they are already collected, and the researchers do not need to organize and finance data collection from individual patients or

search medical records in their hospitals. In addition, they can serve as a sampling frame or a reference for researchers who would like to select a sample or evaluate the representativeness of their own sample. According to the definition of the Statistical Office of the European Union EUROSTAT, these data are of good quality, which is understood broadly as relevance, accuracy, timeliness and punctuality, accessibility and clarity, coherence and comparability [14]. Definitions of variables and domains are already fixed, unambiguous and comprehensive. There are no sampling errors and missing data; data are available immediately after inclusion, including long comparable time series, comparability over time and between domains, coherence with other health statistics.

In Poland, the National Health Fund (NHF) registry is the only public payer for healthcare services which covers the whole country, i.e. 16 administrative regions. The NHF has the right to process the personal data of the patients, with an emphasis on monitoring their health as well as their needs for health services, drugs and medical devices [15]. As for now, no special database concerning pediatric AIS was established in Poland thus Polish NHF is considered as a reliable source of epidemiological data from all over the country. In consequence, such data may enable real-time monitoring of the health status of patients: comorbidity, disease burden, as well as therapeutic patterns. Analyzes based on such comprehensive data will help in defining health priorities, and specially help in appropriate assignation of public funds. Full-scale data on pediatric AIS may provide alternative novel sight into the epidemiological status of AIS in children.

This study aimed to evaluate the prevalence of childhood AIS, as well as the parameters related to AIS hospitalization, including age, gender, region, month and season of the year at admission, duration and costs based on data from NHF registry within a period from 2011 to 2020.

Methods

Study population

For this retrospective study, we included individual, anonymous data from the NHF concerning the hospitalisation of children with AIS registered in Poland in 2011–2020. Data were obtained based on the formal approval from the NHF to implement the “Maps of Health Needs” project.

Inclusion criteria were as follows: hospitalisation due to ICD-10 I63.0–I63.9. We also included patients hospitalized due to I64, which included arterial ischaemic, haemorrhagic and other strokes; age from 0 to 18 years. This inclusion was done after the discussion with the clinically experienced pediatric neurologist.

In the analysed group of children with AIS hospitalised in Poland during 2011–2020, five subgroups according to age were recognised: a) neonates or infants (from birth to the end of 12th month of life), b) toddlers and pre-school children (from the 1st year of life to the end of 6th year of life), c) pre-pubertal children (7th to 10th year of life), d) early pubertal patients (11th to 14th year of life) and e) adolescents (aged 15–18 years).

The patients' consent as well as the formal approval of the Local Ethics Committee were not required because database analysis is not treated in Poland as medical experiment according to Polish law (Act of 5 December 1996 on professions of physicians and dentist (Journal of Laws of 2021, item 790 as amended)). The present study was retrospective and non-invasive as we did not have direct contact with patients during the research and data obtained from the NHF were anonymous.

Hospitalization-related parameters

For each hospitalisation, the following data were available from the NHF database: regional branch of the NHF in which the patient was hospitalised, regional branch of the NHF in which

the patient was registered, age, gender, group of diseases according to the International Classification of Diseases, Tenth Revision (ICD-10) classification, date and type of admission, date and type of discharge, and total cost of the procedure provided were rated by the NHF (amount in Polish zloty). Hospitalisation costs were expressed in EUR and USD during the analysis according to the exchange rate as of 31 December 2020. Based on the registry data, we calculated the following: duration of hospitalisation, month and season of the year at admission, and hospitalisation outside the region of residence.

Statistical analysis

The data were analysed using STATISTICA 13 software (STATSOFT, Tulsa, OK, USA) and R language (version R-3.6.2).

The mean and standard deviation values were estimated for numerical variables and absolute numbers (n) and percentages (%) of the occurrence of items for categorical variables.

The following statistical tests were used:

- Fisher's exact test to compare categorical variables between ICD-10 procedures or year of hospitalisation;
- F test of variance to compare age, duration or costs of hospitalisation between ICD-10 procedures or year of hospitalisation;
- Pearson's correlation coefficient to find correlations between age, duration and costs of hospitalisation.

In analyses of dependent variables against ICD-10 procedures, I63.1 was omitted because it was registered only for one patient. The significance level was 0.05 for all statistical tests.

Results

Diagnosis of AIS in paediatric patients

A total of 622 hospitalisations due to paediatric AIS was registered in Poland in 2011–2020 (Table 1). In all analysed years, stroke not specified as haemorrhage or infarction was the most frequent (31.67%), followed by cerebral infarction, unspecified (18.33%), cerebral infarction due to unspecified occlusion or stenosis of cerebral arteries (15.27%) and other cerebral infarction (14.79%). Furthermore, the ICD-10 procedure of hospitalisations due to AIS did not significantly differ in the analysed period ($p=0.410$).

Gender, age, season of the year and month at admission to hospitals due to paediatric AIS

AIS occurred more common in boys (367, i.e. 59.00%) than girls (255, i.e. 41.00%).

However, the gender of patients hospitalised due to AIS ($p=0.116$) and the ICD-10 procedure ($p=0.891$) did not differ significantly in the analysed period.

The mean age of AIS children hospitalised between 2011 and 2020 in Poland was 10.2 ± 6.0 years. The age of patients hospitalised due to AIS ($p=0.454$) and the ICD-10 procedure ($p=0.201$) did not differ significantly in the analysed period.

Age distribution in the whole group is shown in Figure 1. The most common age subgroups were adolescents (34.73%), followed by toddlers or pre-school children (24.12%), while the least common was the group of neonates or infants (9.81%).

Patients suffered from the disease more common in autumn than in other seasons of the year, while October was a dominant month for the illness (Figure 2). However, month and season of the year at admission and the ICD-10 procedure did not differ significantly in the analysed years (month at admission versus year of hospitalisation $p=0.641$, season of the year at admission versus year of hospitalisation $p=0.536$, month at admission versus ICD-10 procedure $p=0.935$, season of the year at admission versus ICD-10 procedure $p=0.935$).

Region of residence of paediatric patients hospitalised due to AIS

A significant difference was found between the region of residence of paediatric patients with AIS and the year of hospitalisation (Table 2). In the analysis of total years, the highest percentage of paediatric hospitalisations due to AIS took places in Mazowieckie (11.58%), Śląskie (9.97%), Dolnośląskie (9.65%), Wielkopolskie (9.49) and Małopolskie (9.16%), whereas number of hospitalisations due to paediatric AIS were the lowest in Opolskie (2.73%), Lubuskie (3.38%), Lubelskie (3.54%) and Zachodniopomorskie (3.54%).

A total of 57 cases out of 622, i.e. 9.16%, were hospitalised outside the region of residence. Hospitalisation outside the region of residence did not differ significantly in the analysed years ($p=0.286$).

Types of admission and discharge, duration and costs of hospitalisation due to paediatric AIS

The most common type of admission was emergency admission – other cases, while the least common – transfer from another hospital (59.81% and 5.31, respectively) (Table 3).

In case of discharge type, the most common was a referral for further outpatient treatment (47.43%), while the least common discharge was a referral for further treatment in a medical facility other than a hospital, providing medical services such as inpatient and round-the-clock health services (0.80%) (Table 4).

The duration and costs of hospitalisation did not differ significantly in the analysed years ($p = 0.495$ and $p = 0.495$, respectively). Therefore, we analysed the duration and costs of hospitalisation in the entire period together (Table 5). ICD-10 procedures significantly differentiated the duration and costs of hospitalisation ($p<0.001$ and $p<0.001$, respectively).

The most extended hospitalisation was observed for I63.0 and I63.5 procedures. The highest costs of hospitalisations concerned the I63.1 procedure; however, only one child was treated within this procedure, which included thrombectomy. Next, high costs were linked with the I63.4 procedure, while the lowest applied to the I64.

The duration and costs of hospitalisation were positively correlated to each other ($r = 0.525$, $p < 0.001$). The longer the hospitalisation, the higher the average costs of hospitalisation. Age correlated negatively with duration of hospitalisation ($r = -0.154$, $p < 0.001$), whereas there was a positive correlation between age and costs of hospitalisation ($r = 0.133$, $p = 0.008$). The older the patient, the shorter the average duration of the hospitalisation.

Discussion

In the present study, adolescents aged 15–18 years were the most numerous group of patients. The disease occurred less commonly in neonates or infants (from birth to 1st year of life), i.e. in 10%. The frequency of adolescents is in accordance to previously published data on children with AIS diagnosed in Katowice (Southern Poland) [16]. In turn, such a low number of neonates or infants in the present study contradicts the previous data [17]; however, it may result from the fact that AIS in newborns could be classified as other brain diseases using different coding, especially in neonatal departments, because in these cases AIS results from a specific cause and may be scrambled. In the present registry analysis, four patients (i.e. 0.64%) were treated with thrombolysis. They were adolescents aged 17–18 years. An earlier study on American children demonstrated a similar frequency (1.6%) of patients receiving thrombolytic therapy. The patients were only males with a mean age of 11.1 years [18]. Systematic intravenous thrombolysis and endovascular thrombectomy are suggested as potentially effective treatment options in children and adolescents in addition to antiplatelet medication and heparinisation [19].

In the present study, we demonstrated that the experience of thrombolytic therapy in children is very modest, as it was only recently possible to use intravenous thrombolysis in patients aged 16 years and older, and earlier this treatment was only approved in patients over 18 years of age. Since the drug is approved for children > 16 years of age, spreading the benefits and

risks, will not overstate the cost of hospitalization. In the adult population, where stroke is a very widespread problem, IV thrombolysis became the standard treatment for the acute phase of stroke for several years. Given the frequency of stroke in children over 16 years of age, the time needed to gain experience and describe the groups of children treated with thrombolysis will probably be greater.

In the total group, boys with stroke were more common than girls (59% vs 41%). Most researchers also describe a higher incidence of stroke in boys, which relates to both paediatric stroke (risk 1.25) and perinatal stroke (risk 2.0), and this difference remains despite the elimination of trauma as the cause of stroke [20,21]. Earlier study based on Polish children demonstrated that boys were more prevalent in all age subgroups of analysed patients with AIS (i.e. infants–toddlers, children and adolescents). Boys significantly outnumbered girls in the POCI and TACI stroke subgroups, while girls predominated in the LACI stroke subgroup [16]. In the group of 1,979 children with stroke, both ischaemic and haemorrhagic (aged <10 years and from 10 to 19 years) analysed in the study from Taiwan based on the database from the National Health Insurance, 60.2% were boys [22]. An extremely high percentage of boys with AIS (71%) was observed in the recent Chinese study [23]. Similarly, in Danish research, 67% of children with AIS and varicella infection less than 12 months before the onset of symptoms were boys [24]. The explanation of the reason why boys suffer from AIS more often is not clear. This may be due to the higher incidence of infections and head injuries in boys compared to girls, or simply because of the greater number of boys in the general population [25].

Stroke in Polish children occurred most frequently in autumn. This may be due to the higher incidence of infections in this season of the year. The available data demonstrated that infections increased the risk of paediatric stroke [26]. In the Canadian study by Carey et al. [27], infection was found to be a risk factor for stroke in every fifth child with posterior AIS.

The authors observed among others pharyngitis, meningitis, otitis media or history of chicken pox in last 12 months prior to stroke. In turn, in children with AIS from Senegal three patients had meningitis, which was assigned to *Cryptococcus neoformans* (one case) and *Mycobacterium tuberculosis* (two cases) which then revealed a stroke [28]. The general effects of infection lead to temporary thrombophilia, and hypercoagulable states are considered a risk factor for AIS in younger patients [29]. Prothrombotic abnormalities also play a critical role in the aetiology of paediatric AIS [30]. In the study by Cao et al. [23], over 73% of analysed stroke children aged 1 month to 4 years had respiratory tract infection with fever, and 33% had thrombophilia.

In patients from Katowice (Poland), AIS appeared most commonly in winter with the significant differences between types of AIS and seasons of the year at which the AIS occurred, i.e. TACI stroke occurred most frequently in the summer, LACI – during winter or spring, and PACI – winter or autumn [31].

The mean length of hospital stay for children with AIS was 12.9 days. Similarly, Cao et al. [23] reported that children with AIS were hospitalised for 5–20 days, with 13 days on average. Regarding the length of stay of children in hospital - usually more than 5 days is the time which is necessary for all laboratory tests to be carried out to analyse the cause of the stroke. The length and costs of hospitalisation did not differ significantly in the analysed years. This information can be used by the Agency for Health Technology Assessment and Tariff System that prepares and verifies health technology assessments, with particular emphasis on those technologies that are financed from public funds; it also prepares recommendations for the Minister of Health regarding medical technologies (drug and non-drug) in Poland. Information on the length of stay can affect the revision of the cost of treatment by the above Agency and the change in NHF tariffs.

Many factors influence the hospitalisation costs of adult patients, including stroke type and severity and the length of hospital stay [32]. In the present registry-based analysis, the duration and costs of hospitalisation due to paediatric stroke did not differ significantly in the analysed years. The ICD-10 procedures affected the duration and costs of hospitalisation. The most expensive procedure according to ICD-10 was I63.1; however, it was applied in the case of only one child. The least expensive procedure was I64. The mean cost of paediatric stroke in Poland in 2011–2020 was USD 891. Perkins et al. [32] reported the mean cost of ischaemic stroke of USD 20,927. The duration of hospitalisation correlated positively with costs of hospitalisation ($r = 0.525$, $p < 0.001$). The type of ICD-10 procedure affected the length of hospitalisation and its costs significantly. The most expensive procedures were related to cerebral infarction due to embolism of precerebral arteries (I63.1), cerebral infarction due to embolism of cerebral arteries (I63.4), cerebral infarction due to unspecified occlusion or stenosis of precerebral arteries (I63.2) and other cerebral infarction (I63.8) procedures, while the lowest costs applied to cerebral infarction due to thrombosis of precerebral arteries (I64) and stroke, not specified as haemorrhage or infarction (I63.0) procedures.

Previously, data from Taiwan registry [22] demonstrated that the hospitalisation care costs for stroke are higher in children than adults as the proportion of haemorrhagic cases is greater in paediatric populations. The authors reported that the mean inpatient costs, including both first and subsequent hospitalizations were higher for patients <10 years and aged 10–19 years (USD 3,565 per case) than for adult patients with stroke (USD 1,933) [22]. In addition, the costs were found to be higher for the recurrent cases. Chen et al. [22] demonstrated also that costs of hospitalization due to ischemic stroke were comparable for children <10 years and those between 10 and 19 years (approximately USD 2400). In our analysis, mean costs of AIS hospitalization were significantly lower and ranged from USD 482 to USD 3410 depending on ICD-10 procedure with mean USD 891 per case.

The present study has some limitations. First of all, the number of cases is relatively low. In a Taiwan study concerning a comparable period, almost 2,000 children with AIS were identified [22]. Therefore, we may infer that AIS belong to rare diseases in Poland. Second, low number of children with AIS in smaller (population wise) regions (Lubelskie, Lubuskie, Opolskie, Warmińsko-Mazurskie and Zachodniopomorskie) may be alarming. This may indicate the weaknesses of the health service and failure to detect the disease, especially in the Opolskie and Lubuskie regions. Third, NHF data include only hospitalisations financed by public funds. In Poland, hospitalisation costs are covered by public funds; however, they may also be paid directly by patients. Therefore, AIS in children is unlikely to be treated privately. In addition, the study has a retrospective character which may result in limited access to some specific patient information. Last, the cost analysis concerns only single hospitalization due to AIS. The costs do not include treatment of the consequences of AIS, i.e. subsequent hospitalizations and rehabilitation.

Our study is the first analysis of the paediatric patients with AIS based on the NHF registry in Poland, which covers the entire population. This allowed for including a sizeable group of patients hospitalised in all regions of Poland due to AIS in contrast to the clinical studies most often coming from one medical centre and performed on small groups.

Conclusions

In the present study, the occurrence of AIS varies between regions of Poland. AIS is more common in boys than girls and more often found in adolescents (15–18 years) than younger children. Every tenth hospitalisation due to AIS is away from the patient's place of residence. Most often emergency admission, and the extract, most often referred for further outpatient treatment. The mean hospital stay due to paediatric AIS is about two weeks, with an average cost of hospitalisation of approximately USD 900.

It is noteworthy to use health registries for population analyses of various diseases and population distributions of demographic and hospitalisation-related characteristics.

Preprint




Figure 1 Age of paediatric patients hospitalised due to arterial ischaemic stroke in Poland in 2011–2021 (N=622).




Figure 2 Month and season of the year at hospital admission due to paediatric arterial ischaemic stroke in Poland in 2011–2021 (N=622)

Tables

Table I. Paediatric hospitalisations due to arterial ischaemic stroke acc. to ICD-10 in Poland in 2011–2020, n (%)

ICD-10 code	Diagnosis	Total (N=622)	2011 (N=59)	2012 (N=51)	2013 (N=49)	2014 (N=73)	2015 (N=55)	2016 (N=70)	2017 (N=56)	2018 (N=66)	2019 (N=65)	2020 (N=77)
I63.0	Cerebral infarction due to thrombosis of precerebral arteries	17 (2.73)	2 (3.39)	2 (3.92)	0 (0)	4 (5.48)	1 (1.82)	2 (2.86)	2 (3.57)	1 (1.52)	2 (3.08)	1 (1.3)
I63.1	Cerebral infarction due to embolism of precerebral arteries	1 (0.16)	0 (0)	0 (0)	0 (0)	1 (1.37)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
I63.2	Cerebral infarction due to unspecified occlusion or stenosis of precerebral arteries	13 (2.09)	2 (3.39)	1 (1.96)	0 (0)	1 (1.37)	2 (3.64)	2 (2.86)	2 (3.57)	1 (1.52)	0 (0)	2 (2.6)
I63.3	Cerebral infarction due to thrombosis of cerebral arteries	45 (7.23)	6 (10.17)	7 (13.73)	2 (4.08)	2 (2.74)	7 (12.73)	2 (2.86)	4 (7.14)	4 (6.06)	5 (7.69)	6 (7.79)
I63.4	Cerebral infarction due to embolism of cerebral arteries	27 (4.34)	4 (6.78)	1 (1.96)	2 (4.08)	2 (2.74)	3 (5.45)	1 (1.43)	3 (5.36)	1 (1.52)	4 (6.15)	6 (7.79)
I63.5	Cerebral infarction due to unspecified occlusion or stenosis of cerebral arteries	95 (15.27)	13 (22.03)	8 (15.69)	11 (22.45)	6 (8.22)	7 (12.73)	16 (22.86)	5 (8.93)	11 (16.67)	9 (13.85)	9 (11.69)
I63.6	Cerebral infarction due to cerebral venous thrombosis, nonpyogenic	21 (3.38)	2 (3.39)	3 (5.88)	2 (4.08)	2 (2.74)	2 (3.64)	2 (2.86)	1 (1.79)	3 (4.55)	3 (4.62)	1 (1.3)
I63.8	Other cerebral infarction	92 (14.79)	2 (3.39)	7 (13.73)	9 (18.37)	9 (12.33)	9 (16.36)	7 (10)	7 (12.5)	11 (16.67)	14 (21.54)	16 (20.78)
I63.9	Cerebral infarction, unspecified	114 (18.33)	12 (20.34)	10 (19.61)	6 (12.24)	15 (20.55)	12 (21.82)	17 (24.29)	12 (21.43)	8 (12.12)	12 (18.46)	10 (12.99)
I64	Stroke, not specified as haemorrhage or infarction	197 (31.67)	16 (27.12)	12 (23.53)	17 (34.69)	31 (42.47)	12 (21.82)	21 (30)	20 (35.71)	26 (39.39)	16 (24.62)	26 (33.77)

Table II. Region of residence of paediatric patients hospitalised due to arterial ischaemic stroke in Poland in 2011–2020, n (%)

Patient region of residence	Total (N=622)	2011 (N=59)	2012 (N=51)	2013 (N=49)	2014 (N=73)	2015 (N=55)	2016 (N=70)	2017 (N=56)	2018 (N=66)	2019 (N=65)	2020 (N=78)
Dolnośląskie	60 (9.65)	12 (20.34)	4 (7.84)	5 (10.20)	6 (8.22)	1 (1.82)	8 (11.43)	9 (16.07)	6 (9.09)	3 (4.62)	6 (7.69)
Kujawsko-pomorskie	32 (5.14)	2 (3.39)	2 (3.92)	2 (4.08)	3 (4.11)	4 (7.27)	3 (4.29)	2 (3.57)	5 (7.58)	0 (0.00)	9 (11.54)
Lubelskie	22 (3.54)	0 (0.00)	4 (7.84)	1 (2.04)	3 (4.11)	1 (1.82)	3 (4.29)	2 (3.57)	1 (1.52)	2 (3.08)	5 (6.41)
Lubuskie	21 (3.38)	3 (5.08)	4 (7.84)	2 (4.08)	3 (4.11)	2 (3.64)	0 (0.00)	2 (3.57)	3 (4.55)	1 (1.54)	1 (1.28)
Łódzkie	33 (5.31)	0 (0.00)	6 (11.76)	1 (2.04)	5 (6.85)	2 (3.64)	4 (5.71)	4 (7.14)	4 (6.06)	4 (6.15)	3 (3.85)
Małopolskie	57 (9.16)	1 (1.69)	4 (7.84)	3 (6.12)	12 (16.44)	7 (12.73)	5 (7.14)	6 (10.71)	5 (7.58)	3 (4.62)	11 (14.10)
Mazowieckie	72 (11.58)	4 (6.78)	10 (19.61)	7 (14.29)	10 (13.70)	2 (3.64)	14 (20.00)	4 (7.14)	8 (12.12)	6 (9.23)	7 (8.97)
Opolskie	17 (2.73)	1 (1.69)	0 (0.00)	3 (6.12)	1 (1.37)	4 (7.27)	2 (2.86)	0 (0.00)	3 (4.55)	2 (3.08)	1 (1.28)
Podkarpackie	41 (6.59)	8 (13.56)	2 (3.92)	1 (2.04)	5 (6.85)	0 (0.00)	8 (11.43)	5 (8.93)	1 (1.52)	5 (7.69)	6 (7.69)
Podlaskie	28 (4.50)	5 (8.47)	4 (7.84)	6 (12.24)	1 (1.37)	3 (5.45)	0 (0.00)	3 (5.36)	2 (3.03)	0 (0.00)	4 (5.13)
Pomorskie	45 (7.23)	2 (3.39)	1 (1.96)	1 (2.04)	2 (2.74)	5 (9.09)	6 (8.57)	4 (7.14)	1 (1.52)	16 (24.62)	7 (8.97)
Śląskie	62 (9.97)	4 (6.78)	1 (1.96)	4 (8.16)	7 (9.59)	7 (12.73)	9 (12.86)	9 (16.07)	13 (19.70)	3 (4.62)	5 (6.41)
Świętokrzyskie	26 (4.18)	7 (11.86)	5 (9.80)	2 (4.08)	3 (4.11)	4 (7.27)	0 (0.00)	2 (3.57)	2 (3.03)	0 (0.00)	1 (1.28)
Warmińsko-mazurskie	25 (4.02)	6 (10.17)	1 (1.96)	5 (10.20)	1 (1.37)	2 (3.64)	0 (0.00)	2 (3.57)	2 (3.03)	5 (7.69)	1 (1.28)
Wielkopolskie	59 (9.49)	4 (6.78)	1 (1.96)	3 (6.12)	9 (12.33)	9 (16.36)	8 (11.43)	2 (3.57)	7 (10.61)	13 (20.00)	3 (3.85)
Zachodniopomorskie	22 (3.54)	0 (0.00)	2 (3.92)	3 (6.12)	2 (2.74)	2 (3.64)	0 (0.00)	0 (0.00)	3 (4.55)	2 (3.08)	8 (10.26)
Hospitalisations outside the region of residence	57 (9.16)	10 (16.95)	3 (5.88)	2 (4.08)	5 (6.85)	9 (16.36)	5 (7.14)	5 (8.93)	6 (9.09)	7 (10.77)	5 (6.41)

Table III. Admission procedure of paediatric patients hospitalised due to AIS in Poland in 2011–2020, n (%)

Type of admission	Total (N=622)
Transfer from another hospital	33 (5.31)
Scheduled admission on the basis of a referral	104 (16.72)
Emergency admission – other cases	372 (59.81)
Emergency admission as a result of a transfer by an emergency medical team	111 (17.85)
Admission of a newborn as a result of childbirth in a given hospital	2 (0.32)

Table IV. Discharge procedure of paediatric patients hospitalised due to AIS in Poland in 2011–2020, n (%)

Type of discharge	Total (N=622)
Referral for further treatment in a medical facility other than a hospital, providing medical services such as inpatient and round-the-clock health services	5 (0.80)
Referral for further treatment in another hospital	144 (23.155)
Referral for further outpatient treatment	295 (47.43)
Discharge on request	8 (1.29)
Completion of the therapeutic or diagnostic process	158 (25.40)
Patient death	11 (1.77)
Patient arbitrarily left the healthcare facility providing inpatient and round-the-clock medical services before the end of the therapeutic or diagnostic process	1 (0.16)

Table V. Duration and costs of hospitalisation due to paediatric AIS acc. to ICD-10 in Poland in 2011–2020

ICD-10	Diagnosis	Duration of hospitalisation (days), M±SD	Costs of hospitalisation, M±SD		
			PLN	EUR*	USD*
	Total	12.9±12.1	3,288±3,324	729±737	891±901
I63.0	Cerebral infarction due to thrombosis of precerebral arteries	17.3±8.7	3,291±1,290	730±286	892±350
I63.1	Cerebral infarction due to embolism of precerebral arteries	9.0±0.0	1,2584±0.0	2790±0	3410±0
I63.2	Cerebral infarction due to unspecified occlusion or stenosis of precerebral arteries	15.2±12.7	4,331±2,783	960±617	1,174±754
I63.3	Cerebral infarction due to thrombosis of cerebral arteries	15.1±12.4	4,108±3,225	911±715	1,113±874
I63.4	Cerebral infarction due to embolism of cerebral arteries	15.7±17.1	5,935±11,41	1,316±253	1,608±3093
I63.5	Cerebral infarction due to unspecified occlusion or stenosis of cerebral arteries	16.9±11.6	3,753±2,330	832±517	1,017±631
I63.6	Cerebral infarction due to cerebral venous thrombosis, nonpyogenic	16.3±12.1	4,041±2,856	896±633	1,095±774
I63.8	Other cerebral infarction	15.4±12.7	4,182±2,636	927±584	1,133±714
I63.9	Cerebral infarction, unspecified	12.5±12.3	3,559±2,862	789±635	964±776
I64	Stroke, not specified as haemorrhage or infarction**	8.4±9.9	1,778±1,327	394±294	482±360

* EUR 1 = PLN 4.51 and USD 1 = PLN 3.69; exchange rate as of 31 December 2020

Table/Figure legends

Figure 1 Age of paediatric patients hospitalised due to arterial ischaemic stroke in Poland in 2011–2021 (N=622).

Figure 2 Month and season of the year at hospital admission due to paediatric arterial ischaemic stroke in Poland in 2011–2021 (N=622)

Table I. Paediatric hospitalisations due to arterial ischaemic stroke acc. to ICD-10 in Poland in 2011–2020, n (%)

Table II. Region of residence of paediatric patients hospitalised due to arterial ischaemic stroke in Poland in 2011–2020, n (%)

Table III. Admission procedure of paediatric patients hospitalised due to AIS in Poland in 2011–2020, n (%)

Table IV. Discharge procedure of paediatric patients hospitalised due to AIS in Poland in 2011–2020, n (%)

Table V. Duration and costs of hospitalisation due to paediatric AIS acc. to ICD-10 in Poland in 2011–2020

Preprint

References

1. Ferriero DM, Fullerton HJ, Bernard TJ, et al. Management of Stroke in Neonates and Children: A Scientific Statement From the American Heart Association/American Stroke Association. *Stroke*. 2019; 50(3): e51-e96.
2. Chiang KL, Cheng CY. Epidemiology, risk factors and characteristics of pediatric stroke: a nationwide population-based study. *QJM*. 2018; 111(7): 445-454.
3. Gerstl L, Weinberger R, Heinen F, et al. Arterial ischemic stroke in infants, children, and adolescents: results of a Germany-wide surveillance study 2015-2017. *J Neurol*. 2019; 266(12): 2929-2941.
4. Fink M, Slavova N, Grunt S, et al. Posterior Arterial Ischemic Stroke in Childhood. *Stroke*. 2019; 50(9): 2329-2335.
5. Lopez-Vicente M, Ortega-Gutierrez S, Amlie-Lefond C, et al. Diagnosis and management of pediatric arterial ischemic stroke. *J Stroke Cerebrovasc Dis*. 2010; 19: 175-183.
6. Nowak-Gottl U, Gunther G, Kurnik K, Strater R, Kirkham F. Arterial ischemic stroke in neonates, infants, and children: an overview of underlying conditions, imaging methods, and treatment modalities. *Semin Thromb Hemost* 2003; 29: 405-414.
7. Kopyta I, Sarecka-Hujar B, Skrzypek M. Post-stroke epilepsy in Polish paediatric patients. *Dev Med Child Neurol*. 2015; 57(9): 821-8.
8. Westmacott R, Askalan R, MacGregor D, Anderson P, Deveber G. Cognitive outcome following unilateral arterial ischaemic stroke in childhood: effects of age at stroke and lesion location. *Dev Med Child Neurol* 2010; 52: 386-93.
9. Stacey A, Toolis C, Ganesan V. Rates and Risk Factors for Arterial Ischemic Stroke Recurrence in Children. *Stroke*. 2018; STROKEAHA.117.020159, <https://doi.org/10.1161/STROKEAHA.117.020159>
10. Heron M. Deaths: leading causes for 2010. *Natl Vital Stat Rep*. 2013: p. 1-96.
11. Lin YW, Sung FC, Lin MH, et al. Medical Costs of Stroke Care Between Women With and Without Dysmenorrhea: A Population-Based Comparison. *Front Public Health*. 2021; 9: 699359.
12. Łabuz-Roszak B, Starostka-Tatar A, Lasek-Bal A, Gierlotka M, Gąsior M, Skrzypek M. Diagnostics, treatment and secondary prevention of ischemic stroke in the Silesian Province, Poland between 2009 and 2015. *Neurol Neurochir Pol*. 2018; 52(2): 235-242.
13. Lee M, Wu YL, Ovbiagele B. Trends in Incident and Recurrent Rates of First-Ever Ischemic Stroke in Taiwan between 2000 and 2011. *J Stroke*. 2016; 18(1): 60-65.
14. Bergdahl M, Ehling M, Elvers E, et al. *Handbook on Data Quality Assessment Methods and Tools*. Wiesbaden, 2007.
15. Zapalska A, Dziurda D, Podolec Z, Tronczyńska D. Towards an Effective Access into Polish National Health Data - Just a Step Away from a Wealth of Information. *Journal of Health Policy and Outcomes Research* 2012; 2: 86-93.

16. Kopyta I, Dobrucka-Głowacka A, Cebula A, Sarecka-Hujar B. Does the Occurrence of Particular Symptoms and Outcomes of Arterial Ischemic Stroke Depend on Sex in Pediatric Patients?-A Pilot Study. *Brain Sci.* 2020; 10(11): 881.
17. Rivkin MJ, Bernard TJ, Dowling MM, Amlie-Lefond C. Guidelines for Urgent Management of Stroke in Children. *Pediatr Neurol.* 2016; 56: 8-17.
18. Janjua N, Nasar A, Lynch JK, Qureshi AI. Thrombolysis for ischemic stroke in children: data from the nationwide inpatient sample. *Stroke.* 2007 Jun;38(6):1850-4.
19. Meyer S, Poryo M, Flotats-Bastardas M, Ebrahimi-Fakhari D, Yilmaz U. Schlaganfall bei Kindern und Jugendlichen [Stroke in children and adolescents]. *Radiologe.* 2017; 57(7): 569-576. [In German]
20. Fullerton HJ, Wu YW, Zhao S, Johnston SC. Risk of stroke in children: ethnic and gender disparities. *Neurology.* 2003; 61(2): 189-194.
21. Grunt S, Mazenauer L, Buerki SE, et al. Incidence and outcomes of symptomatic neonatal arterial ischemic stroke. *Pediatrics.* 2015; 135(5): e1220-1228.
22. Chen PC, Chien KL, Chang CW, et al. More hemorrhagic and severe events cause higher hospitalization care cost for childhood stroke in Taiwan. *J Pediatr.* 2008; 152(3): 388-393.
23. Cao Q, Yang F, Zhang J, Liang H, Liu X, Wang H. Features of Childhood Arterial Ischemic Stroke in China. *Fetal Pediatr Pathol.* 2019; 38(4): 317-325.
24. Helmuth IG, Mølbak K, Uldall PV, Poulsen A. Post-varicella Arterial Ischemic Stroke in Denmark 2010 to 2016. *Pediatr Neurol.* 2018; 80: 42-50.
25. Sarecka-Hujar B, Kopyta I. The Impact of Sex on Arterial Ischemic Stroke in Young Patients: From Stroke Occurrence to Poststroke Consequences. *Children (Basel).* 2021; 8(3): 238.
26. Fullerton HJ, Hills NK, Elkind MS, et al. Infection, vaccination, and childhood arterial ischemic stroke: Results of the VIPS study. *Neurology.* 2015; 85(17): 1459-1466.
27. Carey S, Wrogemann J, Booth FA, Rafay MF. Epidemiology, Clinical Presentation, and Prognosis of Posterior Circulation Ischemic Stroke in Children. *Pediatr Neurol.* 2017; 74: 41-50.
28. Ndiaye M, Lengue F, Sagna SD, et al. Childhood arterial ischemic stroke in Senegal (West Africa). *Arch Pediatr.* 2018; 25(6): 351-354.
29. Maino A, Algra A, Peyvandi F, Rosendaal FR, Siegerink B. Hypercoagulability and the risk of recurrence in young women with myocardial infarction or ischaemic stroke: a cohort study. *BMC Cardiovasc Disord.* 2019; 19(1): 55.
30. Bernard TJ, Goldenberg NA, Tripputi M, Manco-Johnson MJ, Niederstadt T, Nowak-Göttl U. Anticoagulation in childhood-onset arterial ischemic stroke with non-moyamoya arteriopathy: findings from the Colorado and German (COAG) collaboration. *Stroke.* 2009; 40(8): 2869-2871.
31. Sarecka-Hujar B, Kopyta I, Raczkiewicz D. Risk factors, types and outcomes of arterial ischemic stroke in Polish pediatric patients: a retrospective single-center study. *Arch Med Sci.* 2018; 17(1): 62-70.

32. Perkins E, Stephens J, Xiang H, Lo W. The cost of pediatric stroke acute care in the United States. *Stroke*. 2009; 40(8): 2820-2827.

Preprint