

Predictive value of imaging manifestations of supratentorial hemorrhage in hematoma enlargement and clinical prognosis

Keywords

Sensitivity and Specificity, Tomography, Computed Tomography Angiography, X-Ray Computed, Cerebral Angiography, Cerebral Hemorrhage, Hematoma

Abstract

Introduction

Intracerebral hemorrhage (ICH) is an important cause of death and disability. This study aimed to explore the imaging indicators identifying hematoma expansion in primary ICH and to provide a basis for its clinical treatment.

Material and methods

Hematoma expansion was evaluated by plain computed tomography (CT) scan and multi-detector-row CT angiography (MDCTA).

Results

This study included a total of 203 patients with intracerebral hemorrhage. The size of the hematoma at the time of admission was 32.5–92.3 mL (average, 45.5 mL). Original or reconstructed MDCTA images with contrast extravasation were available in 35 cases (17.2%). Patients were divided into two groups based on whether or not hematoma expansion had occurred: hematoma expansion group (n = 87) and no hematoma expansion group (n = 116). Patients with spot signs in the two groups were 37.93% and 3.45%, respectively, and those with spot sign ± blend sign ± black hole sign ± island sign (%) were 48.30% and 1.72%, respectively, with significant statistical differences. The sensitivity, specificity, and positive and negative predictive values of hematoma enlargement were 37.93%, 98.27%, and 94.29% and 67.86%, respectively. The sensitivity, specificity, and positive and negative predictive values of blend density sign ± irregular sign in predicting hematoma enlargement in ICH patients were 97.7%, 76.7%, and 75.9% and 97.8%, respectively.

Conclusions

Blend density sign ± irregular sign can be used as substitute signs. The more irregular and uneven density the hematoma, the more likely it is that the hematoma will be enlarged.

Predictive value of imaging manifestations of supratentorial hemorrhage in hematoma enlargement and clinical prognosis

Bai Qingke^{1#}, Wang Feng^{2#}, Zhao Zhenguo³, Zheng Ping^{4*}

1. Department of Neurology, Shanghai Pudong New Area People's Hospital

2. Department of Neurology, Shanghai Seventh People's Hospital

3. Department of Neurosurgery, Shanghai Pudong New Area People's Hospital

4. Department of Radiography, Shanghai Pudong New Area People's Hospital

These authors contribute equally to the work.

* Correspondence author: Dr. Zheng Ping, Email: jojo_ras@126.com

Address: 490, Chuanhuan Nan Rd, Shanghai, 201299, P.R.C.

Abstract

Objective: Intracerebral hemorrhage (ICH) is an important cause of death and disability. This study aimed to explore the imaging indicators identifying hematoma expansion in primary ICH and to provide a basis for its clinical treatment.

Methods: Hematoma expansion was evaluated by plain computed tomography (CT) scan and multi-detector-row CT angiography (MDCTA).

Results: This study included a total of 203 patients with intracerebral hemorrhage. The size of the hematoma at the time of admission was 32.5–92.3 mL (average, 45.5 mL). Original or reconstructed MDCTA images with contrast extravasation were available in 35 cases (17.2%). Patients were divided into two groups based on whether or not hematoma expansion had occurred: hematoma expansion group (n = 87) and no hematoma expansion group (n = 116). Patients with spot signs in the two groups were 37.93% and 3.45%, respectively, and those with spot sign ± blend sign ±

black hole sign \pm island sign (%) were 48.30% and 1.72%, respectively, with significant statistical differences. The sensitivity, specificity, and positive and negative predictive values of hematoma enlargement were 37.93%, 98.27%, and 94.29% and 67.86%, respectively. The sensitivity, specificity, and positive and negative predictive values of blend density sign \pm irregular sign in predicting hematoma enlargement in ICH patients were 97.7%, 76.7%, and 75.9% and 97.8%, respectively.

Conclusion: Blend density sign \pm irregular sign can be used as substitute signs. The more irregular and uneven density the hematoma, the more likely it is that the hematoma will be enlarged.

Key words: Cerebral Angiography; Cerebral Hemorrhage; Sensitivity and Specificity; Hematoma; Computed Tomography Angiography; Tomography, X-Ray Computed.

Introduction

Primary intracerebral hemorrhage (ICH) is an important cause of death and disability. The incidence of intracerebral hemorrhage in the Asian stroke population has been reported to range from 20–30%, which was significantly higher, even twice as high, as that reported from western countries [1]. This difference in incidence may be related to the distribution of risk factors such as genes, environment, and hypertension [2,3]. The neurological function often deteriorates rapidly within a few hours of onset of ICH [4]; therefore, it is vital to rapidly diagnose and treat this condition when it occurs.

Hematoma expansion has been considered to be one of the most important determinants of neurological deterioration and poor clinical outcomes in the early stage of primary ICH [5]. After ICH, secondary injury occurs as a result of expansion of the hematoma and peri-hematoma edema [6]. Hematoma expansion is an important risk factor involved in the deterioration of neurological function and poor prognosis in the early stage of ICH [7,8]. The aim of the present study was to retrospectively investigate the signs of contrast extravasation (spot sign) and other CTA image signs in patients with ICH obtained during the hyperacute period of primary ICH and their relationship with the location, size, morphology, and clinical prognosis of hematomas. We hope that the results can help us explore the imaging indicators of hematoma expansion in primary ICH and provide a basis for clinical monitoring of disease changes and strengthening antihypertensive therapy or timely selection of surgical intervention.

Methods

Study population and design

This study retrospectively analyzed data of patients with hemorrhagic stroke who had been treated at the Department of Neurology and Neurosurgery, Pudong New Area People's Hospital in Shanghai, from January 2017 to December 2019.

The inclusion criteria were as follows: (1) Patients who were 18–80 years, (2) those who arrived at the hospital within 6 h of onset; and (3) those with primary intracerebral hemorrhage confirmed by emergency CT examination.

The exclusion criteria were as follows: (1) Patients with deep coma (Glasgow coma score [GCS] of 3–5 points); (2) those who were allergic to contrast medium; (3) patients with previous thyroid disease or renal dysfunction; (4) those with congenital or acquired coagulation dysfunction; (5) those who did not complete CT plain scan and MDCTA.

This study was approved by the Ethics Committee of the Pudong New Area People's Hospital in Shanghai. As this was a retrospective study, informed consent was waived.

Imaging examination

Whole-brain CT scan was performed at the emergency admission, and further CTA was performed after hemorrhage was confirmed. A 64-row CT (Philips, the Netherlands) was used for examination. Patients received a 70 mL ioversol dose administered intravenously through the median elbow vein using a high-pressure syringe at a rate of 5 mL/s. CTA parameters were set to 120 kV and 300 mA. Imaging operations were completed by a technician with more than 5 years of working experience. All the images were uploaded to Neusoft picture archiving and communication system for analysis. The obtained imaging results were observed and

evaluated by two experienced neuroradiologists.

Clinical indicators and definition

Clinical data of the patients, including demographics, medical history at the time of admission, and clinical features, were collected for use in this study. Demographic characteristics included gender and age; medical history included history of drinking, hypertension, and diabetes; and the clinical characteristics included systolic blood pressure, diastolic blood pressure, GCS [9,10], and the National Institutes of Health Stroke Scale (NIHSS) score that were obtained at the time of admission.

According to the standard proposed by Delgado Almandoz et al. [11], hematoma expansion is defined as an increase in the size of the hematoma in ICH patients on subsequent CT scans by >12.5 mL or $>33\%$ compared with the first CT, when the first CT scan is within 6 hours of ICH onset and the second CT scan is performed within 24 hours after the first CT. The volume of the hematoma was evaluated using the method of three-dimensional drawing [12], which was especially suitable to measure irregular hematoma.

On imaging, when a black hole (low-density area) in the hematoma that was completely surrounded by an adjacent high-density hematoma, and satisfies the following criteria: (1) the shape was different, but not connected with adjacent brain tissues; (2) there was a clear boundary; (3) and CT values of the two different density areas in the hematoma differed by at least 28 HU, it was then regarded as a black hole sign [13,14].

On imaging, the presence of relatively low-density areas and adjacent high-density areas in the same hematoma was defined as a blend sign [15], provided

the following conditions were satisfied: (1) there was an obvious boundary between the low- and high-density areas that can be recognized by the naked eye, (2) the CT values of the two different density areas in the hematoma differed at least 18 HU, and (3) the low-density area was not completely enclosed by the high-density area. A blend sign was considered when all the above criteria were simultaneously met.

The black hole sign and blend sign were identified on plain CT scan images, and contrast medium extravasation signs and spot signs were identified on original or reconstructed MDCTA images, and spot sign scoring was completed.

Two CTA scans were performed for each ICH patient, including a CTA phase and a delayed phase (5 min after the CTA phase). The region of interest (ROI) was set with a diameter of 10 mm, and the CT value was calculated. When the CT value in the ROI at the delayed phase was more than 10% higher than that in the CTA phase, this phenomenon was defined as a leakage sign [16].

An enhanced lesion in the hematoma on the original CTA image was referred to as the spot sign [11]. According to the standard of spot sign proposed by Delgado et al. in 2009, a spot sign was considered to be present provided: 1) there was ≥ 1 enhanced lesion with concentrated contrast medium in the extravasation area of the intracranial hematoma; 2) compared with its density, the attenuation of the peripheral hematoma was greater than 120 HU, 3) the CTA spot sign should be separated from the blood vessels in the hematoma, and 4) the spot sign should be located within the intracranial hematoma.

CT re-examination was then performed at 24 hours, 2 days, and after 90 days. NIHSS scoring, 90-day quality of life, Barthel index (BI) [17](range, 0 [severe disability] to 100 [no disability]), and mRS scoring [18–22] (range, 0 [no symptoms] to 5 [severe disability]) were performed at the time of admission and at 24 hours, 7

days, and 14 days.

Mortality was observed during hospitalization, and patients who survived were followed up to observe their quality of life at 6 months after surgery.

The patients were categorized into one of five grades based on their activities of daily living (ADL) score [23]: grade 1, complete recovery of the family's ability to perform daily life activity; grade 2, the ability to perform daily life activities independently and partial recovery of social life; grade 3, some assistance was needed for daily life activities; grade 4, the patient retained consciousness, but assistance was needed for all daily life activities, and grade 5, survival in the vegetative state. Grades 1, 2, and 3 were good effects while grades 4 and 5 were bad effects.

Treatment

According to the Chinese guidelines for the diagnosis and treatment of ICH (2014), craniotomy or minimally invasive surgery was performed to surgically remove the supratentorial hematoma in patients with lobar hemorrhage of >30 mL and within 1 cm of the cortical surface. Patients who presented with supratentorial ICH with a hematoma volume ranging from 20–40 mL, a GCS score of ≥ 9 , and were within 72 hours of onset could be treated with minimally invasive surgery with or without liquidation by thrombolytic drugs for drainage of the hematoma after strict selection. In cases where patients had severe cerebral hemorrhage exceeding 40 mL, in whom the disturbance of consciousness worsened due to the space-occupying effect of the hematoma, craniotomy was performed to remove the hematoma.

Statistical analyses

Statistical analysis was performed using SPSS software version 25.0 (IBM,

Armonk, NY, USA). Continuous variables that conformed to normal distribution were represented by means \pm SD, and continuous variables that did not conform to the normal distribution were represented by the median (range or interquartile [IQR] range). Classification variables were represented by n (%). For continuous variables with normal distribution, *t* test and analysis of variance (ANOVA) were used, and non-normally distributed continuous variables were tested by the Mann-Whitney *U* test or the Kruskal-Wallis test. The chi-square test or Fisher's exact test were used for statistical analysis of classified variables.

Preprint

Results

We analyzed 203 patients with ICH in the present study, of which 87 had hematoma expansion while 116 had no hematoma expansion. The average size of the hematoma at the first admission was 32.5–92.3 mL (average 45.5 mL). A total of 35 cases (17.2%) had contrast medium extravasation signs on original or reconstructed MDCTA images, and spot sign scoring was performed. The relevant imaging manifestations were recorded, and hematomas with a volume >33% or 12.5 mL compared with the first scan image as defined previously was considered as a hematoma expansion. Table 1 showed the baseline data of 203 cases with ICH in the hematoma expansion and no expansion groups, and Table 2 and supplementary Figures 1–4 presented the clinical and imaging data.

A total of 69 cases received skull drilling for hematoma removal. The drainage volume ranged from 20–92 mL with an average of 46.5 mL, accounting for 40–70% of the hematoma. The postoperative NIHSS score was 3–24 points, with an average of 8.5 points. Of these, 4 cases underwent craniotomy after skull drilling and drainage due to hematoma expansion.

Table 3 showed that the sensitivity, specificity, and positive and negative predictive values of hematoma expansion were 37.93%, 98.27%, and 94.29% and 67.86%, respectively. The sensitivity, specificity, and positive and negative predictive values of blend density signs \pm irregular signs in predicting hematoma expansion in ICH patients were 97.7%, 76.7%, and 75.9% and 97.8%, respectively.

Five and seventeen patients in the surgical and non-surgical treatment groups died after surgery during hospitalization (mortality rate, 10.84%). The causes of death were brainstem failure caused by hematoma expansion or rebleeding in 12 cases, acute renal failure in 3 cases, and multiple organ failure in 7 cases. At the 6-month

postsurgical follow-up, 54 cases survived well after surgery in the hematoma expansion group (good rate, 78.3%), and 2 of the 18 in the no surgery hematoma expansion group showed good survival (good rate, 11.11%). The 181 patients who survived were followed up with postoperative CT re-examination. The results of the postoperative scan revealed a gradual decrease in hematoma size or disappearance of most of the hematoma, until finally encephalomalacia foci remained.

Preprint

Discussion

Hematoma expansion is a very common feature of ICH and occurs in 40% of patients with ICH, and it is considered an independent predictor of disease deterioration and poor prognosis [7]. Liu et al. [24] and Wu et al. [25] attempted to predict hematoma expansion in patients with ICH using support vector machine method with good results. Other studies have used imaging findings such as presence of a satellite sign or spot sign on CT scans to predict hematoma expansion [26]. Currently, however, no unified diagnostic standard exists, and diagnosis of ICH primarily depends on CTA and CT examination. We retrospectively studied CT and MDCTA findings of 203 patients and compared various imaging findings between two groups (one with hematoma expansion and one without hematoma expansion). In our study, the results indicated that blend density \pm irregular sign could be used as substitute signs on CT and CTA scans to predict hematoma expansion.

Recent studies have confirmed that some CTA and CT manifestations are known to be associated with hematoma expansion, such as spot sign, leakage sign, blend sign, black hole sign and island sign [26]. These imaging manifestations provide a more effective method to diagnose high-risk patients with ICH with hematoma expansion. Therefore, this study aimed to analyze the potential correlation between the manifestations of hematoma in CTA and CT scans and early hematoma expansion in patients with ICH, so that a timely and effective clinical intervention may be carried out for ICH patients with hematoma expansion, consequently reducing their mortality rate and improving the prognosis.

In the present study, we showed that spot sign, blend sign, black hole sign, and island sign reflect the heterogeneity of hematomas, and can predict the expansion of some types of hematomas. Together, the spot sign, blend sign, black hole sign, and

island sign show high specificity, but any of these signs exhibit low individual sensitivity. CTA examination in patients with ICH at an early stage was poorly operable, and was unsuitable for patients with obviously impaired renal function. In addition, the imaging standards of point sign, blend sign, black hole sign, and island sign were complicated and difficult for clinicians to grasp and operate quickly. We recommended blend density sign and/or the irregular sign as substitute signs. The sensitivity, specificity, and the positive and negative predictive values of blend density sign and/or irregular sign for predicting hematoma expansion in ICH patients were 97.7%, 76.7%, and 75.9% and 97.8%, respectively. Although the specificity and positive predictive value of these signs were slightly lower, their sensitivity and negative predictive values were relatively high. There was a significant statistical difference between predicting hematoma expansion and no expansion, and the evaluation indicator was simple. The maneuverability was strong. Preliminary studies have demonstrated that the more irregular and uneven the hematoma, the more it is likely that hematoma expansion will occur.

This study has some limitations. First, it is a single-center study that requires external validation of our results. Second, the study had a small sample size. Future multicenter studies with a larger sample size should be conducted on the basis of our study findings.

For the 69 patients that underwent minimally invasive surgery or craniotomy, 5 and 17 patients in the surgery and no surgery groups died during hospitalization (mortality rate, 10.84%). The follow-up conducted 6 months post-surgically showed that 54 and 2 cases survived well in the hematoma expansion surgery and no surgery group (good outcome rates, 78.3% and 11.11%, respectively). Preliminary studies

showed that minimally invasive treatment affected the precise treatment of hematomas in patients with ICH [27].

Conflict of Interest:

The authors declare that they do not have any conflict of interest.

Foundation items: Funded by The Featured Clinical Discipline Project of Shanghai Pudong (PWYst2018-01) and Key Discipline Group Construction Project of Shanghai Pudong (PWZxq2017-02) . Funded by General Project of Shanghai Natural Science Foundation (18ZR14307000) ; 2020 Health Science and Technology Project of Pudong New Area Health Commission (PW2020D-5) ; 2020 Science and Technology Development Fund of Pudong New Area Special Fund for People 's Livelihood Scientific Research (PKJ2020-Y-15)

Preprint

Table 1. Baseline data of 203 cases of cerebral hemorrhage in the two groups.

Clinical baseline data	Hematoma expansion group (n = 87)	No hematoma expansion group (n = 116)	<i>P</i> value
Age (year)	59.25 ± 9.46	56.17 ± 8.59	0.175
Male (%)	55.17 (48/87)	58.62 (68/116)	0.505
Hypertension (%)	83.90 (73/87)	81.90 (95/116)	0.536
Atrial fibrillation (%)	9.20 (8/87)	8.33 (10/116)	0.578
Diabetes (%)	12.64 (11/87)	9.48 (11/116)	0.732
Stoke side (left/right)	45/42	61/55	0.425
Time of onset (h)	2.30 ± 1.46	2.15 ± 1.63	0.233

There was no significant difference in the baseline data of patients with cerebral hemorrhage between the two groups ($P > 0.05$).

Table 2. Clinical and imaging data of 203 cases of cerebral hemorrhage in the two groups

Items	Hematoma expansion	No hematoma expansion	<i>P</i> value
	group (n = 87)	group (n = 116)	
Baseline NIHSS	11.53 ± 4.82	10.98 ± 4.11	0.296
24-h NIHSS	17.74 ± 4.36	12.08 ± 4.73	0.008
7-day NIHSS	15.05 ± 4.23	5.68 ± 4.96	0.000
CTA spot sign (%)	37.93 (33/87)	3.45 (2/116)	0.000
Blend sign (%)	11.50 (10/87)	2.59 (3/116)	0.000
Black hole sign (%)	13.79 (12/87)	3.45 (4/116)	0.000
Island sign	10.34 (9/87)	2.59 (2/116)	0.000
Spot sign ± blend sign ± black hole sign ± island sign (%)	48.30 (42/87)	1.72 (6/116)	0.000
Irregular hematoma sign (%)	87.36 (76/87)	21.55 (25/116)	0.000
Blend density sign (%)	93.10 (81/87)	19.83 (23/116)	0.000
Irregular hematoma sign ± blend density sign (%)	97.7 (85/87)	23.28 (27/116)	0.000
Mortality rate (%)	16.09 (14/87)	6.90 (8/116)	0.005
90 d mRS (0 or 1) (%)	49.32 (36/73)	66.67 (72/108)	0.009
90d BI (≥90) (%)	52.05 (38/73)	68.52 (74/108)	0.006
Good survival at 6 m (grade 1, 2, and 3 of ADL scoring)	54.80 (40/73)	71.30 (77/108)	0.009

Abbreviations: NIHSS = National Institutes of Health Stroke Scale, CTAA = CT angiography, mRS = modified Rankin scale, BI = Barthel index, ADL = activities of daily living

There was no significant difference in the baseline NIHSS score between the two groups ($P > 0.05$), but the 24-h and 7-day NIHSS scores were both significantly better in the no hematoma expansion group than in the hematoma expansion group ($P < 0.01$).

Table 3 ROC prediction of each variable

Variable	Sensitivity	Specificity	PPV	NPV
Hematoma expansion	37.93%	98.27%	94.29%	67.86%
Blend density sign \pm Irregular sign	97.7%	76.7%	75.9%	97.8%

Supplementary figures and legends

Supplementary Figure 1

The patient was a woman with a history of 8 years of hypertension. She was admitted to the hospital due to sudden left limb paralysis for 2 h. The NIHSS score was 16 points.

Figs. 1–4. The first CT scan showed high-density hemorrhagic shadow (size $3.0 \times 7.30 \times 3.0 \text{ cm}^3$) in the right basal ganglia. The periphery of the hematoma was irregular and the island sign can be seen around the hematoma. The right lateral ventricle was slightly displaced by compression.

Figs. 5–8. The original CTA images reveal a spot sign in the hematoma.

Figs. 9–12. CT re-examination after 12 h showed that the hematoma had expanded and ruptured into the right ventricle. The hematoma had an uneven density, which can be visualized as a blend sign and black hole sign.

Figs. 13–16 and 17–20. Results of the CT re-examination after 12, 24, and 72 h revealed further expansion of the hematoma and rupture into the ventricle, and a new island sign appeared. The ventricle and midline structure were obviously compressed to the left, and the ventricle was enlarged and exhibited hydrocephalus. Seven days later, the patient died due to pulmonary infection and respiratory and circulatory failure.

Supplementary Figure 2

The patient was a man with a history of 12 years of hypertension. He was admitted due to weakness of the right limb for 1 hour and had an NIHSS score of 10 points.

Figs. 1–4. The first CT scan shows a high-density hemorrhagic shadow ($2.1 \times$

5.0 × 3.0 cm³) in the left basal ganglia. The periphery of the hematoma was regular. There was no island sign around the hematoma, and no obvious compression or displacement of the lateral ventricle and midline were visible.

Figs. 5–8. The original CTA images showed no spot sign in the hematoma.

Figs. 9–12. CT re-examination after 72 h showed changes in hematoma absorption. The density was uniform, there was no blend sign and black hole sign. The 72-h NIHSS score was 6 points, and the 90-d mRS score and BI score were 1 and 90 points, respectively.

Supplementary figure 3

The patient was a woman who was admitted due to sudden dizziness and left limb paralysis for 3 hours. Her mental state was blurred, and the GCS and NIHSS were 13 and 18 points respectively.

Figs. 1–4. The first CT scan showed high-density hemorrhagic shadow in the right basal ganglia, the size was 2.5 × 4.0 × 4.0 cm³, the periphery of the hematoma was irregular, an atypical island sign was seen around the hematoma, and the right lateral ventricle was slightly compressed.

Fig. 5–8. Original CTA images show the presence of a spot sign in the hematoma. CTA images were intra-hematoma via vascular images.

Fig. 9–16. Re-examination of CT after 12 h showed that the hematoma had expanded and ruptured into the right ventricle; it had an uneven internal density, and a new blend sign and island sign appeared.

Figs. 17–20. Re-examination of CT 24 hours after surgery showed that the effect was good. The 90d mRS score and BI score were 2 and 85 points, respectively.

Supplementary figure 4

The patient was a woman who was admitted to the hospital due to sudden headache, aphasia, and mild weakness of the right limb for 4 hours. The GCS score was 11 points (E4V1M6), and the NIHSS was 14 points.

Figs. 1–4. Original CTA images show thick and twisted blood vessels around the intracerebral hematoma in the left parietal lobe. The reconstructed CTA images show an arteriovenous malformation in the left parietal lobe. CT re-examination after 7 days showed that the hematoma was absorbed. The 90d mRS and BI scores were 0 and 95 points, respectively.

References


- 1 Khan NA, McAlister FA, Pilote L, *et al.* Temporal trends in stroke incidence in South Asian, Chinese and white patients: A population based analysis. *Plos One* 2017;12:e0175556. doi:10.1371/journal.pone.0175556
- 2 Chaturvedi S, Greenberg SM. Intracerebral hematoma expansion: making predictions (about the future). *Neurology* 2011;76:1204–5. doi:10.1212/wnl.0b013e31821435d0
- 3 Xie N, Li X, Zhong Q, *et al.* Association of systolic blood pressure and pulse pressure with microalbuminuria in treatment-naïve hypertensive patients. *Archives Medical Sci Ams* 2019;15:832–6. doi:10.5114/aoms.2018.77727
- 4 Li N, Wang Y, Wang W, *et al.* Contrast extravasation on computed tomography angiography predicts clinical outcome in primary intracerebral hemorrhage: a prospective study of 139 cases. *Stroke* 2011;42:3441–6. doi:10.1161/strokeaha.111.623405
- 5 Cao Z, Li S, Yang H, *et al.* Associations of behaviors, biological phenotypes and cardiovascular health with risks of stroke and stroke subtypes: A prospective cohort study. *Eclinicalmedicine* 2021;33:100791. doi:10.1016/j.eclinm.2021.100791

- 6 Ljubisavljevic S, Ignjatovic A, Ljubisavljevic M. Headache secondary to nontraumatic brain hemorrhage: a single-center, retrospective clinical study. *Arch Med Sci* Published Online First: 2019. doi:10.5114/aoms.2019.90356
- 7 Morotti A, Li Q, Mazzoleni V, *et al.* Non-contrast CT markers of intracerebral hemorrhage expansion: The influence of onset-to-CT time. *Int J Stroke* 2022;;174749302211427. doi:10.1177/17474930221142742
- 8 Zimmer S, Meier J, Minnerup J, *et al.* Prognostic Value of Non-Contrast CT Markers and Spot Sign for Outcome Prediction in Patients with Intracerebral Hemorrhage under Oral Anticoagulation. *J Clin Medicine* 2020;9:1077. doi:10.3390/jcm9041077
- 9 Berker H de, Berker A de, Aung H, *et al.* 265 Use of support vector machine to predict discharge destination on an acute stroke ward. *J Neurology Neurosurg Psychiatry* 2022;93:A90.1-A90. doi:10.1136/jnnp-2022-abn.292
- 10 Odderson IR. The National Institutes of Health Stroke Scale and Its Importance in Acute Stroke Management. *Phys Med Rehabil Clin* 1999;10:787–800. doi:10.1016/s1047-9651(18)30162-1
- 11 Almandoz JED, Yoo AJ, Stone MJ, *et al.* Systematic characterization of the computed tomography angiography spot sign in primary intracerebral hemorrhage identifies patients at highest risk for hematoma expansion: the spot sign score. *Stroke* 2009;40:2994–3000. doi:10.1161/strokeaha.109.554667
- 12 Divani AA, Majidi S, Luo X, *et al.* The ABCs of Accurate Volumetric Measurement of Cerebral Hematoma. *Stroke* 2011;42:1569–74. doi:10.1161/strokeaha.110.607861
- 13 Arba F, Rinaldi C, Boulouis G, *et al.* Noncontrast Computed Tomography Markers of Cerebral Hemorrhage Expansion: Diagnostic Accuracy Meta-Analysis. *Int J Stroke Official J Int Stroke Soc* 2021;17:17474930211061640. doi:10.1177/17474930211061639
- 14 Boulouis G, Morotti A, Charidimou A, *et al.* Noncontrast Computed Tomography Markers of Intracerebral Hemorrhage Expansion. *Stroke* 2017;48:1120–5. doi:10.1161/strokeaha.116.015062
- 15 Sporns PB, Schwake M, Schmidt R, *et al.* Computed Tomographic Blend Sign Is Associated With Computed Tomographic Angiography Spot Sign and Predicts Secondary Neurological Deterioration After Intracerebral Hemorrhage. *Stroke* 2018;48:131–5. doi:10.1161/strokeaha.116.014068


- 16 Orito K, Hirohata M, Nakamura Y, *et al.* Leakage Sign for Primary Intracerebral Hemorrhage: A Novel Predictor of Hematoma Growth. *Stroke* 2016;47:958–63. doi:10.1161/strokeaha.115.011578
- 17 Group TE. EuroQol - a new facility for the measurement of health-related quality of life. *Health Policy* 1990;16:199–208. doi:10.1016/0168-8510(90)90421-9
- 18 Qingke B, Ping Z, Jianying Z, *et al.* Clinical comparison of intravenous thrombolysis and bridging artery thrombectomy in hyperacute ischemic stroke with unknown time of onset. *Archives Medical Sci Ams* 2021;17:1804–10. doi:10.5114/aoms/143513
- 19 Ping Z, Huiyu S, Min L, *et al.* Explainable machine learning for long-term outcome prediction in two-center stroke patients after intravenous thrombolysis. *Front Neurosci-switz* 2023;17:1146197. doi:10.3389/fnins.2023.1146197
- 20 Hutanu A, Iancu M, Dobreanu M, *et al.* Extended lipid profile in Romanian ischemic stroke patients in relation to stroke severity and outcome: a path analysis model. *Arch Med Sci* 2021;17:864–73. doi:10.5114/aoms.2019.89302
- 21 Zeng J, Zheng P, Xu J, *et al.* Prediction of motor function by diffusion tensor tractography in patients with basal ganglion haemorrhage. *Archives of medical science : AMS* 2011;7:310–4. doi:10.5114/aoms.2011.22083
- 22 Ping Z, Min L, Qiuyun L, *et al.* Prognostic nomogram for the outcomes in acute stroke patients with intravenous thrombolysis. *Front Neurosci-switz* 2022;16:1017883. doi:10.3389/fnins.2022.1017883
- 23 Cabañero-Martínez MJ, Cabrero-García J, Richart-Martínez M, *et al.* The Spanish versions of the Barthel index (BI) and the Katz index (KI) of activities of daily living (ADL): A structured review. *Arch Gerontol Geriat* 2009;49:e77–84. doi:10.1016/j.archger.2008.09.006
- 24 Liu J, Xu H, Chen Q, *et al.* Prediction of hematoma expansion in spontaneous intracerebral hemorrhage using support vector machine. *Ebiomedicine* 2019;43:454–9. doi:10.1016/j.ebiom.2019.04.040
- 25 Wu T-C, Liu Y-L, Chen J-H, *et al.* The Added Value of Intraventricular Hemorrhage on the Radiomics Analysis for the Prediction of Hematoma Expansion of Spontaneous Intracerebral Hemorrhage. *Diagnostics* 2022;12:2755. doi:10.3390/diagnostics12112755
- 26 Huang Y-W, Huang H-L, Li Z-P, *et al.* Research advances in imaging markers for predicting hematoma expansion in intracerebral hemorrhage: a narrative review. *Front Neurol* 2023;14:1176390. doi:10.3389/fneur.2023.1176390

27 He G-N, Guo H-Z, Han X, *et al.* Comparison of CT black hole sign and other CT features in predicting hematoma expansion in patients with ICH. *J Neurol* 2018;265:1883–90. doi:10.1007/s00415-018-8932-6

Preprint




1




2




3




4




5



6




7




8




9




10




11




12




Supplementary Figure 1



Supplementary Figure 2



Supplementary Figure 3



Supplementary Figure 4