

# The correlation between coagulation biomarkers, bone turnover markers, and bone mineral density in Chinese adults aged $\geq 50$ years

Zhengming Wang<sup>1,2,3</sup>, Rui Wang<sup>3,4,5</sup>, Haiya Ge<sup>1,2</sup>, Yong Gu<sup>3,6,7</sup>, Sicheng Xian<sup>7</sup>, Laijun Yan<sup>1,2</sup>, Guoqing Du<sup>1</sup>, Zhibi Shen<sup>1</sup>, Shuaijie Lv<sup>3</sup>, Hongsheng Zhan<sup>1</sup>, Zhengyan Li<sup>1,2</sup>

<sup>1</sup>Shi's Center of Orthopedics and Traumatology, Shuguang Hospital Affiliated to Shanghai University of Traditional Chinese Medicine, Shanghai, China

<sup>2</sup>Institute of Traumatology and Orthopedics, Shanghai Academy of Traditional Chinese Medicine, Shanghai, China

<sup>3</sup>Center of Orthopedics and Traumatology, The First Affiliated Hospital of Zhejiang Chinese Medical University (Zhejiang Provincial Hospital of Chinese Medicine), Zhejiang, China

<sup>4</sup>Guanghua Clinical Medical College, Shanghai University of Traditional Chinese Medicine, Shanghai, China

<sup>5</sup>Department of Orthopedic Surgery, Shanghai Guanghua Hospital of Integrated Traditional Chinese and Western Medicine, Shanghai, China

<sup>6</sup>Translational Medical Innovation Center, Zhangjiagang TCM Hospital Affiliated to Nanjing University of Chinese Medicine, Jiangsu, China

<sup>7</sup>Department of Orthopedics, Zhangjiagang TCM Hospital Affiliated to Nanjing University of Chinese Medicine, Jiangsu, China

**Submitted:** 24 October 2023; **Accepted:** 20 February 2024

**Online publication:** 12 March 2024

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms/184346>

Copyright © 2024 Termedia & Banach

## Abstract

**Introduction:** Osteoporosis patients undergoing surgery often have significant bleeding, which is closely related to coagulation function. This study aimed to assess the correlation between bone turnover markers (BTMs), bone mineral density (BMD), and coagulation biomarkers.

**Material and methods:** Cases meeting the inclusion criteria were included from our center from January 2020 to December 2021. The relationships between coagulation biomarkers, BTMs, and BMD were investigated by Spearman correlation analysis and canonical correlation analysis (CCA). Consequently, age, bone mass stage, 25-hydroxyvitamin D (25-(OH)D), and BMD were included as independent variables in the logistic regression model. Furthermore, the international normalized ratio (INR), fibrinogen, platelets, and D-dimer levels were set as dependent variables in the regression analysis. Finally, multivariable regressions were conducted to adjust for differences in demographics and comorbidities among groups.

**Results:** A total of 306 patients were included in the study. Significant differences were observed in prothrombin time, fibrinogen, INR, D-dimer, and platelets ( $p < 0.05$ ). D-dimer exhibited the most significant positive association with age and the most pronounced negative correlation with the BMD of Ward's triangle ( $r = 0.322$ ,  $p < 0.001$ ;  $r = -0.298$ ,  $p < 0.001$ ; respectively). 25-(OH)D and calcium were related to coagulation biomarkers. Multivariate analysis demonstrated that age was positively associated with D-dimer levels (odds ratio [OR] = 1.046,  $p = 0.007$ ), while the BMD of Ward's triangle was negatively related to D-dimer levels (OR = 0.004,  $p = 0.0017$ ).

**Conclusions:** For the Chinese population aged 50 and above, attention should be paid to alterations in coagulation biomarkers.

**Key words:** coagulation, bone mineral density, bone turnover marker, correlation.

## Corresponding authors:

Zhengyan Li  
Shi's Center of  
Orthopedics and  
Traumatology  
Shuguang Hospital  
Affiliated to Shanghai  
University of  
Traditional Chinese  
Medicine  
Shanghai, China  
E-mail: 62023230@shutcm.edu.cn

Hongsheng Zhan  
Shi's Center of Orthopedics  
and Traumatology  
Shuguang Hospital  
Affiliated to Shanghai  
University of  
Traditional Chinese  
Medicine  
Shanghai, China  
E-mail:  
zhanhongsheng@shutcm.edu.cn

## Introduction

Osteoporosis is a systemic skeletal disorder characterized by diminished bone mass and the deterioration of bone microarchitecture, resulting in bone fragility and fracture susceptibility [1–3]. The condition is highly prevalent in middle-aged and older populations, showing a higher incidence in women [4]. Fractures, particularly of the hip and lumbar spine, are the most common complication of osteoporosis. Globally, 8.9 million cases of osteoporosis-related fractures are reported annually [5], the majority of which necessitate surgical interventions such as hemiarthroplasty, total arthroplasty, open reduction and internal fixation, and vertebroplasty [6, 7]. These fractures impose significant economic and health-care burdens on individuals and governments.

Minimizing perioperative hemorrhage is of paramount importance to ensure procedural efficacy and improve therapeutic outcomes [8]. Despite the widespread use of tranexamic acid, controlling bleeding remains a challenge in surgery [9]. Therefore, the factors affecting perioperative bleeding require further research. A previous study [10] reported that higher bone mineral density (BMD) was associated with less bleeding during total hip arthroplasty. The amount of bleeding during an operation is largely determined by the coagulation function. Hence, coagulation function is hypothesized to be associated with bone turnover or mass.

Prothrombin time (PT) and the international normalized ratio (INR) are the most common coagulation tests, providing an assessment of the extrinsic coagulation pathway [11]. Conversely, activated partial thromboplastin time (APTT) measures the activity of the intrinsic coagulation pathways [12]. Fibrinogen is a soluble glycoprotein that plays a crucial role in platelet aggregation and interactions with various plasma proteins, participating in the blood clotting response to tissue injury [13]. Moreover, D-dimer is a highly sensitive marker of intravascular blood clots and hypercoagulability and is associated with both arterial and venous thrombotic events [14]. Collectively, fibrinogen, APTT, TT, PT, and INR provide a comprehensive assessment of patients' coagulation status [15]. Additionally, platelet (PLT) activation and aggregation play an essential role in hemostasis and thrombosis [16]. These parameters were measured as coagulation biomarkers in this study. Consequently, this study aimed to evaluate the relationship between bone turnover markers (BTMs), BMD, and coagulation biomarkers.

## Material and methods

### Patients

The study was approved by the Ethics Committee and written informed consent was waived

(2023-KLS-223-01). A total of 306 eligible patients from our hospital were enrolled between January 2020 and December 2021. All diagnoses were established according to the World Health Organization's criteria for osteoporosis [17]. The exclusion criteria were as follows: estimated creatine clearance < 60 ml/min, diabetes, dyslipidemia, tumors, hepatitis virus infection, fractures within the past year, age < 50 years, and body mass index (BMI) > 36.0 kg/m<sup>2</sup>. Additionally, none of the patients had a history of prior bone-active medication or treatments that could impact bone mineral metabolism (such as steroids, diuretics, thyroid hormones, anticonvulsant drugs, and lithium) or coagulation. Furthermore, no patient reported a history of secondary osteoporosis.

### Laboratory examination

The markers in this study were measured following established protocols [18, 19]. Patients were fasted for a minimum period of 8 h, and blood samples were collected from the cubital vein. The obtained venous blood serum was analyzed using a fully automated biochemical instrument. All BTMs were analyzed at a clinical laboratory using a COBAS E601 automated analyzer (Roche Diagnostics, Germany) via the electrochemoluminescence immunoassay method, following the manufacturer's instructions. In addition, the coagulation biomarkers were all measured at our central coagulation laboratory, using the Sysmex CN-6000 System (Siemens Healthineers, GmbH, Marburg, Germany). All tests were carried out according to the manufacturer's instructions. The normal range of laboratory test indexes is displayed in Supplementary Table S1.

### Evaluation of bone mineral density

Dual-energy X-ray absorptiometry scans were performed to assess BMD in all subjects using the Prodigy Advance system (General Electric Company, Fairfield, CT, USA). These scans evaluated the femoral neck, Ward's triangle, total hip, and lumbar spine 1 to 4, and were conducted within two weeks of the laboratory tests. The acquired scan data were transferred to a computer for detailed analysis, including the determination of bone mineral content. BMD (g/cm<sup>2</sup>) was calculated based on the bone area and bone mineral content scores, and T-scores were calculated for all measured anatomical regions. The scans were performed by an independent operator who was blinded to the analytical outcomes.

### Statistical analysis

All statistical analyses were conducted using SPSS version 25.0 (IBM, Armonk, NY, USA). Data

conforming to a normal distribution were compared by an unpaired *t*-test; otherwise, the non-parametric Mann-Whitney *U* test was utilized. Normally distributed data were presented as mean  $\pm$  standard deviation, whereas non-normally distributed data were presented as median and interquartile range. Differences in categorical data were assessed using the  $\chi^2$  test, and the results are expressed as *n* (%).

Spearman correlation analysis was employed to assess the correlation between patient information and coagulation evaluation indexes. Additionally, the relationships between patient information (independent variables, X) and coagulation evaluation indexes (dependent variables, Y) were examined using canonical correlation analysis (CCA) [20]. The canonical correlation coefficient represents the maximum possible correlation between linear combinations of variables in the independent variable set X and dependent variable set Y [21]. Consequently, this study focused solely on the first set of canonical correlations. First, significant influencing factors were identified by logistic regression analysis. Sub-

sequently, multivariate logistic regression was conducted to evaluate the factors with *p*-values  $< 0.05$  in the univariate logistic regression, thereby identifying the variables significantly affecting the outcome. The significance level was set at 0.05, and confidence intervals were reported at the 95% level.

## Results

### Participant characteristics

The demographics and clinical characteristics of all subjects are listed in Table I. The median age of all participants was 64.0 years, ranging from 50 to 90. The mean BMI was 23.5 (4.0) and ranged from 15.8 to 35.6. The study included 56 male participants and 97 individuals diagnosed with hypertension. Apart from 25-hydroxyvitamin D (25-(OH)D), all laboratory tests, including coagulation biomarkers and BTMs, were within the normal range. The median level of 25-(OH)D was 43.9, with a minimum of 7.0 and a maximum of 144.0. The mean BMD of the entire group of participants exhibited the osteopenia level.

**Table I.** General demographic and clinical characteristics of subjects

Variables	Overall (N = 306)	Normal (N = 46)	Osteopenia (N = 129)	Osteoporosis (N = 131)	P-value
Age [years]	64.0 (13.0)	57.5 (11.0)	61.0 (13.0)	68.0 (12.0)	<b>&lt; 0.001</b>
BMI [kg/m <sup>2</sup> ]	23.5 (4.0)	25.2 $\pm$ 2.7	23.6 (3.8)	22.7 (4.4)	<b>&lt; 0.001</b>
Male (n/%)	56/18.3	17/37.0	28/21.7	11/8.4	<b>&lt; 0.001</b>
Hypertension (n/%)	97/31.7	15/32.6	34/26.4	48/36.6	0.205
Coagulation biomarker:					
PT [s]	11.0 (1.0)	10.9 (1.0)	10.9 (1.0)	11.2 (1.0)	<b>0.010</b>
APTT [s]	28.3 (3.0)	28.2 (3.0)	28.3 $\pm$ 2.0	28.5 (3.0)	0.772
Fibrinogen [g/l]	3.1 (1.0)	3.1 $\pm$ 0.8	3.1 (1.0)	3.3 (1.0)	<b>0.009</b>
INR	0.9 (0)	0.9 (0)	0.9 (0)	0.9 $\pm$ 0.1	<b>0.005</b>
D-dimer [mg/l]	0.4 (1.0)	0.3 (0)	0.3 (0)	0.5 (1.0)	<b>&lt; 0.001</b>
Platelets [10 <sup>9</sup> /l]	195.0 (74.0)	198.0 $\pm$ 59.5	212.0 (70.0)	190.0 (75.0)	<b>0.031</b>
MPV [fl]	9.1 (2.0)	9.4 $\pm$ 1.5	8.9 (2.0)	9.1 (2.0)	0.672
BTMs:					
$\beta$ -CTX [ng/l]	626.0 (410.0)	524.9 (424.1)	607.7 (435.0)	685.6 $\pm$ 277.9	<b>0.013</b>
25-(OH)D [nmol/l]	43.9 (36.8)	52.8 $\pm$ 25.6	42.3 (37.5)	43.8 (36.7)	0.336
PINP [ $\mu$ g/l]	53.8 (32.1)	48.1 (26.4)	51.1 (33.0)	59.7 (31.1)	<b>0.014</b>
Osteocalcin [ $\mu$ g/l]	17.4 (9.1)	16.0 (7.7)	17.5 (8.4)	18.4 (10.0)	0.088
Calcium [mmol/l]	2.31 $\pm$ 0.12	2.32 $\pm$ 0.12	2.33 $\pm$ 0.12	2.29 $\pm$ 0.13	<b>0.045</b>
BMD:					
Lumbar spine [g/cm <sup>2</sup> ]	0.9 (0.3)	1.2 (0.2)	1.0 (0.2)	0.8 (0.1)	<b>&lt; 0.001</b>
Femoral neck [g/cm <sup>2</sup> ]	0.8 $\pm$ 0.1	1.0 (0.1)	0.8 $\pm$ 0.1	0.7 (0.1)	<b>&lt; 0.001</b>
Ward's triangle [g/cm <sup>2</sup> ]	0.6 $\pm$ 0.1	0.8 $\pm$ 0.1	0.6 (0.1)	0.6 $\pm$ 0.1	<b>&lt; 0.001</b>
Total hip [g/cm <sup>2</sup> ]	0.8 $\pm$ 0.1	1.0 (0.1)	0.9 $\pm$ 0.1	0.7 $\pm$ 0.1	<b>&lt; 0.001</b>

BMI – body mass index, PT – prothrombin time, APTT – activated partial thromboplastin time, INR – international normalized ratio, MPV – mean platelet volume, BTMs – bone turnover markers,  $\beta$ -CTX –  $\beta$ -cross-linked C-terminal telopeptide of type I, 25-(OH)D – 25-hydroxyvitamin D, PINP – procollagen type I N propeptide. BMD – bone mineral density, **boldface** indicates *p*-value  $< 0.05$ .

Table II. Correlation of coagulation biomarkers with characteristics, bone turnover markers, and bone mineral density

Variables	PT		APTT		Fibrinogen		INR		D-dimer		PLT		MPV	
	r	P	r	P	r	P	r	P	r	P	r	P	r	P
Characteristics:														
Age	0.188	<b>0.001</b>	-0.052	0.360	0.111	0.510	0.192	<b>0.001</b>	0.332	<b>&lt; 0.001</b>	-0.103	0.073	-0.035	0.548
BMI	-0.074	0.195	-0.076	0.182	-0.075	0.193	-0.083	0.145	-0.001	0.983	-0.040	0.489	0.011	0.852
Sex	0.067	0.243	-0.095	0.097	0.067	0.240	0.074	0.195	0.032	0.576	0.055	0.335	-0.104	0.069
Hypertension	0.040	0.485	-0.033	0.569	0.057	0.323	0.037	0.515	0.087	0.130	-0.034	0.558	0.064	0.262
Stage of bone mass	0.164	<b>0.004</b>	0.040	0.491	0.169	<b>0.003</b>	0.180	<b>0.002</b>	0.225	<b>&lt; 0.001</b>	-0.090	0.115	0.010	0.864
BTMs:														
β-CTX	0.005	0.934	0.010	0.855	-0.026	0.654	0.013	0.821	0.017	0.770	0.043	0.451	0.012	0.829
25-(OH)D	0.121	<b>0.034</b>	-0.179	<b>0.002</b>	0.084	0.144	0.139	<b>0.015</b>	0.021	0.717	0.196	<b>0.001</b>	-0.216	<b>&lt; 0.001</b>
PINP	0.058	0.311	0.020	0.724	0.021	0.719	0.064	0.263	0.046	0.424	0.016	0.786	0.036	0.532
Osteocalcin	-0.076	0.186	0.058	0.309	-0.055	0.334	-0.070	0.222	-0.041	0.475	0.034	0.559	-0.016	0.775
Calcium	-0.158	<b>0.006</b>	-0.128	<b>0.025</b>	0.074	0.198	-0.149	<b>0.009</b>	-0.132	<b>0.021</b>	0.169	<b>0.003</b>	0.036	0.529
BMD:														
Lumbar spine	-0.114	<b>0.047</b>	-0.037	0.515	-0.144	<b>0.012</b>	-0.131	<b>0.022</b>	-0.168	<b>0.003</b>	0.047	0.411	-0.038	0.512
Femoral neck	-0.101	0.079	0.059	0.300	-0.162	<b>0.005</b>	-0.111	0.052	-0.249	<b>&lt; 0.001</b>	0.067	0.242	0.007	0.906
Ward's triangle	-0.127	<b>0.026</b>	0.052	0.367	-0.172	<b>0.003</b>	-0.135	<b>0.018</b>	-0.298	<b>&lt; 0.001</b>	0.077	0.177	0.029	0.613
Total hip	-0.151	<b>0.008</b>	-0.031	0.592	-0.191	<b>0.001</b>	-0.168	<b>0.003</b>	-0.230	<b>&lt; 0.001</b>	0.063	0.276	-0.032	0.580

BTMs – bone turnover markers, BMD – bone mineral density, BMI: body mass index, PT – prothrombin time, APTT – activated partial thromboplastin time, INR – international normalized ratio, PLT – platelets, MPV – mean platelet volume, β-CTX – β-cross-linked C-terminal telopeptide of type I, 25-(OH)D – 25-hydroxyvitamin D, PINP – procollagen type I N propeptide. **Boldface** indicates p-value < 0.05.

### Comparison of bone mineral density characteristics

Patients were divided into three groups according to the established osteoporosis diagnostic criteria: 46 cases were assigned to the normal group, 129 cases to the osteopenia group, and 131 cases to the osteoporosis group (Table I). Except for the prevalence of hypertension ( $p = 0.205$ ), significant differences were observed in age, BMI, and gender among the three groups (all  $p < 0.001$ ). In terms of coagulation, several indicators demonstrated significant differences among the three groups, including PT, fibrinogen, INR, D-dimer, and PLT (all  $p < 0.05$ ); in contrast, APTT and MPV showed no significant difference (mean platelet volume,  $p = 0.772$ ,  $p = 0.672$ , respectively). Consistently, procollagen type I N propeptide (PINP, a bone formation biomarker) and  $\beta$ -cross-linked C-terminal telopeptide of type I ( $\beta$ -CTX, a bone resorption biomarker) showed significant differences and increasing trends across the three groups ( $p = 0.014$ ,  $p = 0.013$ , respectively). Furthermore, the levels of 25-(OH)D and osteocalcin exhibited similar patterns across the three groups ( $p = 0.336$ ,  $p = 0.088$ , respectively). However, significant differences in calcium levels were observed among the three groups ( $p = 0.045$ ). Notably, BMD exhibited significant differences in various anatomical regions among the three groups, including the L1-4 of the lumbar spine, femoral neck, Ward's triangle, and total hip (all  $p < 0.001$ ).

### Correlation of coagulation biomarkers with BTMs and BMD

Spearman's rank correlation coefficient values are presented in Table II. Positive correlations were observed between age and bone mass stage with PT, INR, and D-dimer (all  $p < 0.05$ ). Furthermore, a significant positive association was found between bone mass status and fibrinogen ( $r = 0.169$ ,  $p = 0.003$ ). Within the osteopenia group, age was significantly correlated with D-dimer ( $r = 0.298$ ,  $p = 0.001$ , Supplementary Table SII). In the osteoporosis group, age was positively correlated with PT, INR, and D-dimer levels (all  $p < 0.05$ , Supplementary Table SII).

In addition, a positive correlation was observed between 25-(OH)D and PT, INR, and PLT within the context of the association between coagulation biomarkers and BTMs (Table II). However, an inverse relationship was found between 25-(OH)D, APTT, and MPV. Moreover, calcium was significantly negatively correlated with PT, APTT, INR, and D-dimer, while it was positively correlated with PLT (all  $p < 0.05$ ). The relationship between coagulation biomarkers and BTMs in the three groups is presented in Supplementary Table SIII.

Furthermore, the relationship between coagulation biomarkers and BMD was explored, revealing negative correlations between PT, fibrinogen, INR, and D-dimer and BMD at various anatomical sites, including the lumbar spine, femoral neck, Ward's triangle, and total hip. Importantly, these correlations were statistically significant (all  $p < 0.05$ , Table II), except for the association between femoral neck, PT, and INR. Within each group, only the BMD of the total hip in the osteoporosis group displayed a negative correlation with the D-dimer levels ( $r = -0.207$ ,  $p = 0.018 < 0.05$ , Supplementary Table SIV).


### Canonical correlation analysis

To identify the most influential factors affecting coagulation, patient information was divided into three categories as independent variables. The patients' basic characteristics (age, BMI, sex, stage of bone mass, and hypertension) were denoted as X1, the BTMs ( $\beta$ -CTX, 25-(OH)D, PINP, and osteocalcin) were denoted as X2, and BMD at various sites (lumbar spine 1-4, femoral neck, Ward's triangle, and total hip) was denoted as X3. Three canonical functions were constructed using common coagulation evaluation indices as the dependent variable Y (Figure 1). Finally, the most highly correlated canonical correlation coefficients for the three canonical functions were 0.328 (X1 and Y,  $p = 0.007$ ), 0.370 (X2 and Y,  $p < 0.001$ ), and 0.268 (X3 and Y,  $p = 0.048$ ), respectively (Table III, Supplementary Table SV).

Highly influential variables were selected based on the structural coefficients. The structural coefficients of the three canonical functions are displayed in Figure 1. In the first canonical function, age showed the largest contribution to the canonical variable X1 ( $r = -0.811$ ), while fibrinogen ( $r = -0.670$ ) and PLT ( $r = 0.629$ ) contributed the most to the canonical variable Y. Moreover, the largest contributions to the canonical variable X2 were attributed to osteocalcin and 25-(OH)D ( $r = -0.882$  and  $0.840$ , respectively), and the most relevant contribution of the related canonical variable Y was INR ( $r = 3.673$ ). The BMD of Ward's triangle ( $r = -0.843$ ) was the key factor influencing the canonical variable X3, and fibrinogen ( $r = 0.788$ ) was the most essential element defining the canonical variable Y in the third canonical function.

### Logistic regression analysis of the characteristics, BTMs, BMD, and coagulation biomarkers

Based on the above statistical results, age and stage of bone mass were included as independent variables in X1 for the logistic regression analysis, 25-(OH)D in X2, and all variables in X3. Subsequently, the variables INR, fibrinogen, PLT, and D-dimer were transformed into binary variables



**Figure 1.** Canonical correlation analysis of patient information and coagulation biomarkers. **A** – Correlation coefficients between patients’ characteristics and coagulation biomarkers. **B** – Correlation coefficients between bone turnover markers and coagulation biomarkers. **C** – Correlation coefficients between bone mineral density and coagulation biomarkers

BMI – body mass index, PT – prothrombin time, APTT – activated partial thromboplastin time, FIB – fibrinogen, INR – international normalized ratio, PLT – platelets, MPV – mean platelet volume,  $\beta$ -CTX –  $\beta$ -cross-linked C-terminal telopeptide of type I, 25-(OH)D – 25-hydroxyvitamin D, PINP – procollagen type I N propeptide.

based on whether their levels were within the normal range. These binary variables served as the dependent variables in the analysis. The data underwent initial univariate logistic regression analysis (Table IV, Supplementary Tables SVI–VIII).

The multivariate analysis of INR, fibrinogen, and PLT did not reveal any significant differences ( $p > 0.05$ , Supplementary Tables SVI–VIII). However, the multivariate analysis of D-dimer identified two statistically significant variables, namely age

**Table III.** Canonical correlation analysis of the three sets of variables

Variables	Canonical correlation coefficient	Wilk's	F	P-value
X1 and Y	0.328	0.822	1.694	<b>0.007</b>
X2 and Y	0.370	0.717	2.917	<b>&lt; 0.001</b>
X3 and Y	0.268	0.870	1.496	<b>0.048</b>

X1 indicates patients' basic characteristics (age, body mass index, sex, stage of bone mass, and hypertension). X2 indicates the bone turnover markers ( $\beta$ -cross-linked C-terminal telopeptide of type I, 25-hydroxyvitamin D, procollagen type I N propeptide, osteocalcin, and calcium). X3 indicates bone mineral density in various anatomical regions (lumbar spine 1-4, femoral neck, Ward's triangle, and total hip). The dependent variable Y indicates coagulation biomarkers (prothrombin time, activated partial thromboplastin time, fibrinogen, international normalized ratio, platelets, and mean platelet volume). **Boldface** indicates  $p$ -value  $< 0.05$ .

(OR = 1.046,  $p = 0.007$ ) and the BMD of Ward's triangle (OR = 0.004,  $p = 0.017$ , Table IV).

### Discussion

Despite the implementation of various interventions aimed at addressing cardiovascular events, these conditions remain common in individuals of middle and advanced age [22]. Additionally, particular attention should be paid to perioperative thrombosis and bleeding risks in elderly patients undergoing surgical procedures [23]. This study explored the factors affecting blood coagulation in a specific population over 50 years old. The results indicated statistically significant differences in coagulation biomarkers, including PT, fibrinogen, INR, D-dimer, and PLT, among the normal, osteopenia, and osteoporosis groups (all  $p < 0.05$ ). Given the increased risk of organ dysfunction and mortality in older individuals, bleeding presents a major surgical challenge [24]. Therefore, patients aged over 50 undergoing surgery should be provided with additional measures to optimize perioperative management. Identifying predictors of bleeding may facilitate tailored individual interventions before surgery, enhancing patient safety and reducing the need

for blood transfusions [25, 26]. Moreover, individuals with abnormal bone mass should receive additional hematological monitoring in their daily lives to mitigate the risk of spontaneous bleeding or thrombotic events.

Furthermore, correlation coefficient analysis revealed a significant positive correlation between age and bone mass stage with PT, INR, and D-dimer (all  $p < 0.05$ ). Notably, D-dimer exhibited the most significant positive association with age and the most pronounced negative correlation with Ward's triangle bone mineral density. Additionally, among the bone metabolism indicators, only 25-(OH)D was related to coagulation biomarkers.

CCA involves the multivariate evaluation of the interaction between two groups of variables, each of which contains multiple variables [27]. This method enables the exploration of complex interactions and the determination of the significance of each variable's contribution [28]. In this study, a CCA approach was employed to investigate the canonical correlation between seven coagulation indicators (Y) and patient characteristics (X). The study aimed to comprehensively assess patient information, which was categorized into three distinct groups: patients' fundamental characteristics (X1, five variables), BTMs (X2, five variables), and BMD of various

**Table IV.** Logistic regression analysis between D-dimer and characteristics, bone turnover markers, and bone mineral density

Variables	Univariate analysis		Multivariate analysis	
	OR (95% CI)	P-value	OR (95% CI)	P-value
Age	1.069 (1.041–1.099)	<b>&lt; 0.001</b>	1.046 (1.012–1.081)	<b>0.007</b>
Stage of bone mass:				
Normal	Reference			
Osteopenia	0.271 (0.121–0.606)	<b>0.001</b>		
Osteoporosis	0.383 (0.227–0.646)	<b>&lt; 0.001</b>		
25(OH)D	0.999 (0.990–1.008)	0.829		
Calcium	0.090 (0.012–0.694)	<b>0.021</b>		
Lumbar spine	0.158 (0.044–0.564)	<b>0.004</b>		
Femoral neck	0.016 (0.002–0.112)	<b>&lt; 0.001</b>		
Ward's triangle	0.008 (0.001–0.054)	<b>&lt; 0.001</b>	0.004 (0.000–0.360)	<b>0.017</b>
Total hip	0.037 (0.007–0.203)	<b>&lt; 0.001</b>		

25-(OH)D – 25-hydroxyvitamin D, OR – odds ratio, CI – confidence interval; **boldface** indicates  $p$ -value  $< 0.05$ .

anatomical regions (X3, four variables). Based on the correlation and CCA analysis results, INR, fibrinogen, PLT, and D-dimer were selected as dependent variables, whereas age, stage of bone mass, 25-(OH)D, calcium, and BMD in four anatomical regions were designated as independent variables.

The results of the logistic regression analysis revealed that advanced age significantly influenced D-dimer levels. Generally, the elderly are more susceptible to thromboembolism due to higher coagulation activation and a predisposition to a 'prethrombotic state', characterized by elevated coagulation activation peptide levels and a weakened fibrinolytic system [29]. However, despite D-dimer being a widely used marker for various thrombotic conditions, its levels tend to naturally increase with age, particularly in individuals over 80; thus, its diagnostic value is limited to a certain patient population [30, 31]. Therefore, further studies are warranted to determine the potential predictive value of age for thrombosis in the elderly population.

Ward's triangle refers to a radiolucent area between the principal compressive, secondary compressive, and primary tensile trabeculae in the femoral neck [32]. Typically, this area exhibits the lowest BMD within the femoral head. Interestingly, the findings of our study indicated that a decrease in BMD within Ward's triangle may be correlated with a reduced risk of thrombosis. After controlling for other variables, no significant link was found between D-dimer levels and BMD in other anatomical regions. Consequently, elderly individuals with osteoporosis may be exposed to a higher risk of spontaneous bleeding. As reported by Zhu *et al.* [10], increased local bone density was associated with reduced bleeding during surgery. The variability in bleeding may be attributed to the rich intraosseous vascular conditions within Ward's triangle. Hence, despite the diminished coagulation function observed in individuals with osteoporosis, the extent of bleeding remains limited. However, D-dimer lacks specificity and can be elevated in various disorders [33]. Therefore, additional clinical indicators are required to assess the need for appropriate interventions [34].

Among BTMs, vitamin D is thought to significantly influence vascular physiology via direct effects on endothelial cells and indirect pathways, including endocrine and immunomodulatory mechanisms [35]. Nevertheless, our regression analysis revealed that the 25-(OH)D levels showed no significant effect on the odds ratio of coagulation biomarkers (all  $p > 0.05$ ), aligning with certain previous studies [36, 37]. The role of vitamin D in cardiovascular disorders remains controversial [38]. Besides, calcium is not only a biomarker of bone metabolism but is also involved in the intrinsic and extrinsic pathways of blood coagulation.

Previous epidemiological studies reported an association between calcium levels and vascular risk [39]. The present correlation research also showed that serum calcium was negatively correlated with PT, APTT, and D-dimer, and positively correlated with platelets. These findings supported other research results indicating that an increase in calcium concentration can shorten coagulation time [39, 40]. This may be related to platelets expressing the calcium-sensing receptor, with changes in serum calcium potentially influencing blood coagulability [41]. However, the multivariate regression analysis of this study indicated that calcium and D-dimer showed no statistically significant association with coagulation. Nonetheless, the lack of association might be attributed to the plasma being collected into calcium chelator-containing tubes for laboratory analysis; the samples would be unsuitable for examining the influence of physiological changes in serum calcium on whole-blood coagulation [42]. Therefore, a new method for measuring blood coagulation is required to comprehensively evaluate the coagulation system in natural whole blood, to clarify the relationship between calcium and coagulation.

This study presents the first comprehensive analysis of the relationship between coagulation biomarkers and patient characteristics, BTMs, and BMD. Nevertheless, the limitations of this study should be acknowledged. First, this single-center study employed a retrospective design, potentially introducing confounding factors, selection biases, and disparities. Second, the relatively small sample size could introduce bias into our analysis. Third, our study exclusively involved Chinese subjects, limiting the direct generalizability of our findings to other populations. Therefore, larger, multicenter prospective studies are required to validate the predictive ability of BTMs and BMD for coagulation disorders in individuals over 50 years of age. In addition, considering the inherent complexity of *in vivo* hemostasis, our exploration of hemostasis via coagulation biomarker measurements only provides an estimation of the relationship. Finally, due to the relatively weak correlations found in this study, and the presence of multiple confounding factors, further high-quality research is needed to confirm the association between bone mass status and coagulation biomarkers.

In conclusion, for the Chinese population aged 50 and above, attention should be paid to changes in coagulation biomarkers.

## Acknowledgments

Zhengming Wang and Rui Wang contributed equally to this work and share first authorship.

This work was supported by the Shanghai Municipal Health Commission (20MC1920600, shsl-



czdzk03901, and ZY(2021-2023)-0209-02), the National Natural Science Foundation (82074466 and 82174403), and Shanghai University of Traditional Chinese Medicine (GJ2023017).

We thank the Home for Researchers editorial team ([www.home-for-researchers.com](http://www.home-for-researchers.com)) for language editing service.

### Conflict of interest

The authors declare no conflict of interest.

### References

- Compston JE, McClung MR, Leslie WD. Osteoporosis. *Lancet* 2019; 393: 364-76.
- Johnston CB, Dagar M. Osteoporosis in older adults. *Med Clin North Am* 2020; 104: 873-84.
- Aspray TJ, Hill TR. Osteoporosis and the ageing skeleton. *Subcell Biochem* 2019; 91: 453-76.
- Reid IR. A broader strategy for osteoporosis interventions. *Nat Rev Endocrinol* 2020; 16: 333-9.
- Johnell O, Kanis JA. An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporos Int* 2006; 17: 1726-33.
- Florschutz AV, Langford JR, Haidukewych GJ, Koval KJ. Femoral neck fractures: current management. *J Orthop Trauma* 2015; 29: 121-9.
- Prost S, Pesenti S, Fuentes S, Tropiano P, Blondel B. Treatment of osteoporotic vertebral fractures. *Orthop Traumatol Surg Res* 2021; 107: 102779.
- Singleton M, Ghisi D, Memtsoudis S. Perioperative management in complex spine surgery. *Minerva Anestesiol* 2022; 88: 396-406.
- Wang X, Xu T, Wang R, et al. Perioperative blood loss can be reduced if total knee arthroplasty was performed in the si hour-period, compared with the wei hour-period: a retrospective cohort study. *Evid Based Complement Alternat Med* 2021; 2021: 9990413.
- Zhu W, Wei Z, Zhou T, et al. Bone density may be a promising predictor for blood loss during total hip arthroplasty. *J Clin Med* 2022; 11: 3951.
- Dorgalaleh A, Favaloro EJ, Bahraini M, Rad F. Standardization of prothrombin time/international normalized ratio (PT/INR). *Int J Lab Hematol* 2021; 43: 21-8.
- van Ommen CH, Peters M. The bleeding child. Part I: primary hemostatic disorders. *Eur J Pediatr* 2012; 171: 1-10.
- Wang X, Yang Y, Yu L, et al. Association between fibrinogen level and length of stay in patients with lower extremity atherosclerotic disease: a retrospective cohort study. *Sci Rep* 2023; 13: 11872.
- Kleinegris MC, ten Cate H, ten Cate-Hoek AJ. D-dimer as a marker for cardiovascular and arterial thrombotic events in patients with peripheral arterial disease. A systematic review. *Thromb Haemostasis* 2013; 110: 233-43.
- Hayward CP, Moffat KA, Liu Y. Laboratory investigations for bleeding disorders. *Semin Thromb Hemost* 2012; 38: 742-52.
- Sang Y, Roest M, de Laat B, de Groot PG, Huskens D. Interplay between platelets and coagulation. *Blood Rev* 2021; 46: 100733.
- Kanis JA. Assessment of fracture risk and its application to screening for postmenopausal osteoporosis: synopsis of a WHO report. WHO Study Group. *Osteoporos Int* 1994; 4: 368-81.
- Liu Q, Dong H, Li Y, et al. Apolipoprotein-A is a potential prognostic biomarker for severe aplastic anemia patients treated with ATG-based immunosuppressive therapy: a single-center retrospective study. *Lipids Health Dis* 2022; 21: 93.
- Shi Z, Jin H, Ding Q, et al. Bone turnover markers may predict the progression of osteonecrosis of the femoral head in aged males. *Ann Transl Med* 2019; 7: 626.
- Koebnick C, Black MH, Wu J, et al. A diet high in sugar-sweetened beverage and low in fruits and vegetables is associated with adiposity and a pro-inflammatory adipokine profile. *Br J Nutr* 2018; 120: 1230-9.
- Ridderstolpe L, Gill H, Borga M, Rutberg H, Ahlfeldt H. Canonical correlation analysis of risk factors and clinical outcomes in cardiac surgery. *J Med Syst* 2005; 29: 357-77.
- Walli-Attai M, Rosengren A, Rangarajan S, et al. Metabolic, behavioural, and psychosocial risk factors and cardiovascular disease in women compared with men in 21 high-income, middle-income, and low-income countries: an analysis of the PURE study. *Lancet* 2022; 400: 811-21.
- Xu D, Hu X, Zhang H, et al. Analysis of risk factors for deep vein thrombosis after spinal infection surgery and construction of a nomogram preoperative prediction model. *Front Cell Infect Microbiol* 2023; 13: 1220456.
- Dawson D, Milligan D, Callachand F, Cusick L. Hip hemi-arthroplasty vs total hip replacement for displaced intra-capsular hip fractures: retrospective age and sex matched cohort study. *Ulster Med J* 2018; 87: 17-21.
- Wan S, Sparring V, Cabrales DA, Jansson K, Wikman A. Clinical and budget impact of treating preoperative anemia in major orthopedic surgery-a retrospective observational study. *J Arthroplasty* 2020; 35: 3084-8.
- Belay ES, Penrose CT, Ryan SP, Bergen MA, Bolognesi MP, Seyler TM. Perioperative selective serotonin reuptake inhibitor use is associated with an increased risk of transfusion in total hip and knee arthroplasty. *J Arthroplasty* 2019; 34: 2898-902.
- Lu G, Dong Z, Huang B, et al. Determination of weight loss effectiveness evaluation indexes and establishment of a nomogram for forecasting the probability of effectiveness of weight loss in bariatric surgery: a retrospective cohort. *Int J Surg* 2023; 109: 850-60.
- Li J, Gao W, Zuo W, Liu X. Relationship between quality of life and social support in parents of children with congenital anorectal malformations in China. *J Pediatr Nurs* 2020; 53: e87-92.
- Nurmohamed MT, Büller HR, ten Cate JW. Physiological changes due to age. Implications for the prevention and treatment of thrombosis in older patients. *Drugs Aging* 1994; 5: 20-33.
- Bicciè FG, Farcomeni A, Gaudio C, Pignatelli P, Tanzilli G, Pastori D. D-dimer for risk stratification and antithrombotic treatment management in acute coronary syndrome patients: asystematic review and metanalysis. *Thromb J* 2021; 19: 102.
- Zhao X, Li J, Tang X, et al. D-dimer as a thrombus biomarker for predicting 2-year mortality after percutaneous coronary intervention. *Ther Adv Chronic Dis* 2020; 11: 2040622320904302.
- Pei YF, Hu WZ, Hai R, et al. Genome-wide association meta-analyses identified 1q43 and 2q32.2 for hip Ward's triangle areal bone mineral density. *Bone* 2016; 91: 1-10.
- Innocenti F, Lazzari C, Ricci F, Paolucci E, Agishev I, Pini R. D-dimer tests in the emergency department: current insights. *Open Access Emerg Med* 2021; 13: 465-79.

34. Wu MA, Bizzi E, Brucato A. Inflammatory puzzle: unveiling the role of D-Dimer in acute pericarditis and thromboinflammation. *Eur J Intern Med* 2023; 116: 41-2.
35. Towler DA. Calcitropic hormones and arterial physiology: "D"-lightful insights. *J Am Soc Nephrol* 2007; 18: 369-73.
36. Blondon M, Cushman M, Jenny N, et al. Associations of serum 25-hydroxyvitamin D with hemostatic and inflammatory biomarkers in the multi-ethnic study of atherosclerosis. *J Clin Endocrinol Metab* 2016; 101: 2348-57.
37. Hyppönen E, Berry D, Cortina-Borja M, Power C. 25-Hydroxyvitamin D and pre-clinical alterations in inflammatory and hemostatic markers: a cross sectional analysis in the 1958 British Birth Cohort. *PLoS One* 2010; 5: e10801.
38. Pelczyńska M, Grzelak T, Sperling M, Bogdański P, Papek-Musiałik D, Czyżewska K. Impact of 25-hydroxyvitamin D, free and bioavailable fractions of vitamin D, and vitamin D binding protein levels on metabolic syndrome components. *Arch Med Sci* 2017; 13: 745-52.
39. Reid IR. Cardiovascular effects of calcium supplements. *Nutrients* 2013; 5: 2522-9.
40. Shin S, Kim KJ, Chang HJ, et al. Impact of serum calcium and phosphate on coronary atherosclerosis detected by cardiac computed tomography. *Eur Heart J* 2012; 33: 2873-81.
41. House MG, Kohlmeier L, Chattopadhyay N, et al. Expression of an extracellular calcium-sensing receptor in human and mouse bone marrow cells. *J Bone Miner Res* 1997; 12: 1959-70.
42. James MF, Roche AM. Dose-response relationship between plasma ionized calcium concentration and thrombelastography. *J Cardiothorac Vasc Anesth* 2004; 18: 581-6.