

Diagnostic accuracy of non-invasive investigations for coronary artery disease in hypertensive patients: a meta-analysis

Ibadete Bytyci^{1,2,3}, Sefer Bytyqi⁴, Gani Abazi⁵, Gani Bajraktari^{1,2,3}, Michael Y. Henein^{3,6*}

¹Clinic of Cardiology, University Clinical Centre of Kosovo, Prishtina, Kosovo

²Medical Faculty, University of Prishtina, Prishtina, Kosovo

³Institute of Public Health and Clinical Medicine, Umeå University, Sweden

⁴Department of Data Science, Riinvest College, Prishtina, Kosovo

⁵Harvard University, Boston, USA

⁶Molecular and Clinic Research Institute, St George University, London, and Brunel University, UK

***Corresponding author:**

Prof. Michael Y. Henein
Institute of Public Health
and Clinical Medicine
Umeå University
Umeå, Sweden
Phone: +46 90 785 14 31
E-mail: michael.henein@
umu.se

Submitted: 27 April 2024; **Accepted:** 15 May 2024

Online publication: 6 June 2024

Arch Med Sci 2025; 21 (1): 167–178

DOI: <https://doi.org/10.5114/aoms/188782>

Copyright © 2024 Termedia & Banach

Abstract

Introduction: Non-invasive assessment of coronary artery disease (CAD) in patients with hypertensive heart disease is still a major clinical challenge. The aim of this meta-analysis was to evaluate the diagnostic accuracy of different non-invasive techniques in detecting significant (> 50%) CAD in hypertensive patients.

Material and methods: We systematically searched selected electronic databases from inception until February 27th 2024. The main clinical endpoint was the diagnostic accuracy of non-invasive investigations including exercise electrocardiography test (EET), stress echocardiography (SE) and myocardial perfusion imaging (MPI). Random effects summary receiver operating characteristic analysis was performed.

Results: Twenty-five papers with a total of 3812 patients with systemic hypertension and suspected or known significant CAD were finally included in the meta-analysis. The diagnostic accuracy of SE was the highest compared to the two other investigations (85%, 72%, 53%, $p < 0.05$ for all). SE had higher sensitivity 85% (83–87%) and specificity 81% (79–83%) compared to MPI (sensitivity 81% (77–83%), specificity 64% (59–68%)) and EET (sensitivity: 53% (50–65%) and specificity 51% (47–54%)). All three tested investigations had higher accuracy in patients with multivessel disease compared to those with single-vessel disease ($p < 0.05$ for all). The diagnostic accuracy of SE and MPI was not significantly impacted by left ventricular hypertrophy ($p > 0.05$ for both) compared to EET, which was lower in patients with hypertrophy compared to those without (41% vs. 58%, $p = 0.03$). These results were consistent irrespective of the stress echo modality, physical, exercise or pharmacological ($p > 0.05$).

Conclusions: This meta-analysis reveals the powerful diagnostic value of stress echocardiography in detecting significant coronary artery disease in hypertension patients.

Key words: non-invasive methods, stress echocardiography, myocardial perfusion, stress electrocardiography, arterial hypertension, coronary artery disease.

Introduction

Coronary artery disease (CAD) is the leading cause of morbidity and mortality worldwide [1]. Arterial hypertension (AH) is a well-established risk factor for the development and progression of atherosclerosis and CAD [2]. The pathophysiology of hypertension involves chronic mechanical stress on the arterial wall, leading to increased shear stress, turbulence, and oscillatory shear index. Together with the other conventional risk factors, these processes contribute to the formation of atherosclerotic lesions and subsequent development of CAD [1, 3]. Assessing the prognosis of hypertensive patients is crucial in clinical practice because it is closely associated with a two-fold increase in the risk of developing CAD. Hence, prevention, detection, and management of AH are essential in reducing its burden on society and in improving patient outcome [4]. Non-invasive diagnostic investigations of CAD play a crucial role in the evaluation and optimization of clinical management of these patients [5, 6]. However, the exercise electrocardiography test (EET) faces a significant challenge due to its high rate of false-positive responses. This is primarily due to reduced coronary flow reserve, even when obstructive CAD is not present [7, 8]. On the other hand, functional tests in the form of stress echocardiography (SE) and myocardial perfusion imaging (MPI) provide superior diagnostic accuracy with higher sensitivity and specificity for detecting significant CAD [9–11]. Although new diagnostic techniques and advanced therapeutic approaches have been developed, the non-invasive diagnosis of CAD in hypertensive patients often raises uncertainties, mostly due to left ventricular hypertrophy (LVH) [12]. This does not preclude the important role of the recently developed high accuracy coronary computed tomography angiography (CCTA) [13, 14]. The aim of this meta-analysis was to evaluate the diagnostic accuracy of different non-invasive functional investigations in detecting significant (> 50%) CAD in hypertensive patients.

Material and methods

Search strategy and selection criteria

We adhered to the 2020 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [15]. As our study was a meta-analysis, obtaining Institutional Review Board (IRB) approval or patient informed consent was unnecessary.

To shape the clinical question and develop the search strategy, we utilized a PECOS model which included parameters for population, intervention, comparison, outcomes, and study design (Supplementary Table S1).

The following databases were searched until February 27th, 2024: PubMed-Medline, EMBASE, Scopus, Google Scholar, the Cochrane Central Registry of Controlled Trials, and ClinicalTrials.gov. The following keywords were used: “Coronary artery disease” OR “CAD” OR “Ischemic heart disease” OR “IHD” AND “Arterial hypertension” AND “Non-invasive diagnostic” OR “Stress echocardiography” OR “Stress echo” OR “Dipyridamole echocardiography” OR “Dobutamine stress echocardiography” OR “Exercise electrocardiography test” OR “EET” OR “Myocardial perfusion imaging” OR “MPI” AND “Accuracy” (Supplementary Table SII).

Additional searches for potential studies were conducted by reviewing the references of review articles and abstracts from relevant congresses such as the scientific sessions of the European Society of Cardiology (ESC), the European Atherosclerosis Society (EAS), the American Heart Association (AHA), the American College of Cardiology (ACC), and the European Association of Cardiovascular Imaging (EACVI). The wild-card term “*” was used to increase the search sensitivity. The search was limited to human studies published in English, without any applied filters. Each article was evaluated independently by two reviewers (S.B. and I.B.). When there was uncertainty about the suitability of a paper the senior investigator (M.Y.H.) was consulted. The remaining articles were assessed in full text by the same two researchers who extracted and analyzed the data. Usually, data were compared with the original articles and then errors were corrected. The Newcastle-Ottawa Scale (NOS) tool was used to assess the risk of bias in each study. This tool assesses three domains: selection, compatibility and exposure of reported results. The investigators judged the risk of bias in each study as “low”, “high”, or “unclear” [16, 17]. Articles were considered eligible if they reported the diagnostic accuracy in patients with AH: a) studies investigating diagnostic accuracy of non-invasive methods in AH patients; b) studies reporting diagnostic accuracy of significant CAD and/or in comparison of left ventricle hypertrophy or number of diseased vessels; and c) enrolled population of adults aged ≥ 18 years. Exclusion criteria were: i) insufficient statistical data to test the diagnostic accuracy, ii) patients without AH, iii) studies not in humans, and iv) ongoing studies (unless they had reported relevant interim results).

Outcome variables

The main clinical endpoint was the diagnostic accuracy of the non-invasive functional investigations including exercise electrocardiography test (EET), stress echocardiography (SE) and myocardial perfusion imaging (MPI). The secondary endpoint was the effect of left ventricular hyper-

trophy (LVH), number of diseased vessels and the different CAD territories on diagnostic accuracy. In addition, we evaluated the diagnostic accuracy of SE in patients using two stress modalities: pharmacologic and exercise SE. Moreover, we assessed the accuracy of dobutamine (Dob) compared with dipyridamole (Dip) stress echocardiography.

Significant CAD was defined as $\geq 50\%$ coronary luminal stenosis of at least one major epicardial vessel on conventional coronary angiography. Multivessel disease (MVD) was defined as luminal stenosis in at least two major coronary arteries or in one coronary artery in addition to $\geq 50\%$ stenosis of the left main stem.

Data synthesis and statistical analysis

The meta-analysis was conducted using RevMan software (Review Manager [RevMan] Version 5.1, The Cochrane Collaboration, Copenhagen, Denmark), with two-tailed $p < 0.05$ considered significant. The baseline characteristics are reported as median and range. Mean and standard deviation (SD) values were estimated using the method described by Hozo *et al.* [18]. Forest plots were used as the standard method to present the results of individual studies and meta-analyses. We used hierarchical summary receiver operating characteristic (ROC) analysis with the Rutter and Gatsonis model to assess the baseline diagnostic accuracy of the non-invasive investigations [19]. The diagnostic random-effects model was used to compute summary sensitivity and specificity, along with 95% confidence intervals, based on true positive (TP), true negative (TN), false positive (FP), and false negative (FN) values. The summary point obtained from the hierarchical ROC analysis was then used to calculate the positive likelihood ratio (LR+), negative likelihood ratio (LR-), positive predictive value (PPV), negative predictive value (NPV), and diagnostic odds ratio (DOR). For studies that did not provide optimal cut-off values, we created the ROC curve and identified the optimal cut-off as the point on the ROC curve closest to 0.1 in x - y coordinates [19]. The potential interaction between demographic and clinical indices with accuracy of non-invasive investigations was further explored by meta-regression. Meta-regression coefficients and corresponding p -values are reported. We used the random-effects model for the meta-analyses, and heterogeneity between studies was evaluated using the Cochrane Q test and I^2 index. $I^2 < 25\%$ indicated low, 25–50% moderate, and $> 50\%$ high heterogeneity. To assess the additive (between-study) component of variance, the reduced maximum likelihood method (tau2) incorporated the occurrence of residual heterogeneity into the analysis [20]. Publication bias was assessed using visual inspections of funnel plots and Egger's test.

Results

Characteristics of included studies

The initial search identified 2468 articles, which were screened for relevance, resulting in 1278 articles. After a stringent selection process, the meta-analysis included 25 studies comprising 3812 hypertensive patients with known or suspected CAD [21–45] (Supplementary Figure S1). Diagnostic accuracy in 10 studies was evaluated only by one method [21, 23, 25, 30, 32, 35, 40, 42, 44, 45] and the remaining 15 papers reported two or more diagnostic methods [22, 24, 26, 27, 28, 29, 31, 33, 34, 36, 37, 38, 39, 41, 43]. Fourteen studies reported LVH [21, 23, 24, 26, 28, 29, 30, 31, 33, 34, 41, 42, 43, 45] and 9 reported the number of diseased vessels and/or the different territories of CAD [26, 27, 29, 30, 34, 35, 36, 42, 43]. The characteristics of the included studies are shown in Table I.

The mean age of patients was 52.2 ± 7.8 years and 44.5% were female. 51.8% of the patients had CAD, 42.3% were smokers, 19.7% diabetic and 37.2% had dyslipidemia (Supplementary Table SIII).

Diagnostic accuracy of non-invasive methods in detecting significant CAD

SE with a summary sensitivity of 87%, summary specificity 81%, and 85% accuracy was the most powerful predictor of significant CAD followed by MPI with summary sensitivity of 81%, summary specificity of 64% and accuracy of 72%. EET was the investigation with the lowest accuracy with summary sensitivity of 53%, summary specificity of 51% and accuracy of 53% (Figures 1, 2). Positive and negative predictive values were, respectively, 82% and 87% for SE, 71% and 76% for MPI and 56% and 48% for EET (Table II). The predictive value of significant CAD for SE measured by different modalities (pharmacologic vs. physical exercise) was also assessed. The results showed no difference between the accuracy of the two techniques (84% vs. 90%; $p = \text{NS}$; Figure 3 A, Supplementary Figure S2). Although dobutamine SE had modestly higher summary sensitivity (85% vs. 80%) and lower summary specificity (81% vs. 91%) compared to dipyridamole SE, the accuracy was not different between the two SE modalities (83% vs. 85% Figure 3 B, Supplementary Figure S2).

Effect of left ventricular hypertrophy on diagnostic accuracy of non-invasive methods

Left ventricular hypertrophy did not have a significant impact on the diagnostic accuracy of the

Table I. Main characteristics of studies included in the meta-analysis

Study (trial year)	Study design	Location	Population	Sample size	Non-invasive modalities	LV hypertrophy	Prediction [months]
Wroblewski 1982	Retrospective observational study	USA	AH with CAD suspected	37	EET	R	Significant SCAD
Picano 1988 [22]	Prospective observational study	Italy	AH with CAD suspected	63	EET Exercise echocardiography	NR	Significant CAD
Prisant 1992 [23]	Prospective observational study	USA	AH with CAD suspected	92	MPI	R	Significant CAD
Massie 1993 [24]	Prospective observational study	USA	AH with CAD suspected	226	EET MPI	R	Significant CAD
Tsuda 1993 [25]	Prospective observational study	Japan	AH with CAD suspected	51	EET	NR	Significant CAD
Senior 1996 [26]	Prospective observational study	UK	AH with CAD suspected	43	EET Dob echocardiography	R	Significant CAD MVD
Schillaci 1997 [27]	Prospective observational study	Italy	AH with CAD suspected	40	EET Dip echocardiography MPI	NR	Significant CAD MVD
Astarita 1998 [28]	Prospective observational study	Italy	AH with CAD suspected	42	Dip echocardiography MPI	R	Significant CAD
Elhendy 1998 [29]	Prospective observational study	Egypt Netherlands	AH with CAD suspected	84	Dob echocardiography MPI	R	Significant CAD MVD
Ho 1998 [30]	Prospective observational study	Taiwan	AH with known CAD suspected	101	Dob echocardiography	R	Significant CAD MVD
Fragasso 1999 [31]	Prospective observational study	Italy	AH with CAD suspected	101	Dob echocardiography Dip echocardiography MPI	R	Significant CAD
Arrif 2000 [32]	Prospective observational study	UK	AH with CAD suspected	30	Dob echocardiography	NR	Significant CAD
Maltagliati 2000 [33]	Prospective observational study	Italy	AH with CAD suspected	59	EET Exercise echocardiography	R	Significant CAD
Pasierski 2001 [34]	Prospective observational study	Poland	AH with CAD suspected	197	EET Dob echocardiography Exercise echocardiography	R	Significant CAD MVD
Elhendy 2001 [35]	Prospective observational study	USA	AH with CAD suspected	137	MPI	NR	Significant CAD MVD
Astarita 2001 [36]	Prospective observational study	Italy	AH with CAD suspected	53	Dip echocardiography MPI	NR	Significant CAD MVD
Cortigiani 2003 [37]	Prospective observational study	Italy	AH with CAD suspected	35	EET Dip echocardiography	NR	Significant CAD

Table I. Cont.

Study (trial year)	Study design	Location	Population	Sample size	Non-invasive modalities	LV hypertrophy	Prediction [months]
Milosavljevic 2005 [38]	Prospective observational study	Serbia	AH with CAD known or suspected	127	Dip echocardiography Dob echocardiography	NR	Significant CAD
Nishida 2005 [39]	Retrospective observational study	Japan	AH with CAD suspected	83	EET MPI	NR	Significant CAD
Michaelides 2007 [40]	Prospective observational study	Greece	AH with CAD suspected	179	MPI	NR	Significant CAD
Aggeli 2007 [41]	Prospective observational study	Greece	AH with CAD suspected	50	Stress echocardiography MPI	R	Significant CAD
Michaelides 2009 [42]	Prospective observational study	Greece	AH with CAD suspected	382	EET	R	Significant CAD MVD
Lu 2010 [43]	Prospective observational study	Italy	AH with CAD suspected	76	EET Dob echocardiography Exercise echocardiography MPI	R	Significant CAD MVD
Cortigiani 2011 [44]	Retrospective observational study	Italy	AH with CAD suspected	1411	Dob echocardiography	NR	Significant CAD
Zimarino 2016 [45]	Retrospective observational study	Italy	AH with CAD suspected	113	EET	R	Significant CAD

AH – arterial hypertension, CAD – coronary artery disease, EET – exercise electrocardiography test, Dob – dobutamine, Dip – dipyridamole, MPI – myocardial perfusion imaging, R – reported, NR – non-reported.

two stress techniques, SE and MPI. In both investigations the accuracy of the diagnostic indices was similar in patients with and without hypertrophy. The accuracy was not different in SE (85% vs. 81%, $p > 0.05$) or MPI (75% vs. 71%, $p > 0.05$). However, EET accuracy was lower in patients with LVH compared to those without (41% vs. 58%, $p = 0.03$; Figure 4 A, Supplementary Figure S3).

Effect of number of diseased vessels on diagnostic accuracy of investigations

Regardless of the non-invasive diagnostic investigation used, patients with MVD had higher sensitivity, specificity, positive predictive value, negative predictive value, and accuracy compared to those with single-vessel disease. The summary sensitivity of SE was 85%, summary specificity 86% and accuracy 86% in detecting MVD while in single vessel disease respective values were 71%, 84% and 78%. Likewise, the accuracy of MPI and EET in detecting MVD compared to single vessel disease was significantly higher (81% vs. 71% and 60% vs. 50%; $p < 0.05$ for both) (Figure 4 B, Supplementary Figure S4).

The regional accuracy of SE and MPI was similar for LAD and LCx artery stenosis, but there was a trend towards lower sensitivity for RCA stenosis. This may be due to the RCA's anatomical position, which makes it more challenging to visualize and assess accurately, potentially leading to reduced sensitivity and specificity. Additionally, collateral circulation can obscure stenosis detection, complicating accurate diagnosis with these imaging techniques (Supplementary Figure S5).

Interaction of demographic and clinical indices with accuracy of non-invasive methods

Demographic and clinical indices were tested for possible interaction with accuracy of the non-invasive investigations. Positive interactions were found for diabetes mellitus ($\beta = 0.411$; $p = 0.02$) and smoking ($\beta = 0.229$; $p = 0.03$) with sensitivity of SE and EET ($\beta = 0.416$ and $\beta = 0.299$; $p < 0.05$, respectively) and diabetes with sensitivity of MPI ($\beta = 0.281$; $p = 0.03$). A negative interaction was found only between diabetes ($\beta =$




Figure 1. Summary sensitivity and specificity of non-invasive investigations

-0.228; $p = 0.03$) and the specificity of SE. Other factors did not have any interaction with the specificity of other modalities (Figure 5, Supplementary Figures S6–S8).

Risk of bias assessment

The assessment of the risk of bias in the included studies using NOS showed that most studies had a moderate to high quality level in defining objectives and primary outcomes (Supplementary Table SIV). Also, there was no evidence of publication bias, evaluated by Egger’s test, for our findings.

Discussion

Findings

The findings of this meta-analysis, which included 25 studies and a total of 3,812 hypertensive patients with either known or suspected coronary artery disease, are as follows: a) Stress echocardiography is a superior predictor of significant (> 50%) CAD in patients with hypertension compared to the two other non-invasive functional investigations, EET and MPI. b) The accuracy of the two SE modalities was similar, even when dobutamine or dipyridamole was used. c) The accuracy of the three non-invasive diagnostic investigations was higher in patients with multivessel coronary disease compared to those with single-vessel disease. d) The diagnostic accuracy of SE and MPI was not impacted by left ventricular hypertrophy; however, that of EET was lower in patients with hypertrophy compared to those without.

Data interpretation

Despite advances in early diagnosis and treatment of AH, related morbidity and mortality rates continue to be of significant concern. Evaluating the prognosis of hypertensive patients is a critical aspect of clinical practice due to the strong relationship between hypertension and the risk of developing CAD [2, 3], even when subclinical. Accurate assessment of the diagnostic value of non-invasive investigations for detecting CAD is essential for effective management of patients with hypertension [4, 5]. Our meta-analysis shows that SE has significantly higher accuracy than myocardial perfusion imaging and exercise stress testing, and thus should provide incremental diagnostic value for the detection and localization of significant CAD.

EET is considered a potentially useful diagnostic tool for CAD screening due to its simplicity, low




Figure 2. Diagnostic accuracy of non-invasive investigations

Table II. Diagnostic accuracy of non-invasive methods in detecting significant CAD

Methods	Sensitivity	Specificity	PPV	NPV	Accuracy
EET:	53 (50 to 56)	51 (47 to 54)	56 (54 to 58)	48 (46 to 50)	53 (50 to 55)
LVH-	57 (49 to 62)	61 (53 to 70)	71 (63 to 77)	44 (37 to 52)	58 (52 to 62)
LVH+	49 (41 to 58)	40 (30 to 50)	55 (45 to 63)	29 (22 to 37)	41 (35 to 47)
Mono vessel	49 (33 to 65)	52 (40 to 64)	37 (25 to 51)	64 (50 to 76)	50 (41 to 60)
Multivessel	67 (53 to 79)	54 (61 to 75)	54 (42 to 65)	67 (53 to 78)	60 (50 to 67)
SE:	87 (85 to 89)	81 (79 to 83)	82 (78 to 83)	87 (85 to 88)	85 (82 to 87)
Pharmacologic	86 (84 to 88)	80 (78 to 83)	81 (79 to 82)	86 (83 to 88)	84 (81 to 86)
Dob	85 (80 to 87)	81 (76 to 85)	82 (78 to 84)	83 (79 to 86)	83 (79 to 85)
Dip	80 (75 to 85)	91 (87 to 94)	89 (85 to 92)	83 (80 to 86)	86 (82 to 88)
Exercise	95 (90 to 98)	87 (81 to 92)	86 (80 to 90)	95 (92 to 98)	90 (87 to 93)
LVH-	80 (71 to 86)	92 (83 to 96)	94 (84 to 97)	74 (65 to 82)	85 (78 to 89)
LVH+	77 (70 to 83)	89 (82 to 94)	92 (87 to 96)	68 (60 to 75)	81 (77 to 85)
Mono vessel	71 (63 to 78)	84 (80 to 90)	76 (68 to 82)	80 (75 to 85)	78 (73 to 82)
Multivessel	85 (77 to 89)	86 (81 to 90)	83 (77 to 88)	86 (80 to 90)	86 (81 to 80)
MPI:	81 (77 to 84)	64 (59 to 68)	71 (68 to 73)	76 (82 to 79)	72 (70 to 75)
LVH-	74 (61 to 85)	89 (74 to 96)	69 (52 to 80)	91 (79 to 97)	80 (70 to 87)
LVH+	72 (63 to 79)	81 (71 to 89)	71 (62 to 79)	81 (70 to 89)	75 (68 to 81)
Mono vessel	60 (55 to 66)	82 (77 to 87)	77 (71 to 82)	67 (62 to 71)	71 (67 to 74)
Multivessel	75 (67 to 82)	86 (80 to 91)	82 (74 to 87)	81 (75 to 86)	81 (77 to 85)

EET – exercise electrocardiography test, Dob – dobutamine, Dip – dipyridamole, MPI – myocardial perfusion imaging, LVH – left ventricular hypertrophy.




Figure 3. Diagnostic accuracy of different modalities of SE: **A** – exercise vs. pharmacologic echocardiography, **B** – dobutamine vs. dipyridamole echocardiography




Figure 4. Effect of LVH (A) and number of CAD (B) vessels on diagnostic accuracy




Figure 5. Summary interaction of demographic and clinical indices with accuracy of non-invasive methods

cost, and safety [46] with significant prognostic power. However, many studies have demonstrated its relatively poor accuracy for detecting CAD, even in symptomatic individuals due to the co-existence of bundle branch blocks, premature test termination, etc. A considerable number of hypertensive patients experience chest pain/discomfort, particularly with exercise. In these patients, EET specificity for diagnosing significant CAD has been proved limited, particularly in those with left ventricular hypertrophy and/or pre-existing ECG abnormalities [47]. In patients without significant ECG abnormalities at rest, frequently seen in hypertension clinics, diagnostic accuracy of EET is even lower [48, 49]. Such poor accuracy has led the recommendations to discourage the use of exercise testing as a screening tool [7, 47, 48]. These findings are supported by our results, which showed that EET has low accuracy in all hypertensive individuals.

Our results have shown respectable accuracy of MPI for detecting CAD, irrespective of the presence of LVH [7, 50] but still inferior to SE. Despite that, the risk of MPI associated radiation exposure, particularly in the young and in women, limits the potential for using MPI as a screening functional investigation in the general population [51]. EET and MPI assess electric and myocardial perfusion signs of ischemia, respectively, particularly transmural. Recently, the development of CCTA has added extra power to the non-invasive armamentarium of diagnosing significant CAD, with its ability to provide a detailed anatomical description of the location and severity of branch stenosis. Whenever in doubt about the stenosis grade, recent guidelines offer class 1 recommendation for functional tests for detecting myocardial ischemia, even without CCTA, in intermediate- to high-risk patients [52]. Indeed, a large study including 1000 patients with possible stable CAD demonstrated higher efficiency of functional tests including SE, positron emission tomography (PET), non-invasive fractional flow reserve (FFRCT) and cardiac magnetic resonance imaging [8] in diagnosing significant CAD. Although these results are important, their wide implementation could be limited because of the known CT-related limitations, including radiation, subjectivity, cost-effectiveness ratio, cost-utility ratio and availability, particularly in developing countries [7, 8, 21]. On the other hand, in our analysis SE was found to be the strongest predictor of significant CAD through accurate evaluation of ischemic myocardial wall motion abnormalities. Such high accuracy is irrespective of the pharmacological modality used, dobutamine or dipyridamole. Based on this finding the recommendation of generalized use of SE as a screening test in hypertensive patients can be justified, particularly because of its additional benefits: having zero

radiation, being patient friendly and avoiding possible claustrophobia of other scanning tubes. The cost-effectiveness ratio and cost-utility ratio, and the need for a certain degree of specialized knowledge, are additional benefits for SE, particularly in these patients. Despite such benefits, it must be mentioned that some of those patients presenting with exertional symptoms might not prove to have significant coronary stenosis as a cause of their symptoms but either vasospastic disease or microcirculation disease or both. While SE might show wall motion abnormalities in the former [53] it might not demonstrate significant evidence for ischemia in the latter [54], based on the limited available evidence in the literature. These findings advocate for broader clinical implementation of SE because of its cost-effectiveness, patient-friendliness, and lack of radiation exposure. This study also provides strong evidence for revising current diagnostic protocols, emphasizing the need for wider use of SE for early diagnosis of coronary artery disease and hence improvement of patient clinical outcomes.

Clinical implication

The findings of our meta-analysis have significant clinical implications. Hypertension is one of the commonest presentations in cardiology clinics and general practice, irrespective of symptoms. While age and other comorbidities may impact development of co-existing symptoms, such patients, particularly those with long standing hypertension and family history of CAD or stroke, warrant further investigations in order to optimize their clinical management and prevent them developing serious acute events. Our results support the generalized use of SE in such individuals, particularly those who cannot tolerate physical exercise and when MPI, with its special requirements, is not available. The use of more sophisticated echocardiographic modalities including speckle tracking echocardiography for the assessment of myocardial deformation is expected to increase the accuracy of SE in diagnosing significant CAD in these patients.

Limitations

Our meta-analysis has obvious limitations. The data obtained do not allow conclusions to be drawn in relation to the severity of CAD stenosis. We could not include in our analysis data on CMR and PET techniques, because of the limited number of available publications, which would undermine the relevance of the statistical analysis. Due to the limited number of published articles on the use of speckle tracking echocardiography for the assessment of myocardial deformation in these patients, this analysis could not be performed. Also, due to

the limited number of available studies on the investigation of hypertensive patients based on the antihypertensive therapy and duration of hypertension, these data were not included in the analysis. Moderate heterogeneity across some studies was evident in various risk of bias and analyses; therefore the random-effects method was applied to lessen the impact of this heterogeneity.

Conclusions

This meta-analysis revealed the powerful diagnostic value of stress echocardiography, irrespective of the used modality, in detecting significant coronary artery disease in patients with systemic hypertension. Future analysis of other functional tests and advanced echocardiographic modalities could provide more specific indications for the optimum patient stratification.

Funding

No external funding.

Ethical approval

Not applicable.

Conflict of interest

The authors declare no conflict of interest.

References

- Roth GA, Mensah GA, Johnson CO, et al. Global burden of cardiovascular diseases and risk factors, 1990–2019: update from the GBD 2019 study. *J Am Coll Cardiol* 2020; 76: 2982-3021.
- Aursulesei Onofrei V, Ceasovschi A, Anghel RC, et al. Subendocardial viability ratio predictive value for cardiovascular risk in hypertensive patients. *Medicina (Kaunas)* 2022; 59: 24.
- Henein M, Mörner S, Lindmark K, Lindqvist P. Impaired left ventricular systolic function reserve limits cardiac output and exercise capacity in HFpEF patients due to systemic hypertension. *Int J Cardiol* 2013; 168: 1088-93.
- Kavousi M, Leening MJ, Nanchen D, et al. Comparison of application of the ACC/AHA guidelines, Adult Treatment Panel III guidelines, and European Society of Cardiology guidelines for cardiovascular disease prevention in a European cohort. *JAMA* 2014; 311: 1416-23.
- Del Pinto R, Giua C, Keber E, Grippa E, Tilotta M, Ferri C. Impact of 2021 ESC Guidelines for Cardiovascular Disease Prevention on Hypertensive Patients Risk: Secondary Analysis of Save Your Heart Study. *High Blood Press Cardiovasc Prev* 2023; 30: 167-73.
- Bytyçi I, Morina D, Bytyqi S, Bajraktari G, Henein MY. Percutaneous coronary intervention is not superior to optimal medical therapy in chronic coronary syndrome: a meta-analysis. *J Clin Med* 2023; 12: 1395.
- Cortigiani L, Bigi R, Landi P, Bovenzi F, Picano E, Sicari R. Prognostic implication of stress echocardiography in 6214 hypertensive and 5328 normotensive patients. *Eur Heart J* 2011; 32: 1509-18.
- Chin D, Battistoni A, Tocci G, Passerini J, Parati G, Volpe M. Non-invasive diagnostic testing for coronary artery disease in the hypertensive patient: potential advantages of a risk estimation-based algorithm. *Am J Hypertens* 2012; 25: 1226-35.
- Iwata K, Ogasawara K. Assessment of the efficiency of non-invasive diagnostic imaging modalities for detecting myocardial ischemia in patients suspected of having stable angina. *Healthcare (Basel)* 2022; 11: 23.
- Ballo H, Doghman F, Hartikainen J, Saraste A, Knuuti J. Speckle-tracking echocardiography for predicting improvement of myocardial contractile function after revascularization: a meta-analysis of prospective trials. *Int J Cardiovasc Imaging* 2023; 39: 541-53.
- Bytyçi I, Alves L, Alves O, Lopes C, Bajraktari G, Henein MY. Left ventricular myocardial and cavity velocity disturbances are powerful predictors of significant coronary artery stenosis. *J Clin Med* 2022; 11: 6185.
- Bhat A, Desai A, Amsterdam EA. Usefulness of high functional capacity in patients with exercise-induced ST-depression to predict a negative result on exercise echocardiography and low prognostic risk. *Am J Cardiol* 2008; 101: 1541-3.
- Cury RC, Leipsic J, Abbara S, et al. CAD-RADS™ 2.0 - 2022 Coronary Artery Disease-Reporting and Data System: An Expert Consensus Document of the Society of Cardiovascular Computed Tomography (SCCT), the American College of Cardiology (ACC), the American College of Radiology (ACR), and the North America Society of Cardiovascular Imaging (NASCI). *J Cardiovasc Comput Tomogr* 2022; 16: 536-57.
- Winther S, Schmidt SE, Rasmussen LD, et al. Validation of the European Society of Cardiology pre-test probability model for obstructive coronary artery disease. *Eur Heart J* 2021; 42: 1401-11.
- Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; 372: n71.
- Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.handbook.cochrane.org.
- Zeng X, Zhang Y, Kwong JS, et al. The methodological quality assessment tools for preclinical and clinical studies, systematic review and meta-analysis, and clinical practice guideline: a systematic review. *J Evid Based Med* 2015; 8: 2-10.
- Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Med Res Methodol*. 2005; 5: 13.
- Rutter CM, Gatsonis CA. A hierarchical regression approach to meta-analysis of diagnostic test accuracy evaluations. *Stat Med* 2001; 20: 2865-84.
- Morton SC, Adams JL, Suttrop MJ, Shekelle PG. *Meta-regression Approaches: What, Why, When, and How?*. Rockville (MD): Agency for Healthcare Research and Quality (US); 2004. Report No.: 04-0033.
- Wroblewski EM, Pearl FJ, Hammer WJ, Bove AA. False positive stress tests due to undetected left ventricular hypertrophy. *Am J Epidemiol* 1982; 115: 412-7.
- Picano E, Lucarini AR, Lattanzi F, et al. Dipyridamole echocardiography in essential hypertensive patients with chest pain. *Hypertension* 1988; 12: 238-43.
- Prisant LM, von Dohlen TW, Houghton JL, Carr AA, Frank MJ. A negative thallium (\pm dipyridamole) stress test excludes significant obstructive epicardial coronary artery disease in hypertensive patients. *Am J Hypertens* 1992; 5: 71-5.

24. Massie BM, Szlachcic Y, Tubau JF, O'Kelly BF, Ammon S, Chin W. Scintigraphic and electrocardiographic evidence of silent coronary artery disease in asymptomatic hypertension: a case-control study. *J Am Coll Cardiol* 1993; 22: 1598-606.
25. Tsuda M, Hatano K, Hayashi H, Yokota M, Hirai M, Saito H. Diagnostic value of postexercise systolic blood pressure response for detecting coronary artery disease in patients with or without hypertension. *Am Heart J* 1993; 125: 718-25.
26. Senior R, Basu S, Handler C, Raftery EB, Lahiri A. Diagnostic accuracy of dobutamine stress echocardiography for detection of coronary heart disease in hypertensive patients. *Eur Heart J* 1996; 17: 289-95.
27. Schillaci O, Moroni C, Scopinaro F, et al. Technetium-99m sestamibi myocardial tomography based on dipyridamole echocardiography testing in hypertensive patients with chest pain. *Eur J Nucl Med* 1997; 24: 774-8.
28. Astarita C, Nicolai E, Liguori E, Gambardella S, Rumolo S, Maresca FS. Dipyridamole-echocardiography and thallium exercise myocardial scintigraphy in the diagnosis of obstructive coronary or microvascular disease in hypertensive patients with left ventricular hypertrophy and angina. *G Ital Cardiol* 1998; 28: 996-1004.
29. Elhendy A, Geleijnse L, van Domburg RT, et al. Comparison of dobutamine stress echocardiography and technetium-99m sestamibi single-photon emission tomography for diagnosis of coronary artery disease in hypertensive patients with and without left ventricular hypertrophy. *Eur J Nucl Med* 1998; 25: 69-78.
30. Ho YL, Wu CC, Lin LC, et al. Assessment of the coronary artery disease and systolic dysfunction in hypertensive patients with the dobutamine-atropine stress echocardiography: effect of the left ventricular hypertrophy. *Cardiology* 1998; 89: 52-8.
31. Fragasso G, Lu C, Dabrowski P, Pagnotta P, Sheiban I, Chierchia SL. Comparison of stress/rest myocardial perfusion tomography, dipyridamole and dobutamine stress echocardiography for the detection of coronary disease in hypertensive patients with chest pain and positive exercise test. *J Am Coll Cardiol* 1999; 34: 441-7.
32. Arrif B, Thom SA, Foale RA, et al. Stress echocardiography for the diagnosis of ischemia in hypertensives. *J Hum Hypertens* 2000; 14: 399-401.
33. Maltagliati A, Berti M, Muratori M, et al. Exercise echocardiography versus exercise electrocardiography in the diagnosis of coronary artery disease in hypertension. *Am J Hypertens* 2000; 13: 796-801.
34. Pasiński T, Szwed H, Malczewska B, et al. Advantages of exercise echocardiography in comparison to dobutamine echocardiography in the diagnosis of coronary artery disease in hypertensive subjects. *J Hum Hypertens* 2001; 15: 805-9.
35. Elhendy A, van Domburg RT, Sozzi FB, Poldermans D, Bax JJ, Roelandt JR. Impact of hypertension on the accuracy of exercise stress myocardial perfusion imaging for the diagnosis of coronary artery disease. *Heart* 2001; 85: 655-61.
36. Astarita C, Pálincás A, Nicolai E, Maresca FS, Varga A, Picano E. Dipyridamole-atropine stress echocardiography versus exercise SPECT scintigraphy for detection of coronary artery disease in hypertensives with positive exercise test. *J Hypertens* 2001; 19: 495-502.
37. Cortigiani L, Bigi R, Rigo F, et al. Diagnostic value of exercise electrocardiography and dipyridamole stress echocardiography in hypertensive and normotensive chest pain patients with right bundle branch block. *J Hypertens* 2003; 21: 2189-94.
38. Milosavljevic J, Ostojic M, Marinkovic J. Dipyridamole-dobutamine stress echocardiography for the detection of myocardial ischemia in patients with hypertension. *Herz* 2005; 30: 215-22.
39. Nishida C, Okajima K, Kudo T, Yamamoto T, Hattori R, Nishimura Y. The relationship between coronary artery calcification detected by non-gated multi-detector CT in patients with suspected ischemic heart disease and myocardial ischemia detected by thallium exercise stress testing. *Ann Nucl Med* 2005; 19: 647-53.
40. Michaelides AP, Tousoulis D, Fourlas CA, et al. Hypertensive patients with false-positive thallium-201 scintigraphic results in the infero-posterior wall are in high risk for coronary artery disease development. *Int J Cardiol* 2007; 117: 178-83.
41. Aggeli C, Christoforatu E, Giannopoulos G, et al. The diagnostic value of adenosine stress-contrast echocardiography for diagnosis of coronary artery disease in hypertensive patients: comparison to Tl-201 single-photon emission computed tomography. *Am J Hypertens* 2007; 20: 533-8.
42. Michaelides AP, Liakos CI, Raftopoulos LG, et al. Electrocardiographic criteria for detecting coronary artery disease in hypertensive patients with ST-segment changes during exercise testing. *J Electrocardiol* 2009; 42: 405-9.
43. Lu C, Lu F, Fragasso G, et al. Comparison of exercise electrocardiography, technetium-99m sestamibi single photon emission computed tomography, and dobutamine and dipyridamole echocardiography for detection of coronary artery disease in hypertensive women. *Am J Cardiol* 2010; 105: 1254-60.
44. Cortigiani L, Rigo F, Galderisi M, et al. Diagnostic and prognostic value of Doppler echocardiographic coronary flow reserve in the left anterior descending artery in hypertensive and normotensive patients. *Heart* 2011; 97: 1758-65.
45. Zimarino M, Montebello E, Radico F, et al. ST segment/heart rate hysteresis improves the diagnostic accuracy of ECG stress test for coronary artery disease in patients with left ventricular hypertrophy. *Eur J Prev Cardiol* 2016; 23: 1632-9.
46. Luu JM, Wei J, Shufelt CL, et al. Clinical practice variations in the management of ischemia with no obstructive coronary artery disease. *J Am Heart Assoc* 2022; 11: e022573.
47. Fowler-Brown A, Pignone M, Pletcher M, Tice JA, Sutton SF, Lohr KN; U.S. Preventive Services Task Force. Exercise tolerance testing to screen for coronary heart disease: a systematic review for the technical support for the U.S. Preventive Services Task Force. *Ann Intern Med* 2004; 140: W9-24.
48. Lal K, Singh N, Kumar A, et al. Association of ischemic electrocardiographic changes in high-altitude areas with coronary angiography. *Med J Armed Forces India* 2021; 77: 403-7.
49. Patel MR, Peterson ED, Dai D, et al. Low diagnostic yield of elective coronary angiography. *N Engl J Med* 2010; 362: 886-95.
50. Gyllenhammar T, Carlsson M, Jögi J, Arheden H, Engblom H. Myocardial perfusion by CMR coronary sinus flow shows sex differences and lowered perfusion at stress in patients with suspected microvascular angina. *Clin Physiol Funct Imaging* 2022; 42: 208-19.
51. Koh AS, Blankstein R. Selecting the best noninvasive imaging test to guide treatment after an inconclusive exercise test. *Curr Treat Options Cardiovasc Med* 2012; 14: 8-23.

52. Gulati M, Levy PD, Mukherjee D, et al. 2021 AHA/ACC/AASE/CHEST/SAEM/SCCT/SCMR Guideline for the Evaluation and Diagnosis of Chest Pain: Executive Summary: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation* 2021; 30: e368-454.
53. Barbarie RF, Dib E, Ahmad M. Stress echocardiography using real-time three-dimensional imaging. *Echocardiography* 2018; 35: 1196-203.
54. Cosyns B, Van Camp G, Droogmans S, Weytjens C, Schoors D, Lancellotti P. Analysis of regional wall motion during contrast-enhanced dobutamine stress echocardiography: effect of contrast imaging settings. *Eur J Echocardiogr* 2009; 10: 956-60.