

# Analysis of the current status of CT diagnosis of sarcopenia

---

## Keywords

computed tomography, sarcopenia, artificial intelligence, skeletal muscle index

---

## Abstract

Sarcopenia is a group of clinical syndromes characterized by decreased skeletal muscle mass and strength. The incidence of sarcopenia in the elderly population remains high, which can lead to many adverse events such as falls, fractures, weakness, and also increases the risk of disability and death, so clear diagnosis and early intervention are essential. Computed tomography (CT) has significant advantages over previous diagnostic techniques such as dual-energy X-ray absorptiometry (DXA), bioelectrical impedance analysis (BIA), and body composition analysis, and has been increasingly used to diagnose sarcopenia. CT images use artificial intelligence (AI) software to automatically segment muscle groups and calculate body composition parameters, which is convenient and fast for diagnosing sarcopenia. In order to increase the effectiveness of sarcopenia clinical diagnosis, this study aimed to evaluate the benefits of CT compared to other diagnostic techniques and the value of skeletal muscle parameters at various spinal levels on CT images.

Preprint

## Analysis of the current status of CT diagnosis of sarcopenia

**Abstract:** Sarcopenia is a clinical syndrome characterized by the reduction of skeletal muscle mass and strength, leading to adverse events such as falls, fractures, frailty, disability, and increased mortality. Compared to previous diagnostic techniques such as Dual-Energy X-ray Absorptiometry (DXA), Bioelectrical Impedance Analysis (BIA), and body composition analysis, Computed Tomography (CT) offers significant advantages. Opportunistic CT imaging, enhanced by artificial intelligence (AI) software, provides a superior diagnostic tool for sarcopenia. AI software can automatically segment muscle groups on opportunistic CT images from different populations, enabling the efficient calculation of body composition parameters and more accurate and rapid diagnosis of sarcopenia. Early intervention may significantly reduce adverse clinical outcomes associated with sarcopenia. This study aims to evaluate the advantages of using CT images compared to traditional diagnostic techniques and to assess the value of skeletal muscle parameters at different spinal levels on opportunistic CT images for diagnosing sarcopenia.

**Keywords:** sarcopenia, computed tomography, artificial intelligence, skeletal muscle index

**Abbreviations:** DXA, Dual-Energy X-ray Absorptiometry; BIA, Bioelectrical Impedance Analysis; CT, Computed Tomography; AI, artificial intelligence; HF, heart failure; FS, frailty syndrome; IWGS, International Working Group on Sarcopenia; SMM, skeletal muscle mass; AWGS, Asian Working Group for Sarcopenia; CSA, cross-sectional area; SMA, skeletal muscle area; SMI, skeletal muscle index; L4, the fourth lumbar vertebra; L3,

the third lumbar vertebra; T10, the tenth thoracic vertebra; L5, the fifth lumbar vertebra; L1, the first lumbar vertebra; T4, the fourth thoracic vertebra; IPF, idiopathic pulmonary fibrosis; T6, the sixth thoracic vertebra; T12, the twelfth thoracic vertebra; C3, the third cervical vertebra; HNC, head and neck cancer; CNN, convolutional neural network; VAT, visceral adipose tissue; SAT, subcutaneous adipose tissue.

Irwin Rosenberg, an American researcher, originally proposed the concept of sarcopenia as a type of geriatric condition in 1989<sup>1</sup>, mainly characterized by decreased muscle mass, weakened muscle strength, and a decrease in individual activity ability. According to a study, sarcopenia affects older adults between the ages of 60 and 80 with a frequency of 11.1% and older adults above the age of 80 at a prevalence of 33.1%<sup>2</sup>. In addition, sarcopenia is closely related to heart failure (HF), frailty, etc. In patients with HF, myocardial contractility decreases, and skeletal muscle blood perfusion is insufficient, resulting in reduced skeletal muscle mass and function. Sarcopenia with frailty syndrome (FS) can also lead to a decline in physical function and quality of life in patients with a variety of diseases such as diabetes and HF<sup>3-5</sup>, as well as an increase in hospitalization rates, length of stay and hospitalization costs, thereby increasing the public health burden—one of the biggest medical challenges society will face in the future.

Studies have shown that sarcopenia is not only associated with aging but also occurs at an earlier age in chronic patients<sup>6</sup>. International Working Group on Sarcopenia (IWGS) 2011 considers sarcopenia to be like most kinds of geriatric syndromes, rarely recognized

by doctors<sup>7</sup>. Sarcopenia must be accurately diagnosed and treated as soon as possible because it can be a significant predictor of poor clinical outcomes in the elderly and patients with many other diseases<sup>8</sup>. We hope that opportunistic CT screening can provide muscle mass information, improve clinicians' understanding of sarcopenia, and provide a basis for clinical screening, identification, diagnosis, treatment, and intervention. Because of the complex diagnostic tools and inconsistent diagnostic criteria for sarcopenia<sup>9</sup>, there are great differences in the clinical diagnosis of sarcopenia<sup>10,11</sup>, and the prevalence rate has a large fluctuation range, so the research on universal diagnostic tools and diagnostic criteria needs to be further carried out.

Many patients with sarcopenia do not exhibit obvious symptoms, but the reduction in muscle mass is significant on imaging. In real life, being used for diagnosing thoracoabdominal diseases, CT is also increasingly utilized for routine health check-ups. Recently, CT is becoming more common in measuring muscles<sup>12-14</sup>. CT measurement of muscle size is a reliable and valid technique that can be used in numerous studies to assess the presence of sarcopenia in healthy adults<sup>15</sup>, and this CT body composition analysis has evolved into an objective measure of physical health in patients with comorbidities. The chest CT analysis we describe for assessing the presence or absence of sarcopenia has the distinct advantage of leveraging the patient's existing opportunistic CT images, combined with an AI-automated analysis process, to improve diagnostic efficiency without increased radiation exposure and additional test costs. The CT imaging technique clinically allows

straightforward body composition analysis and sarcopenia quantification from a single axial image acquired at the different vertebrae, with the advantages and utility of CT imaging technology far greater than that of DXA and BIA<sup>16</sup>.

Many international working groups on sarcopenia have noted that an evaluation of muscle mass and function should be part of the diagnosis. However, the cut-off point of muscle mass is still used as the core diagnostic index<sup>17</sup>. To increase the application value of opportunistic CT screening and to provide more evidence for clinical sarcopenia diagnosis, this paper compares the current CT diagnosis of sarcopenia with other diagnostic techniques.

### **1. Comparison between existing diagnostic methods of muscle volume and CT**

Skeletal muscle mass (SMM) is closely related to skeletal muscle strength and function. Total body skeletal muscle mass, appendicular skeletal muscle mass, or skeletal muscle cross-sectional area of a specific muscle group or body part can reflect SMM. Muscle loss can be divided into primary and secondary causes, with primary sarcopenia referring to a reduction of muscle mediated by aging. Secondary sarcopenia focuses on loss of muscle mass, rather than emphasizing muscle function. With the deepening of the understanding of the pathogenesis of sarcopenia, the statistics of the incidence of this disease are becoming more and more common. Studies have shown that the overall prevalence of sarcopenia in Chinese elderly men is 18%, the total prevalence of elderly men living in the community is 12.9%, and the total prevalence of hospitalized elderly men

is 29.7%. The overall prevalence of sarcopenia in elderly women was 16.4%, the prevalence of community-living elderly women was 11.2%, and the prevalence of hospitalized elderly women was 23.0%<sup>18</sup>.

In the diagnosis of sarcopenia, there are many screening methods used, some for clinical purposes, and some for scientific research. In practical application, appropriate diagnostic techniques can be selected according to the requirements and advantages and disadvantages. Based on the attenuation characteristics of the X-ray beam through different tissues such as muscle and fat, CT examination can accurately distinguish muscle from other body tissues and provide information about muscle mass. Utilizing opportunistic CT images from patients or individuals undergoing health examinations incurs no additional examination costs. These opportunistic CT images may be an overlooked resource; if effectively used in sarcopenia assessment, they could greatly advance clinical diagnostic research for sarcopenia. Although BIA is easy to operate and is useful in primary care, its low accuracy limits its diagnostic application in clinical practice. Due to its outstanding advantages, CT image has gradually replaced traditional diagnostic techniques such as DXA, BIA, and body composition analysis to evaluate muscle mass, and it has become the first choice of more clinical workers.

### 1.1 DXA

DXA assesses fat, bone, and muscle mass by analyzing different attenuation characteristics of X-rays in soft tissue, bone, and muscle. The diagnostic accuracy of

sarcopenia is high, the examination time is short, the cost is low, the radiation is low, and the patient compliance is high, so it has been widely used in early scientific research related to the diagnosis of sarcopenia. DXA diagnosis of sarcopenia typically follows the standards set by the Asian Working Group for Sarcopenia (AWGS)<sup>19</sup>. The reference values of diagnostic cut-off points for men and women were 7.0 kg/m<sup>2</sup> and 5.4 kg/m<sup>2</sup> respectively, which were more consistent with the body composition, genetic background, and ethnic characteristics of the Asian population. However, there are some drawbacks to the practical application of DXA: the equipment is expensive, non-portable, non-mobile, and difficult to obtain in primary medical institutions, which limits its application in large-scale epidemiological studies. The utilization rate of DXA examination equipment in various medical institutions is relatively low, with the number of DXA scanners per million population ranging from 24 in South Korea to less than 1 in Sri Lanka and Vietnam<sup>20</sup>. In addition, because its measurement accuracy is easily affected by the beam hardening effect and soft tissue hydration, it is inappropriate for obese patients or patients with excessive water volume<sup>21</sup>. Therefore, DXA plays a limited role in clinical practice.

## 1.2 BIA

BIA is a painless and non-invasive technology that can measure body composition such as fat, muscle, bone, and water according to the electrical conductivity of the whole body, the advantages are: easy to operate, safe and non-invasive, low cost, fast execution, convenient bedside application. The BIA diagnostic criteria for sarcopenia updated by

AWGS2 were:  $<7.0 \text{ kg/m}^2$  for males and  $<5.7 \text{ kg/m}^2$  for females. However, since human impedance is closely related to water content, BIA is highly susceptible to changes in total water content and body fluids when conducting conductivity analysis, such as environmental temperature, diet, water, diarrhea, and exercise<sup>22</sup>. The accuracy of the BIA is suboptimal, and the validity of its results cannot be unequivocally guaranteed. Moreover, the limited availability of equipment hampers its widespread implementation in large-scale clinical practice. BIA is only employed as a screening tool for sarcopenia when superior alternatives are not readily accessible<sup>23</sup>.

### 1.3 CT

CT examination is routinely used clinically<sup>24</sup>. DXA and BIA are far less accessible as the test device than conventional CT. For optimistic CT images of patients with other diseases, if the information on muscle metabolism and muscle mass can be regularly extracted from the image, opportunistic sarcopenia screening will help in the early detection and follow-up of muscle metabolic diseases. CT, as a cross-sectional imaging technique, can be used to assess many acute and chronic diseases associated with aging, including fractures, frailty, cancer, and cardiometabolic syndrome<sup>25</sup>, and has been associated with many important clinical outcomes, including death<sup>26,27</sup>. CT image files of patients were extracted from different vertebral body levels, such as skeletal muscles of the third lumbar spine, including erector, transversus abdominis muscle, psoas major, internal oblique muscle, and external oblique muscle. The threshold of muscle tissue unit ranges



from -29 to 150 HU. The total skeletal muscle area (cm<sup>2</sup>) can be obtained by semi-automatic mapping using imaging image processing software. Some scholars<sup>28</sup> believe that using the cross-sectional area (cm<sup>2</sup>) of a single skeletal muscle on CT images to predict the skeletal muscle content of the whole body is reproducible and highly correlated with other methods to evaluate skeletal muscle content. CT examination has become the main means to assess muscle mass in patients with many diseases such as cancer<sup>28</sup>, which can display lesions, provide accurate scanning details, and help obtain skeletal muscle parameters such as muscle cross-sectional area (CSA), skeletal muscle area (SMA), skeletal muscle index (SMI), skeletal muscle volume.

CT is the most accurate method to measure muscle mass lately, which is the "gold standard" of morphology<sup>29</sup>. Moreover, the imaging technique can be used for multiple disease stages and follow-ups<sup>30</sup>, so sarcopenia can be evaluated in prospective and retrospective analyses. Analyzing patients' opportunistic CT images to assess muscle mass allows for a more objective quantitative diagnosis, which is easier and more efficient. Deploying opportunistic CT images as a superior examination method can provide new strategies for sarcopenia diagnosis and treatment. A study of 450 trauma patients reported that CT-derived sarcopenia was associated with increased mortality<sup>31</sup>. Another study of 83 patients showed that CT-derived measurements of sarcopenia were predictive of hospital length of stay in older burn patients<sup>32</sup>.

However, the criteria for determining skeletal muscle parameters at the level of CT

images to assess sarcopenia are not uniform at present, and the cut-off point values used at the same level are also different in different studies. The main reason is that the field of view from the scan, the patient's position, and the selection of vertebral segments are different. There is not much difference in the concept and content of sarcopenia. Clinically, when diagnosing sarcopenia based on CT images, a unified measurement posture, a unified analysis method, and a multivariate prediction rule combined with the characteristics of the patients can be used to estimate muscle CSA better<sup>33</sup>. The diagnostic criteria of AWGS are mainly referred to in our country. The following focuses on the development of CT in diagnosing sarcopenia and discusses the diagnostic value of AI-driven opportunistic CT images.

## **2. CT image and diagnosis of sarcopenia**

### **2.1 L3 SMI**

Regarding the CT diagnosis of sarcopenia, the initial research mainly focused on the psoas major. Bokshan et al.<sup>34</sup> diagnosed sarcopenia by dividing the cross-sectional area of the psoas major muscle at the fourth lumbar vertebra (L4) in the abdominal CT images by the square of the height. Later, more studies found that the muscle mass at the third lumbar vertebra (L3) in abdominal CT images was closely related to the SMM of the limbs, and the quantitative measurement of skeletal muscle and parameters related to skeletal muscle space fat at L3 vertebral body level by CT had high repeatability. CT can accurately estimate skeletal muscle volume and whole-body muscle mass by measuring the muscle

cross-sectional area of the L3 layer with three-dimensional imaging technology<sup>35,36</sup>.

Therefore, the most commonly used technique is to evaluate all skeletal muscle on axial sections at the L3 vertebral level<sup>37</sup>. These studies also further validated the reliability of quantitative CT measurements for L3 vertebral plane skeletal muscle analysis. SMI is the muscle mass of limbs after height correction [muscle mass of limbs (kg) / height of 2 (m<sup>2</sup>)]. On CT images, SMI can be represented by the CSA of muscle tissue (cm<sup>2</sup>) / height (m<sup>2</sup>) at different vertebral layers. L3 SMA or L3 SMI measured by CT can represent total body skeletal muscle mass and has become a common method for assessing sarcopenia<sup>38-41</sup>. L3 SMI is also the gold standard for the diagnosis of sarcopenia at home and abroad, which is widely used in the detection of sarcopenia in cancer patients. Currently, the reference value of the L3 SMI tangent point used in most cases is  $\leq 39$  cm<sup>2</sup>/ m<sup>2</sup> for females and  $\leq 55$  cm<sup>2</sup>/ m<sup>2</sup> for males<sup>17</sup>.

## 2.2 L1 SMI

Muscle measurement at the L3 level requires patients to undergo abdominal CT examination, which is not suitable for routine evaluation of patients. The skeletal muscle cut-point values from the tenth thoracic vertebra (T10) to the fifth lumbar vertebra (L5), reported by Derstine et al.<sup>42</sup> extended the assessment of sarcopenia to the chest and pelvic examination so that the imaging technique could be applied more widely in the clinic.

Although not all chest CT scan images provided by patients with thoracic malignancies and other diseases include the L3 level, they can provide skeletal muscle parameters and other

body tissue composition information at the first lumbar vertebra (L1) level. Many scholars have found that the muscle mass information of other vertebral segments on CT images can also be used to evaluate sarcopenia<sup>43</sup>, and patients do not need to undergo non-contrast abdominal CT scanning. Chest CT can depict the conditions of the pectoralis major and latissimus dorsi muscles. Therefore, the analysis of skeletal muscle information at different levels of chest CT images is of great significance for confirming sarcopenia. Studies<sup>44</sup> have shown that the cut-point values of L1 SMI determined by chest CT for the diagnosis of sarcopenia are  $46 \text{ cm}^2/\text{m}^2$  (male) and  $29 \text{ cm}^2/\text{m}^2$  (female). At present, this test is widely used in lung cancer patients<sup>45</sup>. Based on a population in central China, one study defined a sex-specific cutoff for L1 SMI with low skeletal muscle mass as  $36.52 \text{ cm}^2/\text{m}^2$  for men and  $27.29 \text{ cm}^2/\text{m}^2$  for women<sup>46</sup>.

### 2.3 T4 SMI

Pectoral muscle size is a reliable and valid piece of information that can be used in large studies to assess the presence or absence of muscle loss in patients with advanced lung disease<sup>47</sup>. Kim et al.<sup>48</sup> used the CSA of chest muscle measured by CT to conduct a correlation study with SMM measured by BIA analysis, the results showed that SMI at the fourth thoracic vertebra (T4) was moderately correlated with L3 SMI, and L1 SMI was also highly correlated with L3 SMI. Because thoracic muscle is easy to label, and its measurement is easier than that of skeletal muscle at the L3 level, T4 SMI can be used as an indicator to detect sarcopenia in patients with thoracic tumors and other types of

patients, and it is increasingly applied in related studies. Low T4 CSA reflected on chest CT images of patients with idiopathic pulmonary fibrosis (IPF), chest tumors, and chronic obstructive pulmonary disease may be a strong risk factor for all-cause death<sup>49,50</sup>, and early diagnosis and intervention can improve the prognosis of patients. In addition, the thoracic muscles from T4 to the sixth thoracic vertebra (T6) were fairly evenly distributed, CSA at each level was strongly correlated with muscle volume<sup>47</sup>, and moderately to highly correlated with measurements of the quadriceps muscle. It is concluded that T4 SMI, T5 SMI, and T6 SMI can effectively reflect the total body muscle volume and be used to diagnose sarcopenia.

#### 2.4 T12 SMI

Chest skeletal muscle measurements can reasonably represent the area of other skeletal muscles in the body. The muscle mass area at the twelfth thoracic vertebra (T12) on a CT image can easily and accurately diagnose sarcopenia. The T12 level muscle measurement has been used more as a clinical tool. Gao et al.<sup>51</sup> used chest CT to measure the cross-sectional area of muscle tissue at the level of the T12 pedicle and calculated T12 SMI. The cut-off values for the diagnosis of sarcopenia can be used according to the criteria proposed by Nemec et al.<sup>52</sup> of Harvard Medical School: 42.6 cm<sup>2</sup>/ m<sup>2</sup> (male) and 30.6 cm<sup>2</sup>/ m<sup>2</sup> (female). In patients with COPD, chest CT measurements provide the size of the pectoral and erector spinal muscles, which correlate with disease-related parameters such as body mass index, lung function, exercise capacity, and dyspnea<sup>53</sup>. Although abdominal CT

is not widely used in sarcopenia diagnosis with T12 SMI, in some specific cases, such as thoracic aneurysm repair<sup>54</sup>, chest CT is superior to abdominal CT in the evaluation of sarcopenia.

## 2.5 C3 SMI

In addition, as research progresses, multiple studies have shown that the assessment of SMM at the third cervical vertebra (C3) level is simple and reliable<sup>55</sup>, that measurement of SMM at the C3 level in head and neck CT imaging may be a viable, economical, accurate, meaningful, and rapid biomarker for the detection of sarcopenia in patients with head and neck cancer (HNC)<sup>38,56</sup>. Chargini et al.<sup>57</sup> proposed a new criterion for diagnosing sarcopenia: for BMI < 25 kg/m<sup>2</sup> and ≥ 25 kg/m<sup>2</sup>, low SMM in men ≤ 6.8 cm<sup>2</sup>/m<sup>2</sup> and ≤ 8.5 cm<sup>2</sup>/m<sup>2</sup>, respectively, low SMM in women ≤ 5.3 cm<sup>2</sup>/m<sup>2</sup> and 6.4 cm<sup>2</sup>/m<sup>2</sup>, respectively. The muscle mass information based on CT measurement technology is reliable, and it can be used as an important predictor of surgical morbidity and mortality, which will have a wider application prospect in the future.

All in all, the thoracoabdominal muscle reduction measurement is an additional robust clinical tool that can be applied to a variety of disease states, especially those patients undergoing routine CT scans. This can be used to assess sarcopenia to reduce the cost of testing and expand the diagnostic strategy. However, it is undoubtedly a time-consuming and onerous task for clinicians to manually segment muscle groups on CT images and calculate muscle volume or area to evaluate muscle mass. So far, artificial intelligence

image analysis software has been gradually put into use. And machine learning algorithms are better than manual measurements of bone and muscle<sup>58,59</sup>.

### **3. The application advantage of artificial intelligence technology**

Machine learning, part of artificial intelligence, can be used to analyze medical big data. It is independent of individual subjective factors and provides the opportunity to frequently assess some relevant variables to help diagnose disease, identify hidden trends, and develop predictive models<sup>60,61</sup>. The application of artificial intelligence and machine learning technologies has enormous potential benefits and will become increasingly irreplaceable in the future development of medicine.

#### **3.1 Reduce labor costs**

Many methods have been successfully used to automatically segment muscle groups on single-layer CT images recently, among which the methods based on atlas and convolutional neural network (CNN)<sup>58,62-66</sup> show great advantages. The traditional method of manually tracking muscle groups to calculate body composition parameters to diagnose sarcopenia is onerous, expensive, and time-consuming. In the era of precision medicine, AI-based analysis relies on differences in CT attenuation values between muscles and other tissues in single-layer CT sections, the measured skeletal muscle parameters can be used as indicators for the diagnosis of sarcopenia. AI graphic analysis software may realize the automation of labor-intensive tasks, and can greatly reduce the workload of clinicians, shorten the diagnosis time, and improve the diagnosis efficiency<sup>28</sup>. A study suggests that

CT-based AI-powered software facilitates the evaluation of objective clinical benefits and prognostic tumor outcomes<sup>67</sup>.

### 3.2 Improve the accuracy of sarcopenia assessment

Artificial segmentation analysis is prone to be limited by inter-observer variability, fat infiltration, observer time, and other factors<sup>68</sup>. AI can overcome these problems<sup>69</sup>, and improve calibration performance<sup>70</sup>. AI can help divide skeletal muscle in a short amount of time and with the same precision as a trained subject<sup>28</sup>. In addition, we found that AI can be very effective in using CT imaging for automated body composition analysis, providing physicians with meaningful imaging biomarkers and further improving the clinical value of radiological reports to improve patient care<sup>71</sup>. Therefore, the use of CT image-based AI models to diagnose sarcopenia is of great clinical value regardless of clinical indications<sup>72</sup>. AI image analysis can accurately segment muscle and adipose tissues<sup>73,74</sup>, to distinguish the relative proportions of various tissues, and automatically calculate muscle and adipose tissue parameters<sup>75</sup>, including visceral adipose tissue (VAT), subcutaneous adipose tissue (SAT), and SMI, etc. Although the specific diagnostic threshold of sarcopenia used by this method is still uncertain, its accuracy in evaluating sarcopenia is certain.

### 3.3 Standardizing medical procedures

The special mathematical reconstruction algorithm in AI graphics software is used to automatically measure the distribution of skeletal muscle components and adipose tissue in different levels, various regions, and individual muscle groups on CT images with high



precision, to achieve a quantitative assessment of muscle mass. This AI algorithm can also achieve standardization and repeatability of medical procedures for sarcopenia diagnosis. It is helpful to simplify the diagnostic process and improve the evaluation criteria of sarcopenia. In conclusion, AI-derived body composition represents an accessible method that can be performed post-processing with imaging data<sup>76</sup>, providing the basis for rapid diagnosis of sarcopenia and assessment of clinical outcomes.

### 3.4 To help predict clinical outcome

When discussing the association between sarcopenia and postoperative infection in patients with colorectal cancer, Lieffers et al.<sup>77</sup> found that the postoperative infection rate in patients with sarcopenia was as high as 23.7% (P=0.025), indicating that sarcopenia could predict postoperative infection, affect hospitalization rehabilitation nursing, and thus prolong postoperative hospitalization. In addition, many studies have shown that sarcopenia is associated with increased morbidity and mortality in patients with various underlying diseases<sup>78</sup>. Sarcopenia may also be a marker of poor prognosis in patients with different diseases such as cancer<sup>39,79,80</sup>. CT-based screening for sarcopenia has predictive value in a vertebral compression fracture in elderly individuals<sup>81</sup>, the survival rate after lung transplantation<sup>82</sup>, the length of ICU stay in colon cancer patients<sup>83</sup>, and the patients' overall survival time in lung cancer<sup>84,85</sup>. The influence of body composition on various clinical outcomes has been broadly studied, and body composition has thus become an accepted risk factor. As this valuable metabolic information can easily be extracted from CT images,

there is a growing demand for automatic software solutions.

Early diagnosis of sarcopenia with a view to early intervention plays an important role in improving the prognosis of patients. For example, patients with advanced lung disease can improve clinical outcomes through pulmonary rehabilitation<sup>9</sup>. AI-assisted image analysis uses automatic tissue segmentation technology to provide a standardized trend, which can explore the continuous changes in cancer-related sarcopenia<sup>86</sup> so that clinicians can obtain additional information needed to develop a more rational treatment plan to prevent or delay adverse consequences, and ultimately improve the clinical outcome of patients<sup>87</sup>. In addition, preoperative muscle mass assessment can help clinicians predict patients' health status, and assist in better classification and targeted therapy for the best treatment results. The development of this automated CT segmentation assessment of body composition will accelerate the study of body composition and ultimately facilitate the integration of body composition measures into clinical care<sup>88,89</sup>.

### 3.5 Assist in determining the cut point value of CT image diagnosis of sarcopenia

In daily clinical practice, the AI automatic segmentation model can simplify the diagnosis of sarcopenia and provide additional image data when CT images are examined for various clinical indications. Due to the variety of diagnostic methods and the difference in the definition of the thresholds between different populations, we need large data sets with multi-center and large sample sizes. With higher-quality data annotation and large-scale AI models, relying on appropriate algorithms and eliminating heterogeneity through

integrated data processing, we believe appropriate diagnostic criteria can be explored for early diagnosis of sarcopenia and effective intervention and treatment for patients.

Using this concept, a study has solved the variability of multi-source heterogeneous data in patients with COVID-19 pulmonary infection from different subordinate medical institutions of the center and successfully established a standard prediction model for the outcome of pulmonary dispersion function after COVID-19<sup>90</sup>. Therefore, an automatic CT image detection model based on deep learning should be built as soon as possible. Optimizing the AI algorithm, revealing the cutting point value of sarcopenia diagnosis and the change rule after the intervention, is conducive to providing a decision-making basis for clinical diagnosis and early prevention.

Loss of muscle mass is an important predictor of adverse clinical outcomes. Screening for sarcopenia and implementing appropriate interventions in elderly individuals is crucial. This article proposes using opportunistic CT imaging as an advantageous method for clinical application. In routine clinical practice, opportunistic CT imaging holds significant clinical value. AI-driven CT image analysis can rapidly and accurately monitor muscle mass and health status, facilitating better risk stratification for different patient groups. This technology optimizes clinical decision-making and reduces healthcare costs. Consequently, future research should focus on high-quality and rigorous methodologies to establish reliable diagnostic thresholds for sarcopenia and automate the diagnostic process. This will enhance the efficiency of sarcopenia diagnosis, leading to early intervention and

delayed disease progression.

## **Statements & Declarations**

**Ethics approval and consent to participate:** Does not include patient-specific clinical information or involve ethical considerations. Related issues such as informed consent statements and data privacy are not applicable.

**Competing interests:** The authors have no relevant financial or non-financial interests to disclose.

**Funding:** The authors declare that no funds, grants, or other support were received during the preparation of this manuscript.

**Authors' contributions:** All authors read and revised the manuscript.

## References

1. Truswell AS. Standardization of nomenclature of body composition in weight loss. *Am J Clin Nutr.* Mar 1998;67(3):492-3. doi:10.1093/ajcn/67.3.492
2. Cao M, Lian J, Lin X, Liu J, Chen C, Xu S, Ma S, Wang F, Zhang N, Qi X, Xu G, Peng N. Prevalence of sarcopenia under different diagnostic criteria and the changes in muscle mass, muscle strength, and physical function with age in Chinese old adults. *BMC Geriatr.* Nov 22 2022;22(1):889. doi:10.1186/s12877-022-03601-7
3. Chew STH, Tey SL, Yalawar M, Liu Z, Baggs G, How CH, Cheong M, Chow WL, Low YL, Huynh DTT, Tan NC. Prevalence and associated factors of sarcopenia in community-dwelling older adults at risk of malnutrition. *BMC Geriatr.* Dec 24 2022;22(1):997. doi:10.1186/s12877-022-03704-1
4. Keevil VL, Romero-Ortuno R. Ageing well: a review of sarcopenia and frailty. *Proc Nutr Soc.* Nov 2015;74(4):337-47. doi:10.1017/s0029665115002037
5. Uchmanowicz I, Rosano G, Piepoli M, Vellone E, Czapla M, Lisiak M, Diakowska D, Prokopowicz A, Aleksandrowicz K, Nowak B, Wleklik M, Faulkner K. The concurrent impact of mild cognitive impairment and frailty syndrome in heart failure. *Archives of Medical Science.* 2023;doi:10.5114/aoms/162369
6. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, Martin FC, Michel JP, Rolland Y, Schneider SM, Topinkova E, Vandewoude M, Zamboni M, European Working Group on Sarcopenia in Older P. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing.* Jul 2010;39(4):412-23. doi:10.1093/ageing/afq034

7. Fielding RA, Vellas B, Evans WJ, Bhasin S, Morley JE, Newman AB, Abellan vKG, Andrieu S, Bauer J, Breuille D. Sarcopenia: An Undiagnosed Condition in Older Adults. Current Consensus Definition: Prevalence, Etiology, and Consequences. International Working Group on Sarcopenia. *Journal of the American Medical Directors Association*. 2011;12(4):249-256.
8. Vitale JA, Messina C, Albano D, Fascio E, Galbusera F, Corbetta S, Sconfienza LM, Banfi G. Appendicular Muscle Mass, Thigh Intermuscular Fat Infiltration, and Risk of Fall in Postmenopausal Osteoporotic Elder Women. *Gerontology*. 2021;67(4):415-424. doi:10.1159/000513597
9. Cruz-Jentoft AJ, Landi F, Schneider SM, Zuniga C, Arai H, Boirie Y, Chen LK, Fielding RA, Martin FC, Michel JP, Sieber C, Stout JR, Studenski SA, Vellas B, Woo J, Zamboni M, Cederholm T. Prevalence of and interventions for sarcopenia in ageing adults: a systematic review. Report of the International Sarcopenia Initiative (EWGSOP and IWGS). Review. *Age and Ageing*. Nov 2014;43(6):748-759. doi:10.1093/ageing/afu115
10. Reiss J, Iglseider B, Alzner R, Mayr-Pirker B, Pirich C, Kassmann H, Kreutzer M, Dovjak P, Reiter R. Consequences of applying the new EWGSOP2 guideline instead of the former EWGSOP guideline for sarcopenia case finding in older patients. *Age Ageing*. Sep 1 2019;48(5):719-724. doi:10.1093/ageing/afz035
11. Bianchi L, Maietti E, Abete P, Bellelli G, Bo M, Cherubini A, Corica F, Di Bari M, Maggio M, Martone AM, Rizzo MR, Rossi AP, Volpato S, Landi F, Investigators GG. Comparing EWGSOP2 and FNIH Sarcopenia Definitions: Agreement and 3-Year Survival Prognostic Value in Older Hospitalized Adults: The GLISTEN Study. *J Gerontol A Biol Sci Med Sci*. Jun 18 2020;75(7):1331-1337. doi:10.1093/gerona/glz249
12. Jing X, Tan L, Fu H, Yang L, Yang M. Associations of ADL Disability With Trunk Muscle Mass and Muscle Quality Indicators Measured by Opportunistic Chest Computed Tomography Imaging Among Older Inpatients. *Front Med (Lausanne)*. 2021;8:743698. doi:10.3389/fmed.2021.743698
13. Lopez PD, Nepal P, Akinlonu A, Nekkalapudi D, Kim K, Cativo EH, Visco F, Mushiyevev S, Pekler G. Low Skeletal Muscle Mass Independently Predicts Mortality in Patients with Chronic Heart Failure after an Acute Hospitalization. *Cardiology*. 2019;142(1):28-36. doi:10.1159/000496460
14. Touban BM, Pavlesen S, Smoak JB, Sayegh MJ, Wang J, Zhao J, Anders MJ. Decreased Lean Psoas Cross-Sectional Area Is Associated With Increased 1-Year All-Cause Mortality in Male Elderly Orthopaedic Trauma Patients. *J Orthop Trauma*. Jan 2019;33(1):e1-e7. doi:10.1097/bot.0000000000001331
15. Tan L, Ji G, Bao T, Fu H, Yang L, Yang M. Diagnosing sarcopenia and myosteatorsis based on chest computed tomography images in healthy Chinese adults. *Insights Imaging*. Nov 6 2021;12(1):163. doi:10.1186/s13244-021-01106-2
16. Beetz NL, Geisel D, Maier C, Auer TA, Shnayien S, Malinka T, Neumann CCM, Pelzer U, Fehrenbach U. Influence of Baseline CT Body Composition Parameters on Survival in Patients with Pancreatic Adenocarcinoma. *J Clin Med*. Apr 22 2022;11(9)doi:10.3390/jcm11092356
17. Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyere O, Cederholm T, Cooper C, Landi F, Rolland Y, Sayer AA, Schneider SM, Sieber CC, Topinkova E, Vandewoude M, Visser M, Zamboni M, Ewgsop, Extended Grp E. Sarcopenia: revised European consensus on definition and diagnosis (vol 48, pg 16, 2019). Correction. *Age and Ageing*. Jul 2019;48(4):1. 601. doi:10.1093/ageing/afz046
18. Chen Z, Li WY, Ho M, Chau PH. The Prevalence of Sarcopenia in Chinese Older Adults: Meta-Analysis and Meta-Regression. *Nutrients*. Apr 24 2021;13(5)doi:10.3390/nu13051441

19. Chen LK, Liu LK, Woo J, Assantachai P, Auyeung TW, Bahyah KS, Chou MY, Chen LY, Hsu PS, Krairit O, Lee JSW, Lee WJ, Lee Y, Liang CK, Limpawattana P, Lin CS, Peng LN, Satake S, Suzuki T, Won CW, Wu CH, Wu SN, Zhang TM, Zeng P, Akishita M, Arai H. Sarcopenia in Asia: Consensus Report of the Asian Working Group for Sarcopenia. Review. *Journal of the American Medical Directors Association*. Feb 2014;15(2):95-101. doi:10.1016/j.jamda.2013.11.025
20. Wang W, Richards G, Rea S. Hybrid data mining ensemble for predicting osteoporosis risk. *Conf Proc IEEE Eng Med Biol Soc*. 2005;2006:886-9. doi:10.1109/iembs.2005.1616557
21. Jimenez-Gutierrez GE, Martinez-Gomez LE, Martinez-Armenta C, Pineda C, Martinez-Nava GA, Lopez-Reyes A. Molecular Mechanisms of Inflammation in Sarcopenia: Diagnosis and Therapeutic Update. *Cells*. Aug 1 2022;11(15)doi:10.3390/cells11152359
22. Herda AA, Cleary CJ. Agreement between multifrequency BIA and DXA for assessing segmental appendicular skeletal muscle mass in older adults. *Aging Clin Exp Res*. Nov 2022;34(11):2789-2795. doi:10.1007/s40520-021-02000-z
23. Cheng KY, Chow SK, Hung VW, Wong CH, Wong RM, Tsang CS, Kwok T, Cheung WH. Diagnosis of sarcopenia by evaluating skeletal muscle mass by adjusted bioimpedance analysis validated with dual-energy X-ray absorptiometry. *J Cachexia Sarcopenia Muscle*. Dec 2021;12(6):2163-2173. doi:10.1002/jcsm.12825
24. Vasilevska Nikodinovska V, Ivanoski S. Sarcopenia, More Than Just Muscle Atrophy: Imaging Methods for the Assessment of Muscle Quantity and Quality. *Rofo*. May 9 2023;Sarkopenie, mehr als nur Muskelschwund: Bildgebende Verfahren zur Beurteilung von Muskelquantitat und -qualitat. doi:10.1055/a-2057-0205
25. Radwan RW, Tang AM, Beasley WD. Computed tomography as a first-line investigation for elderly patients admitted to a surgical assessment unit. *Ann R Coll Surg Engl*. Apr 2018;100(4):285-289. doi:10.1308/rcsann.2017.0231
26. Lenchik L, Lenoir KM, Tan J, Boutin RD, Callahan KE, Kritchevsky SB, Wells BJ. Opportunistic Measurement of Skeletal Muscle Size and Muscle Attenuation on Computed Tomography Predicts 1-Year Mortality in Medicare Patients. *J Gerontol A Biol Sci Med Sci*. Jun 18 2019;74(7):1063-1069. doi:10.1093/gerona/gly183
27. Boutin RD, Lenchik L. Value-Added Opportunistic CT: Insights Into Osteoporosis and Sarcopenia. *AJR Am J Roentgenol*. Sep 2020;215(3):582-594. doi:10.2214/AJR.20.22874
28. Chianca V, Albano D, Messina C, Gitto S, Ruffo G, Guarino S, Del Grande F, Sconfienza LM. Sarcopenia: imaging assessment and clinical application. *Abdom Radiol (NY)*. Sep 2022;47(9):3205-3216. doi:10.1007/s00261-021-03294-3
29. Albano D, Messina C, Vitale J, Sconfienza LM. Imaging of sarcopenia: old evidence and new insights. *Eur Radiol*. Apr 2020;30(4):2199-2208. doi:10.1007/s00330-019-06573-2
30. Boutin RD, Houston DK, Chaudhari AS, Willis MH, Fausett CL, Lenchik L. Imaging of Sarcopenia. *Radiol Clin North Am*. Jul 2022;60(4):575-582. doi:10.1016/j.rcl.2022.03.001
31. Kaplan SJ, Pham TN, Arbabi S, Gross JA, Damodarasamy M, Bentov I, Taitsman LA, Mitchell SH, Reed MJ. Association of Radiologic Indicators of Frailty With 1-Year Mortality in Older Trauma Patients: Opportunistic Screening for Sarcopenia and Osteopenia. *JAMA Surg*. Feb 15 2017;152(2):e164604. doi:10.1001/jamasurg.2016.4604
32. Romanowski KS, Fuanga P, Siddiqui S, Lenchik L, Palmieri TL, Boutin RD. Computed Tomography

- Measurements of Sarcopenia Predict Length of Stay in Older Burn Patients. *J Burn Care Res.* Feb 3 2021;42(1):3-8. doi:10.1093/jbcr/iraa149
33. van Heusden HC, Swartz JE, Chargi N, de Jong PA, van Baal M, Wegner I, de Bree R. Feasibility of assessment of skeletal muscle mass on a single cross-sectional image at the level of the fourth thoracic vertebra. *Eur J Radiol.* Sep 2021;142:109879. doi:10.1016/j.ejrad.2021.109879
34. Bokshan SL, Han AL, DePasse JM, Eltorai AEM, Marcaccio SE, Palumbo MA, Daniels AH. Effect of Sarcopenia on Postoperative Morbidity and Mortality After Thoracolumbar Spine Surgery. Article. *Orthopedics.* Nov-Dec 2016;39(6):E1159-E1164. doi:10.3928/01477447-20160811-02
35. Zopfs D, Bousabarah K, Lennartz S, dos Santos DP, Schlaak M, Theurich S, Reimer RP, Maintz D, Haneder S, Hokamp NG. Evaluating body composition by combining quantitative spectral detector computed tomography and deep learning-based image segmentation. Article. *European Journal of Radiology.* Sep 2020;130:8. 109153. doi:10.1016/j.ejrad.2020.109153
36. Waduud MA, Adusumilli P, Drozd M, Bailey MA, Cuthbert G, Hammond C, Scott JA. Volumetric versus single slice measurements of core abdominal muscle for sarcopenia. Article. *British Journal of Radiology.* 2019;92(1097):7. 20180434. doi:10.1259/bjr.20180434
37. Amini B, Boyle SP, Boutin RD, Lenchik L. Approaches to Assessment of Muscle Mass and Myosteatosis on Computed Tomography: A Systematic Review. *J Gerontol A Biol Sci Med Sci.* Sep 15 2019;74(10):1671-1678. doi:10.1093/gerona/glz034
38. Jovanovic N, Chinnery T, Mattonen SA, Palma DA, Doyle PC, Theurer JA. Sarcopenia in head and neck cancer: A scoping review. *PLoS One.* 2022;17(11):e0278135. doi:10.1371/journal.pone.0278135
39. Lin TY, Chen YF, Wu WT, Han DS, Tsai IC, Chang KV, Ozcakar L. Impact of sarcopenia on the prognosis and treatment of lung cancer: an umbrella review. *Discov Oncol.* Oct 28 2022;13(1):115. doi:10.1007/s12672-022-00576-0
40. Mauro E, Diaz JM, Garcia-Oliveira L, Spina JC, Savluk L, Zalazar F, Saidman J, De Santibanes M, Pekolj J, De Santibanes E, Crespo G, Abraldes JG, Gadano A. Sarcopenia HIBA score predicts sarcopenia and mortality in patients on the liver transplant waiting list. *Hepatol Commun.* Jul 2022;6(7):1699-1710. doi:10.1002/hep4.1919
41. Kim DW, Ahn H, Kim KW, Lee SS, Kim HJ, Ko Y, Park T, Lee J. Prognostic Value of Sarcopenia and Myosteatosis in Patients with Resectable Pancreatic Ductal Adenocarcinoma. *Korean J Radiol.* Nov 2022;23(11):1055-1066. doi:10.3348/kjr.2022.0277
42. Derstine BA, Holcombe SA, Ross BE, Wang NC, Su GL, Wang SC. Skeletal muscle cutoff values for sarcopenia diagnosis using T10 to L5 measurements in a healthy US population. *Sci Rep.* Jul 27 2018;8(1):11369. doi:10.1038/s41598-018-29825-5
43. Arayne AA, Gartrell R, Qiao J, Baird PN, Yeung JM. Comparison of CT derived body composition at the thoracic T4 and T12 with lumbar L3 vertebral levels and their utility in patients with rectal cancer. *BMC Cancer.* Jan 16 2023;23(1):56. doi:10.1186/s12885-023-10522-0
44. Kim EY, Kim YS, Park I, Ahn HK, Cho EK, Jeong YM, Kim JH. Evaluation of sarcopenia in small-cell lung cancer patients by routine chest CT. Article. *Supportive Care in Cancer.* Nov 2016;24(11):4721-4726. doi:10.1007/s00520-016-3321-0


45. Sato S, Sato M, Shinohara H. Significance of preoperative evaluation of skeletal muscle index and immune-nutritional status for patients with early-stage non-small cell lung cancer. *Gen Thorac Cardiovasc Surg*. Jun 2023;71(6):354-362. doi:10.1007/s11748-022-01899-z
46. Liu S, Han X, Li J, Xie X, Yang Y, Jiang W, Liu L, Liu Z. Feasibility of using chest computed tomography (CT) imaging at the first lumbar vertebra (L1) level to assess skeletal muscle mass: a retrospective study. *PeerJ*. 2023;11:e16652. doi:10.7717/peerj.16652
47. Mathur S, Rodrigues N, Mendes P, Rozenberg D, Singer LG. Computed Tomography–Derived Thoracic Muscle Size as an Indicator of Sarcopenia in People With Advanced Lung Disease. *Cardiopulmonary Physical Therapy Journal*. 2017;28(3):99-105. doi:10.1097/cpt.000000000000054
48. Kim YS, Kim EY, Kang SM, Ahn HK, Kim HS. Single cross-sectional area of pectoralis muscle by computed tomography - correlation with bioelectrical impedance based skeletal muscle mass in healthy subjects. Article. *Clinical Physiology and Functional Imaging*. Sep 2017;37(5):507-511. doi:10.1111/cpf.12333
49. Moon SW, Choi JS, Lee SH, Jung KS, Jung JY, Kang YA, Park MS, Kim YS, Chang J, Kim SY. Thoracic skeletal muscle quantification: low muscle mass is related with worse prognosis in idiopathic pulmonary fibrosis patients. *Respir Res*. Feb 15 2019;20(1):35. doi:10.1186/s12931-019-1001-6
50. Kalyani RR, Corriere M, Ferrucci L. Age-related and disease-related muscle loss: the effect of diabetes, obesity, and other diseases. *Lancet Diabetes Endocrinol*. Oct 2014;2(10):819-29. doi:10.1016/S2213-8587(14)70034-8
51. Gao H, Wang J, Zou X, Zhang K, Zhou J, Chen M. High blood urea nitrogen to creatinine ratio is associated with increased risk of sarcopenia in patients with chronic obstructive pulmonary disease. *Exp Gerontol*. Nov 2022;169:111960. doi:10.1016/j.exger.2022.111960
52. Nemeč U, Heidinger B, Sokas C, Chu L, Eisenberg RL. Diagnosing Sarcopenia on Thoracic Computed Tomography: Quantitative Assessment of Skeletal Muscle Mass in Patients Undergoing Transcatheter Aortic Valve Replacement. *Acad Radiol*. Sep 2017;24(9):1154-1161. doi:10.1016/j.acra.2017.02.008
53. Wang Y, Chai L, Chen Y, Liu J, Wang Q, Zhang Q, Qiu Y, Li D, Chen H, Shen N, Shi X, Wang J, Xie X, Li M. Quantitative CT parameters correlate with lung function in chronic obstructive pulmonary disease: A systematic review and meta-analysis. *Front Surg*. 2022;9:1066031. doi:10.3389/fsurg.2022.1066031
54. Panthofer AM, Olson SL, Harris DG, Matsumura JS. Derivation and validation of thoracic sarcopenia assessment in patients undergoing thoracic endovascular aortic repair. Article; Proceedings Paper. *Journal of Vascular Surgery*. May 2019;69(5):1379-1386. doi:10.1016/j.jvs.2018.08.180
55. Bril SI, Wendrich AW, Swartz JE, Wegner I, Pameijer F, Smid EJ, Bol GH, Pothen AJ, de Bree R. Interobserver agreement of skeletal muscle mass measurement on head and neck CT imaging at the level of the third cervical vertebra. *Eur Arch Otorhinolaryngol*. Apr 2019;276(4):1175-1182. doi:10.1007/s00405-019-05307-w
56. de Bree R, Meerkerk CDA, Halmos GB, Makitie AA, Homma A, Rodrigo JP, Lopez F, Takes RP, Vermorken JB, Ferlito A. Measurement of Sarcopenia in Head and Neck Cancer Patients and Its Association With Frailty. *Front Oncol*. 2022;12:884988. doi:10.3389/fonc.2022.884988
57. Charki N, Bril SI, Smid EJ, de Jong PA, de Bree R. Cut-off values for low skeletal muscle mass at the level of the third cervical vertebra (C3) in patients with head and neck cancer. *Quant Imaging Med Surg*. Jun 2022;12(6):3024-3033. doi:10.21037/qims-21-911



58. Barnard R, Tan J, Roller B, Chiles C, Weaver AA, Boutin RD, Kritchevsky SB, Lenchik L. Machine Learning for Automatic Paraspinous Muscle Area and Attenuation Measures on Low-Dose Chest CT Scans. Article. *Academic Radiology*. Dec 2019;26(12):1686-1694. doi:10.1016/j.acra.2019.06.017
59. Lenchik L, Heacock L, Weaver AA, Boutin RD, Cook TS, Itri J, Filippi CG, Gullapalli RP, Lee J, Zagurovskaya M, Retson T, Godwin K, Nicholson J, Narayana PA. Automated Segmentation of Tissues Using CT and MRI: A Systematic Review. *Acad Radiol*. Dec 2019;26(12):1695-1706. doi:10.1016/j.acra.2019.07.006
60. Laino ME, Generali E, Tommasini T, Angelotti G, Aghemo A, Desai A, Morandini P, Stefanini GG, Lleo A, Voza A, Savevski V. An individualized algorithm to predict mortality in COVID-19 pneumonia: a machine learning based study. *Arch Med Sci*. 2022;18(3):587-595. doi:10.5114/aoms/144980
61. Pella D, Toth S, Paralic J, Gonsorcik J, Fedacko J, Jarcuska P, Pella D, Pella Z, Sabol F, Jankajova M, Valocik G, Putrya A, Kirschova A, Plachy L, Rabajdova M, Hunavy M, Kafkova B, Doci I, Timkova S, Dvorožnakova M, Babic F, Butka P, Dimunova L, Marekova M, Paralicova Z, Majernik J, Luczy J, Janosik J, Kmec M. The possible role of machine learning in detection of increased cardiovascular risk patients - KSC MR Study (design). *Arch Med Sci*. 2022;18(4):991-997. doi:10.5114/aoms.2020.99156
62. Kullberg J, Hedstrom A, Brandberg J, Strand R, Johansson L, Bergstrom G, Ahlstrom H. Automated analysis of liver fat, muscle and adipose tissue distribution from CT suitable for large-scale studies. Article. *Scientific Reports*. Sep 2017;7:11. 10425. doi:10.1038/s41598-017-08925-8
63. Lee H, Troschel FM, Tajmir S, Fuchs G, Mario J, Fintelmann FJ, Do S. Pixel-Level Deep Segmentation: Artificial Intelligence Quantifies Muscle on Computed Tomography for Body Morphometric Analysis. Article. *Journal of Digital Imaging*. Aug 2017;30(4):487-498. doi:10.1007/s10278-017-9988-z
64. Weston AD, Korfiatis P, Kline TL, Philbrick KA, Kostandy P, Sakinis T, Sugimoto M, Takahashi N, Erickson BJ. Automated Abdominal Segmentation of CT Scans for Body Composition Analysis Using Deep Learning. Article. *Radiology*. Mar 2019;290(3):669-679. doi:10.1148/radiol.2018181432
65. Polan DF, Brady SL, Kaufman RA. Tissue segmentation of computed tomography images using a Random Forest algorithm: a feasibility study. Article. *Physics in Medicine and Biology*. Sep 2016;61(17):6553-6569. doi:10.1088/0031-9155/61/17/6553
66. Somasundaram E, Deaton J, Kaufman R, Brady S. Fully automated tissue classifier for contrast-enhanced CT scans of adult and pediatric patients. Article. *Physics in Medicine and Biology*. Jul 2018;63(13):15. 135009. doi:10.1088/1361-6560/aac944
67. Borrelli A, Pecoraro M, Del Giudice F, Cristofani L, Messina E, Dehghanpour A, Landini N, Roberto M, Perotti S, Muscaritoli M, Santini D, Catalano C, Panebianco V. Standardization of Body Composition Status in Patients with Advanced Urothelial Tumors: The Role of a CT-Based AI-Powered Software for the Assessment of Sarcopenia and Patient Outcome Correlation. *Cancers (Basel)*. May 29 2023;15(11)doi:10.3390/cancers15112968
68. Herrod PJJ, Boyd-Carson H, Doleman B, Trotter J, Schlichtemeier S, Sathanapally G, Somerville J, Williams JP, Lund JN. Quick and simple; psoas density measurement is an independent predictor of anastomotic leak and other complications after colorectal resection. Article. *Techniques in Coloproctology*. Feb 2019;23(2):129-134. doi:10.1007/s10151-019-1928-0
69. Shen DG, Wu GR, Suk HI. Deep Learning in Medical Image Analysis. In: Yarmush ML, ed. *Annual Review*

- of *Biomedical Engineering*, Vol 19. Annual Reviews; 2017:221-248. *Annual Review of Biomedical Engineering*.
70. Limkin EJ, Sun R, Derclé L, Zacharaki EI, Robert C, Reuze S, Schernberg A, Paragios N, Deutsch E, Ferte C. Promises and challenges for the implementation of computational medical imaging (radiomics) in oncology. Review. *Annals of Oncology*. Jun 2017;28(6):1191-1206. doi:10.1093/annonc/mdx034
71. Hsu TMH, Schawkat K, Berkowitz SJ, Wei JL, Makoyeva A, Legare K, DeCicco C, Paez SN, Wu JSH, Szolovits P, Kikinis R, Moser AJ, Goehler A. Artificial intelligence to assess body composition on routine abdominal CT scans and predict mortality in pancreatic cancer- A recipe for your local application. Article. *European Journal of Radiology*. Sep 2021;142:7. 109834. doi:10.1016/j.ejrad.2021.109834
72. Bedrikovetski S, Seow W, Kroon HM, Traeger L, Moore JW, Sammour T. Artificial intelligence for body composition and sarcopenia evaluation on computed tomography: A systematic review and meta-analysis. Review. *European Journal of Radiology*. Apr 2022;149:9. 110218. doi:10.1016/j.ejrad.2022.110218
73. Ackermans L, Volmer L, Wee L, Brecheisen R, Sanchez-Gonzalez P, Seiffert AP, Gomez EJ, Dekker A, Ten Bosch JA, Damink S, Blokhuis TJ. Deep Learning Automated Segmentation for Muscle and Adipose Tissue from Abdominal Computed Tomography in Polytrauma Patients. Article. *Sensors*. Mar 2021;21(6):13. 2083. doi:10.3390/s21062083
74. Koitka S, Kroll L, Malamutmann E, Oezcelik A, Nensa F. Fully automated body composition analysis in routine CT imaging using 3D semantic segmentation convolutional neural networks. Article. *European Radiology*. Apr 2021;31(4):1795-1804. doi:10.1007/s00330-020-07147-3
75. Beetz NL, Maier C, Segger L, Shnayien S, Trippel TD, Lindow N, Bousabarah K, Westerhoff M, Fehrenbach U, Geisel D. First PACS-integrated artificial intelligence-based software tool for rapid and fully automatic analysis of body composition from CT in clinical routine. *JCSM Clinical Reports*. 2022;7(1):3-11. doi:<https://doi.org/10.1002/crt2.44>
76. Just IA, Schoenrath F, Roehrich L, Heil E, Stein J, Auer TA, Fehrenbach U, Potapov E, Solowjowa N, Balzer F, Geisel D, Braun J, Boening G. Artificial intelligence-based analysis of body composition predicts outcome in patients receiving long-term mechanical circulatory support. *J Cachexia Sarcopenia Muscle*. Feb 2024;15(1):270-280. doi:10.1002/jcsm.13402
77. Liefers JR, Bathe OF, Fassbender K, Winget M, Baracos VE. Sarcopenia is associated with postoperative infection and delayed recovery from colorectal cancer resection surgery. *Br J Cancer*. Sep 4 2012;107(6):931-6. doi:10.1038/bjc.2012.350
78. McDonald ML, Diaz AA, Ross JC, San Jose Estepar R, Zhou L, Regan EA, Eckbo E, Muralidhar N, Come CE, Cho MH, Hersh CP, Lange C, Wouters E, Casaburi RH, Coxson HO, Macnee W, Rennard SI, Lomas DA, Agusti A, Celli BR, Black-Shinn JL, Kinney GL, Lutz SM, Hokanson JE, Silverman EK, Washko GR. Quantitative computed tomography measures of pectoralis muscle area and disease severity in chronic obstructive pulmonary disease. A cross-sectional study. *Ann Am Thorac Soc*. Mar 2014;11(3):326-34. doi:10.1513/AnnalsATS.201307-229OC
79. Liu Y, Ji F, Nguyen MH. Sarcopenia in cirrhosis: epidemiology, diagnosis, management and prognosis. *Curr Opin Gastroenterol*. May 1 2023;39(3):131-139. doi:10.1097/MOG.0000000000000922
80. Jiang C, Wang Y, Fu W, Zhang G, Feng X, Wang X, Wang F, Zhang L, Deng Y. Association between sarcopenia and prognosis of hepatocellular carcinoma: A systematic review and meta-analysis. *Front Nutr*. 2022;9:978110. doi:10.3389/fnut.2022.978110

81. Wang M, Tang H, Chen X, Liu J, Hu N, Cui W, Zhang C, Xie C, Chen X. Opportunistic Muscle Evaluation During Chest CT Is Associated With Vertebral Compression Fractures in Old Adults: A Longitudinal Study. *J Gerontol A Biol Sci Med Sci*. Feb 1 2024;79(2)doi:10.1093/gerona/glad162
82. Friedman RS, Tarasova A, Jain VR, Ye K, Mansour A, Haramati LB. Predictive Value of CT Biomarkers in Lung Transplantation Survival: Preliminary Investigation in a Diverse, Underserved, Urban Population. *Lung*. Dec 2023;201(6):581-590. doi:10.1007/s00408-023-00650-6
83. Olmez T, Karakose E, Bozkurt H, Pence HH, Gulmez S, Aray E, Bulut CI, Sert OZ, Polat E, Duman M. Sarcopenia is associated with increased severe postoperative complications after colon cancer surgery. *Arch Med Sci*. 2021;17(2):361-367. doi:10.5114/aoms.2019.88621
84. Icard P, Schussler O, Loi M, Bobbio A, Lupo AM, Wislez M, Iannelli A, Fournel L, Damotte D, Alifano M. Pre-Disease and Pre-Surgery BMI, Weight Loss and Sarcopenia Impact Survival of Resected Lung Cancer Independently of Tumor Stage. *Cancers (Basel)*. Jan 22 2020;12(2)doi:10.3390/cancers12020266
85. Pekařová A, Pekař M, Daniš D, Nováková Z. CT evaluated sarcopenia signals: Shorter survival for small cell lung cancer patients. *Physiol Res*. Dec 31 2021;70(S3):S381-s386. doi:10.33549/physiolres.934816
86. Dong Q. Fully-automated Segmentation of Muscle Measurement on CT in Detecting Central Sarcopenia: A Trend of Standardization. Editorial Material. *Academic Radiology*. Mar 2020;27(3):321-322. doi:10.1016/j.acra.2019.12.003
87. Rozynek M, Kucybala I, Urbanik A, Wojciechowski W. Use of artificial intelligence in the imaging of sarcopenia: A narrative review of current status and perspectives. Review. *Nutrition*. Sep 2021;89:7. 111227. doi:10.1016/j.nut.2021.111227
88. Cespedes Feliciano EM, Popuri K, Cobzas D, Baracos VE, Beg MF, Khan AD, Ma C, Chow V, Prado CM, Xiao J, Liu V, Chen WY, Meyerhardt J, Albers KB, Caan BJ. Evaluation of automated computed tomography segmentation to assess body composition and mortality associations in cancer patients. *J Cachexia Sarcopenia Muscle*. Oct 2020;11(5):1258-1269. doi:10.1002/jcsm.12573
89. Lee YS, Hong N, Witanto JN, Choi YR, Park J, Decazes P, Eude F, Kim CO, Chang Kim H, Goo JM, Rhee Y, Yoon SH. Deep neural network for automatic volumetric segmentation of whole-body CT images for body composition assessment. *Clin Nutr*. Aug 2021;40(8):5038-5046. doi:10.1016/j.clnu.2021.06.025
90. Chen L, Wu F, Huang J, Yang J, Fan W, Nie Z, Jiang H, Wang J, Xia W, Yang F. Well-Aerated Lung and Mean Lung Density Quantified by CT at Discharge to Predict Pulmonary Diffusion Function 5 Months after COVID-19. *Diagnostics (Basel)*. Nov 23 2022;12(12)doi:10.3390/diagnostics12122921



Preprint