

# Assessment of special hospital meals for diabetic patients. Discrepancies between provided meals, model menus and dietary guidelines: a hospital diet medical investigation (HDMI) study

Alicja M. Jodczyk<sup>1</sup>, Alicja Kucharska<sup>2</sup>, Mariusz Panczyk<sup>3</sup>, Natalia Adamczyk<sup>4</sup>, Karolina Turlej<sup>2</sup>, Piotr Wierzbński<sup>1</sup>, Marcin Barylski<sup>5</sup>, Adam R. Poliwczak<sup>5</sup>, Maciej Janiszewski<sup>6</sup>, Artur Mamcarz<sup>1</sup>, Daniel Śliż<sup>1</sup>

<sup>1</sup>Third Department of Internal Medicine and Cardiology, Medical University of Warsaw, Warsaw, Poland

<sup>2</sup>Department of Human Nutrition, Medical University of Warsaw, Warsaw, Poland

<sup>3</sup>Department of Education and Research in Health Sciences, Medical University of Warsaw, Warsaw, Poland

<sup>4</sup>Polish Society of Lifestyle Medicine, Warsaw, Poland

<sup>5</sup>Department of Internal Medicine, Rehabilitation and Physical Medicine, Medical University of Lodz, Lodz, Poland

<sup>6</sup>Department of Heart Failure and Cardiac Rehabilitation, Medical University of Warsaw, Warsaw, Poland

## Corresponding author:

Alicja M. Jodczyk  
Third Department of  
Internal Medicine  
and Cardiology  
Medical University  
of Warsaw  
Warsaw, Poland  
E-mail: alicja.jodczyk@gmail.com

**Submitted:** 6 May 2024; **Accepted:** 16 July 2024

**Online publication:** 6 August 2024

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms/191298>

Copyright © 2024 Termedia & Banach

## Abstract

**Introduction:** In Poland, it is estimated that more than 2 million individuals have diabetes. Intervening in modifiable risk factors can effectively prevent and delay the onset of type 2 diabetes. Previous reports claimed that the Polish healthcare system did not guarantee proper nutrition and nutritional education. This study aimed to examine the dietary provisions for diabetic patients in Poland's hospitals and evaluate their compliance with the Good Meal in Hospital (*Dobry Positek w Szpitalu*) guidelines.

**Material and methods:** Hospital workers were asked to complete the survey regarding hospital (degree of reference, number of beds in total and internal medicine unit, availability of a special diet for diabetic patients, performing nutritional education and presence of a dietitian) and attach menus from meals received by patients from 10 consecutive days. The menus were analyzed in a computer program for assessment of institutional nutrition and then compared to a self-made model menu and the Good Meal in Hospital guidelines. These are the first such recommendations in Poland that introduce a coherent system to plan meals in hospitals. For each nutritional value, the mean (M), standard deviation (SD), and range (min–max.) were calculated. To assess differences in means for nutrients and products a one-way analysis of variance (ANOVA) followed by Dunnett's two-tailed test was used.

**Results:** Seventy menus from seven hospitals were examined. Five hospitals met eight and two hospitals met seven out of thirteen criteria. The discrepancies particularly concerned excessive levels of saturated fatty acids and mono- and disaccharides, and insufficient amounts of legumes and fish in the hospital diets. Only four out of seven hospitals had a resident dietitian present.

**Conclusions:** Providing meals containing typical nutritional errors representing the pattern of the Polish population during hospitalization may lead to potential post-discharge dietary errors. They may result in deterioration in glycemic regulation and lipid profiles, and heightened susceptibility to complications, including elevated cardiovascular risk.

**Key words:** hospital nutrition, diabetes, clinical care.

## Introduction

Data indicate that 537 million adults are living with diabetes mellitus (DM) worldwide [1]. In Poland, over 2 million people have DM, 25% of them unaware of their condition [2]. Forecasts predict that this number will double in the next 15–20 years [2]. Their projected lifespan is reduced, since DM elevates the likelihood of cardiovascular (CV) diseases and the susceptibility to other illnesses [3]. According to guidelines, lifestyle interventions are paramount in treating DM and in preventing the development of atherosclerotic cardiovascular disease [4]. Diets that favor a higher intake of whole grains and green leafy vegetables and a lower intake of refined grains, red and processed meat, and sugar-sweetened beverages have been linked with reduced risk of type 2 DM [5–8]. Adhering to Mediterranean-style or Dietary Approaches to Stop Hypertension (DASH) or diets which exclude animal products decrease the likelihood of developing type 2 DM and reduce risk of CV complications [9–11]. Nutritional education is a continuous, permanent process and an integral part of the treatment during every doctor's visit or nursing consultation [12]. Standards of Medical Care in Diabetes recommend reeducation of the patient when complication factors influencing self-management such as new health conditions occur [13].

The most recent and reliable data regarding nutrition in Polish hospitals were presented in the report of the Supreme Audit Office from 2018 [14]. It revealed that the healthcare system did not guarantee proper nutrition. There were no nutritional standards, health requirements or methods for assessing the quality of nutrition and rules for employing dietitians in hospital wards. Such gaps in the system and low financial outlays resulted in meals that were inadequate for patients' health conditions, prepared from low-quality raw materials, with unsuitable energy and nutritional values. The food provided by hospitals could harm patients' health. The problem of poor-quality hospital nutrition is also noticeable in other Central European countries, such as the Czech Republic and Germany [15, 16]. Since 1991, hospital nutrition in the Czech Republic has been guided by recommendations of the Ministry of Health. As a result, individual hospitals may offer very different diets. They have to meet criteria for nutrients and energy value but often lack quality and taste and do not fulfil patients' needs. In Germany, the Physicians Association for Nutrition prepared an open letter to the Ministry of Health in which they requested changes in hospital meals. Nutrition there is included in the so-called "non-medical services" and therefore competes for financing with 12 other services (including training and fur-

ther education, IT, administration, laundry, control and cleaning).

The aim of this study was to investigate the nutrition provided to diabetic patients in hospitals located in Poland and assess their adherence to the Good Meal in Hospital recommendations, prepared at the behest of the Polish Ministry of Health [16]. We hypothesized that the menus offered to diabetic patients in Polish hospitals failed to adhere to recommendations and lacked proper balance.

## Material and methods

### Study design and setting

The comparative cross-sectional study was conducted between January and August 2022. Menus for a diabetic diet were collected. Inclusion criteria were: (1) location in Poland (2) being a public hospital, (3) having a general medicine ward, (4) submitting meal plans from 10 consecutive days, which contained all essential information for analysis (precise ingredients and their weight), and (5) completing the entire questionnaire. Exclusion criteria were: (1) being a private hospital, (2) focusing on specific illnesses or patient demographics, e.g. military personnel, (3) sending incomplete surveys or (4) inaccurate and unreadable menus (without the weight of the ingredients or containing only general information, e.g. ham sandwich). The aims of the study, information about anonymity and terms of participation were mentioned before beginning the survey. Completing and submitting the form was regarded as providing informed consent to participate in the study.

### Data collection

Participants – heads of departments, specialists and resident doctors from various districts of Poland – were reached by an email about the study. They were asked to complete a survey created with Google Forms and send the meal menus received by diabetic patients for the upcoming 10 days.

### Research tools

Questions involved: (1) the hospital's degree of reference (1<sup>st</sup>/2<sup>nd</sup>/3<sup>rd</sup> degree) – the first degree provides fundamental medical procedures, the third degree offers the most specialized procedures; (2) the total amount of beds in the hospital and the internal medicine unit; (3) the attendance of a person delivering nutritional education; (4) place of meal preparation – hospital kitchen or outsourced catering services; (5) access to diets tailored for patients with diabetes. Next, the received menus were compared with the self-created model menu. It was prepared using widely accessible

and relatively affordable ingredients to meet the nutritional requirements of individuals with diabetes and at risk of cardiovascular disease. The form and model menu are available in the Supplementary Materials section. Then, all menus were evaluated in the DIETA 6 computer program and compared with the Good Meal in Hospital governmental recommendations. The DIETA 6 computer program was created by the National Institute of Public Health – National Institute of Hygiene in Warsaw and is tailored to the Polish population. The Good Meal in Hospital recommendations are the first such recommendations in Poland that introduce a coherent system to plan meals in hospitals [17]. They introduce an average weighted standard for energy and nutrients for the entire hospital population, regardless of gender, types of diets, their characteristics, recommended and contraindicated foods. A summary of the recommendations is presented in Table I.

### Statistical analysis

Statistical analyses were carried out using Statistica 13.3 software by TIBCO Software in Palo Alto, California, United States. DIETA 6 software was used to calculate the nutritional values of the received menus. For every nutritional value, the mean (M), standard deviation (SD), and range (min-max) were calculated. To assess discrepancies in means for both nutrients and products among different hospitals and the standardized menu, we used a one-way ANOVA followed by a post-hoc Dunnett's two-tailed test. Dunnett's test is a multiple comparison method that contrasts each mean with a single control in many-to-one comparisons. The null hypothesis was rejected when the *p*-value was below 0.05, indicating a statistically significant difference in means.

### Results

A total of 70 daily menus for diabetic patients from seven hospitals underwent analysis (A – 1035 beds, B – 452 beds, C – 683 beds, D – 677 beds, E – 363 beds, F – 176 beds, G – 321 beds). Five of the examined hospitals were third degree reference hospitals (A, C, D, E, G), one was second (B) degree and one first degree (F). Hospitals 1, 2, and 3 were located in the Mazowieckie Voivodeship, hospitals 4, 6, and 7 in the Śląskie Voivodeship, hospital 5 in the Łódzkie Voivodeship. A special diet for diabetic patients was available in every hospital. Nutrition education was available in six hospitals (A, B, D, E, F, G), yet only four of them had a resident dietician (D, E, F, G). Four hospitals undertook the culinary preparations internally (C, D, F, G), while three hospitals opted to engage external catering services (A, B, E).

There was a statistically significant difference in mean: energy, energy from carbohydrates, total carbohydrates per 1000 kcal, fiber per 1000 kcal, mono- and disaccharides per 1000 kcal, and percentage of energy from mono- and carbohydrates. Compared to the standardized menu, there was a statistically greater mean amount of energy in hospitals B, C and lower in hospitals D and E; a greater percentage of energy from carbohydrates in hospitals F and G and lower in hospital B; lower mean fiber per 1000 kcal in hospitals B, D, E; a greater mean amount of carbohydrates per 1000 kcal in hospitals F and G and lower in hospital B; a greater mean amount of mono- and disaccharides per 1000 kcal in hospitals F and G and lower in hospital D; a greater percentage of energy from mono- and disaccharides in hospitals F and G and lower in hospital D. Table II displays precise values.

There was a statistically significant difference in the mean: total protein, percentage of energy

**Table I.** Characteristics of a diet restricting easily digestible carbohydrates based on the Good Meal in Hospital recommendations [17]

The daily caloric value of meals should be 2000–2400 kcal
Protein should represent no more than 10–20% of total energy and the amount 25–50 g/1000 kcal
Fat should represent no more than 20–30% of total energy and the amount 22–33 g/1000 kcal
Saturated fatty acids should represent no more than 10% of energy and the amount < 11 g/1000 kcal
Carbohydrates should represent no more than 45–65% of energy and the amount 113–163 g/1000 kcal
Mono- and disaccharides should represent no more than 10% of energy and the amount < 25 g/1000 kcal
The amount of sodium should be < 2000 mg/day
The amount of fiber should be at least 15 g/1000 kcal
Whole grain cereal products should be provided at least twice a day
Each meal should incorporate vegetables or fruit, with a minimum daily intake of 400 g (excluding potatoes and sweet potatoes). Vegetables should predominate, constituting at least three portions
Legumes or legume preserves should be included in meals at least three times within 10 days
Fish or fish preserves should be included in meals at least three times within 10 days
The number of meals should range from 4 to 6 per day, and it should remain consistent every day.

**Table II.** Energy and carbohydrate content

Hospital/unit	M	SD	Min.	Max.	P-value <sup>a</sup>
Energy [kcal] ( $F = 22.141$ ; $p < 0.001^b$ )					
A	2054.1	234.5	1726.4	2397.1	0.998
B	2266.4	100.7	2111.4	2394.4	0.033
C	2332.9	108.3	2180.1	2524.2	0.002
D	1620.8	65.4	1514.8	1716.5	< 0.001
E	1759.1	116.3	1572.9	1940.6	0.016
F	2130.0	155.3	1881.7	2389.5	0.608
G	2162.3	192.1	1864.2	2439.1	0.344
Reference	2015.8	29.3	1993.1	2055.8	–
Percentage of energy from carbohydrates [kcal] ( $F = 25.600$ ; $p < 0.001^b$ )					
A	53.8	4.3	46.2	59.4	0.900
B	41.8	3.1	38.3	46.9	< 0.001
C	53.7	2.6	50.0	58.0	0.935
D	54.3	1.4	52.2	56.5	0.802
E	52.8	3.3	48.4	58.1	1.000
F	59.4	2.6	55.5	63.6	< 0.001
G	59.6	3.1	55.9	65.3	< 0.001
Reference	52.2	2.6	48.5	55.7	–
Fiber/1000 kcal [g/kcal] ( $F = 7.033$ , $p < 0.001^b$ )					
A	20.6	2.0	16.4	22.5	0.996
B	17.1	1.9	14.9	20.3	0.004
C	18.2	2.6	15.3	22.4	0.044
D	15.6	1.5	13.7	17.6	< 0.001
E	16.5	1.5	13.7	19.8	< 0.001
F	18.2	1.8	15.7	19.8	0.040
G	18.1	2.1	14.1	21.6	0.035
Reference	21.1	1.5	18.8	22.9	–
Carbohydrates/1000 kcal [g/kcal] ( $F = 22.490$ , $p < 0.001^b$ )					
A	144.6	11.7	124.2	159.3	0.939
B	112.7	8.4	103.8	127	< 0.001
C	143.1	7.6	132.6	155.9	0.996
D	143.1	3.7	137.0	148.9	0.997
E	139.7	9	128.2	154.8	1.000
F	157.4	6.7	146.4	167.2	0.003
G	157.9	8.4	146.5	173.7	0.002
Reference	140.7	6.6	131.1	149.5	–
A total of mono- and disaccharides/1000 kcal [g/1000 kcal] ( $F = 20.194$ ; $p < 0.001^b$ )					
A	32.3	6.7	18.5	41.9	0.209
B	21.6	3.6	17.3	26.8	0.293
C	32.3	3.7	26.2	36.7	0.212
D	15.5	3.8	10.4	20.0	0.001
E	27.9	7.6	14.2	42.6	0.999
F	37.6	3.2	32.8	42.1	0.001
G	37.2	3.1	32.8	42.1	0.002
Reference	26.9	2.2	23.8	29.7	–
Percentage of energy from mono- and disaccharides [%] ( $F = 20.194$ , $p < 0.001^b$ )					
A	12.9	2.7	7.4	16.8	0.209
B	8.6	1.4	6.9	10.7	0.293

Table II. Cont.

Hospital/unit	M	SD	Min.	Max.	P-value <sup>a</sup>
C	12.9	1.5	10.5	14.7	0.212
D	6.2	1.5	4.2	8.0	0.001
E	11.2	3.0	5.7	17.0	0.999
F	15.1	1.3	13.1	16.9	0.001
G	14.9	1.3	13.1	16.9	0.002
Reference	10.7	0.9	9.5	11.9	–

M – mean, SD – standard deviation, <sup>a</sup>Dunnett's post hoc test (multiple comparisons to a reference: many-to-one comparisons); <sup>b</sup>one-way analysis of variance.

Table III. Protein content

Hospital/unit	M	SD	Min.	Max.	P-value <sup>a</sup>
Total amount of protein [g] ( $F = 25.246; p < 0.001^b$ )					
A	79.2	10.4	70.6	101.9	< 0.001
B	120.4	8.7	106.2	135	0.033
C	92.1	10.0	74.9	105.5	0.022
D	73.8	11.9	57.4	93.7	< 0.001
E	77.0	7.2	66.6	90.5	< 0.001
F	85.3	5.2	75.6	94.5	< 0.001
G	86.1	5.8	75.2	94.5	< 0.001
Reference	106.1	6.7	99.8	116.0	–
Percentage of energy from protein [%] ( $F = 21.362; p < 0.001^b$ )					
A	15.5	1.4	12.7	18.0	< 0.001
B	21.2	1.1	19.7	22.6	1.000
C	16.0	1.6	12.9	17.6	< 0.001
D	17.9	1.7	14.7	20.0	0.001
E	17.6	1.7	14.8	20.2	< 0.001
F	15.8	1.0	14.3	16.8	< 0.001
G	15.7	1.1	14.1	16.8	< 0.001
Reference	21.3	1.1	20.2	22.8	–
Animal-based protein/1000 kcal [g/kcal] ( $F = 11.049; p < 0.001^b$ )					
A	22.7	4.2	18.0	31.5	0.986
B	38.1	4.4	32.7	45.0	< 0.001
C	22.6	4.5	13.2	27.5	0.981
D	25.2	6.1	16.1	35.4	0.998
E	25.4	4.3	17.9	30.7	0.994
F	23.5	2.9	20.0	28.4	1.000
G	23.4	2.9	19.8	28.4	1.000
Reference	24.1	4.6	18.6	29.1	–
Plant based protein/1000 kcal [g/kcal] ( $F = 29.212; p < 0.001^b$ )					
A	16.0	2.3	12.6	19.3	< 0.001
B	14.6	1.7	12.3	16.8	< 0.001
C	16.7	2.7	13.9	21.7	< 0.001
D	19.9	1.0	18.7	21.4	< 0.001
E	18.3	1.7	14.8	20.5	< 0.001
F	16.5	1.1	14.8	18.0	< 0.001
G	16.5	1.4	13.7	18.0	< 0.001
Reference	28.2	2.2	24.9	30.7	–

M – mean, SD – standard deviation, <sup>a</sup>Dunnett's post hoc test (multiple comparisons to a reference: many-to-one comparisons), <sup>b</sup>one-way analysis of variance.

from protein, animal-based protein per 1000 kcal, and plant-based protein per 1000 kcal. Compared to the standardized menu there was a statistically significantly lower mean amount of total protein in hospitals A, C, D, E, F, G and a greater amount in hospital B; a lower percentage of energy from protein in hospitals A, C, D, E, F, G, a greater mean of animal-based protein per 1000 kcal in hospital B, and a lower mean amount of plant-based protein per 1000 kcal in every hospital. Precise values are presented in Table III.

Table IV displays fat content. A one-way ANOVA revealed that there was a statistically significant difference in the mean: total fat, fat per 1000 kcal, percentage of energy from fat, cholesterol and percentage of energy from saturated fat acids. Compared to the standardized menu there was a statistically significantly higher mean amount of total fat in hospitals B and C; a higher mean amount of fat per 1000 kcal in hospital B, a higher percentage of energy from fat in hospital B, a higher mean amount of cholesterol in hospitals A, B, C; and a higher percentage of energy from saturated fat acids in hospital A, B, C, E, F, G.

The hospitals showed significant differences in mean amounts of: fruits, vegetables, sugar in total, legumes, nuts and seeds, whole grain cereal products, red meat, and processed white and red meat. Compared to the model menu, lower mean amounts of fruits were observed in hospitals B, D, E, lower mean amounts of vegetables in hospitals A, C, D, E, F, G; greater mean amounts of sugar in hospitals A, C, E, F, G, greater mean amounts of potato in hospitals F and G; a lower amount of legumes in every hospital, a greater mean amount of whole grain products in hospital A and lower in hospital D; and a greater mean amount of red meat in hospitals B and E. Only one hospital offered nuts and in a statistically significantly lower amount than the model menu. Other differences were not statistically significant. Every hospital provided red meat; only the reference diet did not. Precise values are presented in Table V.

All hospitals successfully met five criteria concerning intake of protein, carbohydrate, fiber, whole grains, and vegetables and fruits. Hospitals A, B, C, D, E met eight out of thirteen criteria, while hospitals F and G met seven out of thirteen criteria. A summary of the results is presented in Table VI.

## Discussion

The results of our study show a significant discrepancy between the recommendations and meals received by the patients. Although every hospital provided a “diabetic diet”, its quality was inadequate and unadjusted. The lack of appropriate nutritional patterns promoted in the hospital raises the potential for patients to make nutrition-

al mistakes, which could negatively impact the control of DM. According to the recommendations of the Polish Diabetes Association, there is no universal diet for all people with diabetes [12]. Optimal macronutrient proportions for people with diabetes should be determined individually, taking into account age, physical activity, presence of diabetes complications, comorbidities, and personal preferences. Implementing the proper dietary recommendations can be challenging for diabetics, who are generally in good health. For those who undergo hospitalization or treatment, procedures maintaining glycemic control can be even more difficult. Failure to follow prescribed treatments, particularly dietary recommendations, is a critical issue in managing DM [18, 19].

The crucial role of healthcare providers in encouraging individuals with DM to adopt health-promoting behaviors has been highlighted in many studies [20, 21]. Research confirmed improved management of diabetes among patients who received care from a multidisciplinary team [19, 22]. Doctors and nurses play a significant role, as they interact with patients most frequently [23]. They are usually the first to notice glycemic irregularities and dietary and treatment errors. The Polish Diabetes Society emphasizes that education provided by authorized professionals (doctors, diabetes nurses, dietitian, diabetes educators) is crucial for proper diabetes management [12]. Effective communication between doctors and patients enhances patient satisfaction, boosts adherence to treatment plans, and results in better health outcomes [24]. Discussion of self-care successes and failures with physicians enables treatment to be individualized and increases the likelihood of success [25]. Detailed dietary recommendations should be individualized according to the patient’s needs and capabilities. Three out of seven examined hospitals did not have a dietician permanently available on the ward. In two of them, nutritional education was not provided at all. This presents a suboptimal scenario wherein delivering comprehensive assistance to a patient with diabetes becomes challenging.

Analysis of the menus demonstrates that they did not meet the criteria for a diet tailored to the needs of diabetic patients. The menus contained typical nutritional errors representing the nutritional pattern of the Polish population: insufficient intake of whole grain products, legumes and fish and excessive intake of meat and meat preserves [26]. Although all hospitals met the requirements regarding the amount and percentage of carbohydrates, only two out of seven hospitals met the criteria regarding the intake of mono- and disaccharides. Research and recommendations show a lack of sufficient scientific evidence for deter-

Table IV. Fat content

Hospital/unit	M	SD	Min.	Max.	P-value <sup>a</sup>
Fat [g] ( $F = 11.726$ ; $p < 0.001^b$ )					
A	70.7	17.1	51.7	106.6	0.407
B	93.3	11.6	79.8	112.4	< 0.001
C	79.4	10.1	65.3	100.9	0.019
D	50.0	1.6	47.4	52.4	0.529
E	58.4	9.9	39.5	78.2	1.000
F	58.6	12.0	44.9	79.3	1.000
G	59.0	11.7	44.9	81.0	1.000
Reference	60.1	7.6	49.9	70.2	–
Fat/1000 kcal [g/kcal] ( $F = 10.600$ ; $p < 0.001^b$ )					
A	34.1	5.2	28.0	45.7	0.252
B	41.1	4.5	34.9	46.9	< 0.001
C	34.0	3.5	27.5	40.9	0.279
D	30.9	1.5	29.2	33.3	0.998
E	33.0	3.8	25.1	40.3	0.554
F	27.3	3.7	22.1	33.8	0.768
G	27.2	4.0	21.9	33.5	0.727
Reference	29.9	4.1	24.3	35.2	–
Percentage of energy from fat [%] ( $F = 10.607$ ; $p < 0.001^b$ )					
A	30.6	4.6	25	40.9	0.170
B	37.0	3.9	31.1	42	< 0.001
C	30.4	3.2	24.5	36.6	0.224
D	27.8	1.3	26.1	29.5	0.983
E	29.7	3.0	24.5	36.0	0.417
F	24.8	3.3	19.6	30.2	0.916
G	24.7	3.7	19.5	29.9	0.890
Reference	26.5	3.6	21.5	31.3	–
Cholesterol [g] ( $F = 6.167$ ; $p < 0.001^b$ )					
A	329.2	100.6	173.8	445.9	0.024
B	421.3	129.6	305.7	649.0	< 0.001
C	330.9	103.4	193.4	486.9	0.022
D	177.3	93.0	99.1	316.7	1.000
E	258.1	87.8	152.3	412.5	0.469
F	228.4	65.1	180.8	372.3	0.868
G	229.2	64.5	180.8	359.9	0.859
Reference	177.3	97.7	93.4	318.4	–
Percentage of energy from saturated fat acids [%] ( $F = 21.777$ ; $p = 0.001^b$ )					
A	12.8	1.8	10.1	15.5	< 0.001
B	15.7	1.5	14.3	18.6	< 0.001
C	15.0	1.9	12.6	19.4	< 0.001
D	9.3	1.1	7.8	10.9	0.025
E	11.4	1.1	10.3	13.6	< 0.001
F	11.8	1.7	10.0	14.6	< 0.001
G	11.7	1.8	9.2	14.6	< 0.001
Reference	6.5	1.4	4.9	8.6	–

M – mean, SD – standard deviation, <sup>a</sup>Dunnett's post hoc test (multiple comparisons to a reference: many-to-one comparisons), <sup>b</sup>one-way analysis of variance.

**Table V.** Groups of products and sodium intake

Hospital/Unit	M	SD	Min.	Max.	P-value <sup>a</sup>
Fruits [g] ( $F = 6.855$ ; $p < 0.001^b$ )					
A	287.8	52.8	150.0	348.8	0.987
B	168.9	21.0	150.0	202.5	0.025
C	262.1	121.5	138.8	497.3	0.763
D	65.2	85.4	0.0	178.1	< 0.001
E	155.3	90.1	0.0	281.0	0.006
F	240.5	94.2	138.8	450.8	0.433
G	237.4	94.7	138.8	450.8	0.391
Reference	316.9	19.1	300.0	346.2	–
Vegetables [g] ( $F = 15.204$ ; $p < 0.001^b$ )					
A	484.4	89.3	334.5	602.4	0.001
B	761.3	185.0	482.1	1021.3	1.000
C	551.2	175.6	290.7	922.3	0.025
D	424.4	125.1	272.8	641.6	< 0.001
E	384.2	95.0	240.8	540.3	< 0.001
F	337.5	69.4	248.8	472.2	< 0.001
G	327.7	53.3	248.8	413.4	< 0.001
Reference	744.2	124.2	572.6	861.4	–
Fish [g] ( $F = 0.634$ ; $p = 0.726^b$ )					
A	35.3	39.2	0.0	83.3	0.997
B	33.3	88.2	0.0	233.3	0.995
C	19.6	40.9	0.0	120.0	0.839
D	11.9	31.5	0.0	83.3	0.714
E	12.0	37.9	0.0	120.0	0.654
F	8.8	27.9	0.0	88.2	0.571
G	8.8	27.9	0.0	88.2	0.571
Reference	49.0	109.6	0.0	245.0	–
Sugar total [g] ( $F = 44.278$ ; $p < 0.001^b$ )					
A	30.1	9.2	5.1	40.1	< 0.001
B	1.2	1.6	0.0	3.8	1.000
C	33.7	3.6	31.3	43.1	< 0.001
D	1.3	1.6	0.0	3.8	1.000
E	27.4	12.7	2.7	56.2	< 0.001
F	33.3	2.6	31.3	39.3	< 0.001
G	33.1	2.4	31.3	39.3	< 0.001
Reference	2.2	2.5	0.0	5.9	–
Legumes [g] ( $F = 43.320$ ; $p < 0.001^b$ )					
A	3.8	11.9	0.0	37.5	< 0.001
B	0.0	0.0	0.0	0.0	< 0.001
C	9.1	11.9	0.0	28.1	< 0.001
D	0.0	0.0	0.0	0.0	< 0.001
E	1.5	3.1	0.0	7.5	< 0.001
F	0.0	0.0	0.0	0.0	< 0.001
G	0.0	0.0	0.0	0.0	< 0.001
Reference	43.3	10.5	26.3	52.5	–



Table V. Cont.

Hospital/Unit	M	SD	Min.	Max.	P-value <sup>a</sup>
Nuts and seeds [g] ( $F = 350.946$ ; $p < 0.001^b$ )					
A	0.0	0.0	0.0	0.0	–
B	0.0	0.0	0.0	0.0	–
C	1.5	3.4	0.0	10.0	< 0.001
D	0.0	0.0	0.0	0.0	–
E	0.0	0.0	0.0	0.0	–
F	0.0	0.0	0.0	0.0	–
G	0.0	0.0	0.0	0.0	–
Reference	30.0	0.0	30.0	30.0	–
Whole grain cereal products [g] ( $F = 12.651$ ; $p < 0.001^b$ )					
A	252.0	6.3	250.0	270.0	0.009
B	240.0	0.0	240.0	240.0	0.094
C	184.5	47.8	50.0	220.0	0.978
D	130.0	0.0	130.0	130.0	0.003
E	218.0	15.5	200.0	230.0	0.675
F	180.0	0.0	180.0	180.0	0.896
G	187.0	60.7	90.0	340.0	0.995
Reference	196.0	8.9	180.0	200.0	–
Red meat [g] ( $F = 5.374$ ; $p < 0.001^b$ )					
A	12.6	26.5	0.0	102.3	0.668
B	81.5	53.9	0.0	159.9	< 0.001
C	16.6	37.9	0.0	131.2	0.312
D	16.7	52.4	0.0	191.8	0.380
E	56.9	56.3	0.0	132.5	0.001
F	21.3	43.2	0.0	115.1	0.103
G	21.3	43.2	0.0	115.1	0.103
Reference	0.0	0.0	0.0	0.0	–
Processed white and red meat [g] ( $F = 3.534$ ; $p = 0.002^b$ )					
A	34.0	15.2	0.0	51.0	0.366
B	58.1	22.0	28.3	100.7	0.924
C	39.5	15.7	17.0	69.1	0.769
D	26.8	13.1	0.0	45.3	0.103
E	43.0	20.9	14.2	76.8	0.969
F	31.9	11.5	14.4	47.2	0.264
G	31.9	11.5	14.4	47.2	0.264
Reference	49.7	32.1	0.0	76.8	–

M – mean, SD – standard deviation, <sup>a</sup>Dunnett's post hoc test (multiple comparisons to a reference: many-to-one comparisons), <sup>b</sup>one-way analysis of variance.

mining one optimal amount of carbohydrates in the diet of people with diabetes [10, 27]. Emphasis should be placed on high-quality, nutrient-dense carbohydrate sources that are rich in fiber. Intake of mono- and disaccharides should be limited to the minimum [12]. Consistent consumption of an adequate amount of dietary fiber is linked to a reduced risk of all-cause mortality in individuals with diabetes [28, 29]. The primary source of carbohydrates should be whole grain cereal products,

especially those with a low glycemic index (GI). Encouraging carbohydrate intake from vegetables, fruits, whole grains, legumes, and dairy products is recommended over consumption from other carbohydrate sources, especially those containing added fats, sugars, or sodium [4, 27]. Low-GI diets may be useful for glycemic control and may reduce body weight in people with prediabetes or diabetes [30]. A meta-analysis [30] revealed that low-GI diets effectively reduced glycated hemo-

**Table VI.** Summary of recommendations fulfilled by hospitals

Variable	A	B	C	D	E	F	G
Energy: 2000–2400 kcal/day	+	+	+	-	-	+	+
Protein: 25–50 g/1000 kcal and 10–20% of total energy	+	+	+	+	+	+	+
Fat: 22–33 g/1000 kcal and 20–30% of energy in total	-	-	-	+	+	+	+
Saturated fatty acids: No more than 10% of energy and < 11 g/1000 kcal	-	-	-	+	-	-	-
Carbohydrates: 113–163 g/1000 kcal and 45–65% of energy	+	+	+	+	+	+	+
Mono- and disaccharides: < 25 g/1000 kcal and no more than 10% of energy	-	+	-	+	-	-	-
Fiber: 15 g/1000 kcal	+	+	+	+	+	+	+
Sodium: < 2000 mg/day	-	-	-	-	-	-	-
Whole grain cereal products should be given at least twice per day	+	+	+	+	+	+	+
Vegetables or fruit should be added to each meal (minimum 400 g/day excluding potatoes and sweet potatoes); vegetables should predominate (at least 3 portions)	+	+	+	+	+	+	+
Legumes or legume preserves should be served at least 3 times in 10 days	-	-	+	-	+	-	-
Fish or fish preserves should be served at least 3 times in 10 days	+	-	-	-	-	-	-
Number of meals should be the same every day: 4–6 a day	+	+	+	-	+	-	-

(+) recommendation fulfilled, (-) recommendation not fulfilled.

globin, fasting glucose, BMI, total cholesterol, and LDL, but did not impact fasting insulin, HOMA-IR, HDL, triglycerides, or insulin requirements. In principle, every hospital adhered to the requirement of including two whole grain products daily. Upon closer examination, it becomes evident that this primarily consisted of whole-grain bread. Products such as oat flakes, brown rice, millet and buckwheat were either absent or only occasionally featured in other menus.

The fat composition was also inadequate. Three out of seven hospitals exhibited excessive fat intake, and only one met the criteria for limiting saturated fatty acids. According to the recommendations, the quality of fat is more important than its total quantity [12, 19]. Adhering to a Mediterranean eating pattern can enhance glycemic control and blood lipid levels and contribute to the reduction of CV risk [10]. Vegetable fats are recommended [19]. Patients are advised to increase their consumption of foods rich in long-chain omega-3 fatty acids from fatty fish and omega-3 linolenic acid. Evidence indicates that high-monounsaturated fatty acid (MUFA) diets are associated with improved glycemic control and CV disease risk or risk factors [31]. None of the menus contained seeds, nuts or fatty fish, but poultry, meat and butter were included every day in most of the meals. Given this distribution of fat sources, meeting recommendations and achieving a well-balanced diet is unfeasible.

The recommendations for protein intake for patients with diabetes align with those for the general population [12, 19]. All the examined hospitals met the stipulated protein requirements. However, the distribution of protein sources was

not optimal. Animal protein predominated, surpassing plant protein by approximately 1.5 times. The prevalence of animal protein contributes to an elevated intake of saturated fatty acids, with six out of the seven surveyed hospitals falling short of the recommended limit. A closer examination of meal compositions revealed minimal inclusion of plant proteins, with legumes present in trace amounts and the absence of nuts and seeds. Legumes share several characteristics with whole grains that could potentially benefit glycemic control, including the presence of slow-release carbohydrates and a high fiber content [21]. According to the recommendations, diabetic patients should receive 4–6 meals with a 3–4 h break between them [17]. Three out of seven examined hospitals offered only three meals a day, which was insufficient and could lead to deterioration of glycemic control and increased risk of hypoglycemia. According to some interpreters, incorrect nutrition is inconsistent with current medical knowledge and should be considered a violation of the Act on Patient Rights and the Act on the Patient Ombudsman [32]. Medical entities – hospitals and clinics – are responsible for providing the proper diets. Given all the mentioned deficiencies in hospital nutrition, it is necessary to consider whether and how state authorities should oversee its quality. New legal regulations are also needed. The National Health Fund, as the authorized body, would then have a stronger basis and duty for assessing the quality of hospital nutrition and its compliance with the requirements.

Limitations: The largest limitation of the study is the relatively small number of examined hospitals, which may affect its representativeness.

Nonetheless, despite the sample size, the results underscore the severity of the issue from a specific viewpoint. We analyzed 70 menus, which were received by a considerable number of patients. Each hospital's capacity, based on its province and annual occupancy rate, allows us to estimate the total number of patient beds and hospitalization days. For example, hospital 1 has 1,035 beds, hospital 2 has 452 beds, hospital 3 has 683 beds, hospital 4 has 677 beds, hospital 5 has 363 beds, hospital 6 has 176 beds, and hospital 7 has 321 beds. Summing these data helps us determine the number of patients missing out on nutritional education, impacting the healthcare system negatively. Hospitals 1, 2, and 3 are in the Mazowieckie Voivodeship, with an average of 40.8 patients per bed per year, resulting in  $(2,170 \times 40.8)$  88,536 patients. Hospitals 4, 6, and 7 are in the Śląskie Voivodeship, with an average of 36.2 patients per bed per year, resulting in  $(1,174 \times 36.2)$  42,498.8 patients. Hospital 5 is in the Łódzkie Voivodeship, with an average of 42.4 patients per bed per year, resulting in approximately 15,391 patients annually. Adding these numbers, we get an estimated total of 146,425.8 patients. The limited number of hospitals results from the scarce availability of data, complicated access to the person responsible for hospital nutrition and not providing us with necessary data (such as portion sizes, preparation methods, and specific diets). Therefore more research on the topic is needed. Another limitation arises from the operation of the Dieta 6 program, which uses approximations to estimate nutritional values. The program also includes pre-prepared standardized dishes, such as soups, which may differ from the actual meals and slightly impact the values of the meals.

In conclusion, our current data revealed that the so-called "diabetic" diets failed to meet patient requirements. This suggests an insufficient level of public awareness on the subject [21]. Insufficient nutrition education during hospitalization may lead to potential post-discharge dietary errors leading to deterioration in glycemic regulation and lipid profiles, and heightened susceptibility to complications. In the situation where doctors and nurses are often overburdened with work and there is a small number of dietitians on the wards, meals received during a hospital stay are of particular importance – sometimes they are the only form of education. To address these challenges effectively, it is essential to implement policy reforms, initiate extensive educational campaigns, and establish regular and binding quality controls for hospital nutrition. Additionally, more research on this topic is needed to exert pressure on decision-making bodies such as the National Health Fund and the Ministry of Health. By recognizing and actively addressing the disparities un-

covered in this research, we can improve the care of diabetic patients and ultimately enhance their overall health. This preliminary study not only draws attention to present challenges but also establishes a basis for promising future research opportunities.

### Funding

No external funding.

### Ethical approval

This study was conducted according to the guidelines of the Declaration of Helsinki. Although conducting survey studies in Poland does not require bioethical approval, it was acquired retrospectively for publication purposes by the Institutional Review of the Bioethical Committee at the Medical University of Warsaw (date: 11 September 2023, number: AKBE/258/2023).

### Conflict of interest

The authors declare no conflict of interest.

### References

1. International Diabetes Federation. IDF Diabetes Atlas 2021- 10<sup>th</sup> edition. Published 2021. Accessed November 21, 2023. <https://diabetesatlas.org/>
2. Ministry of Health. Diabetes. Published December 14, 2022. Accessed November 21, 2023. <https://archiwum.mz.gov.pl/zdrowie-i-profilaktyka/choroby-cywilizacyjne/cukrzyca/>
3. de Lorgeril M, Salen P, Martin JL, Monjaud I, Delaye J, Marmelle N. Mediterranean diet, traditional risk factors, and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon Diet Heart Study. *Circulation* 1999; 99: 779-85.
4. McDonagh TA, Metra M, Adamo M, et al. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure [published correction appears in *Eur Heart J* 2021; 42: 4901]. *Eur Heart J* 2021; 42: 3599-726.
5. Diabetes in America. 3<sup>rd</sup> edn. Ley SH, Schulze MB, Hivert MF, et al. (eds.) National Institute of Diabetes and Digestive and Kidney Diseases (US), Bethesda (MD) 2018.
6. Ley SH, Hamdy O, Mohan V, Hu FB. Prevention and management of type 2 diabetes: dietary components and nutritional strategies. *Lancet* 2014; 383: 1999-2007.
7. Cicero AFG, Fogacci F. The year in nutrition medicine 2023. *Arch Med Sci* 2023; 19: 1599-1601.
8. Liu D, Shi Q, Cheng G, Huang Q, Li S. Worldwide burden attributable to diet high in red meat from 1990 to 2019. *Arch Med Sci* 2022; 19: 1-15.
9. Liese AD, Nichols M, Sun X, D'Agostino RB Jr, Haffner SM. Adherence to the DASH Diet is inversely associated with incidence of type 2 diabetes: the insulin resistance atherosclerosis study. *Diabetes Care* 2009; 32: 1434-6.
10. Salas-Salvadó J, Bulló M, Estruch R, et al. Prevention of diabetes with Mediterranean diets: a subgroup analysis of a randomized trial [published correction appears in *Ann Intern Med* 2018; 169: 271-2]. *Ann Intern Med* 2014; 160: 1-10.

11. Tonstad S, Stewart K, Oda K, Batech M, Herring RP, Fraser GE. Vegetarian diets and incidence of diabetes in the Adventist Health Study-2. *Nutr Metab Cardiovasc Dis* 2013; 23: 292-9.
12. Araszkiwicz A, Bandurska-Stankiewicz E, Borys S, et al. 2023 Guidelines on the management of patients with diabetes – a position of Diabetes Poland. *Curr Topics Diabetes* 2023; 3: 1-133.
13. Powers MA, Bardsley J, Cypress M, et al. Diabetes self-management education and support in type 2 diabetes: a joint position statement of the American Diabetes Association, the American Association of Diabetes Educators, and the Academy of Nutrition and Dietetics. *J Acad Nutr Diet* 2015; 115: 1323-34.
14. NIK – The Supreme Audit Office Nutrition of Patients in Hospitals. Published April 13, 2018. Accessed November 10, 2023. <https://www.nik.gov.pl/aktualnosci/nik-o-zywieniu-w-szpitalach.html>.
15. prof. MUDr. Roman Havlík, Ph.D., ředitel Fakultní nemocnice Olomouc. Projekt Nemocniční strava. Published February 27, 2019. Accessed June 29, 2024.: <https://www.nemocnicnistrava.cz/pdf/projekt-Nemocnicni-strava.pdf>
16. Dr. Kristin Hünninghaus, Prof. Dr. Gustav Dobos, Dr. Markus Keller, et al. Offener Brief: Ernährung im Krankenhaus – wichtig für Gesundheit und Umwelt. Accessed June 29, 2024. <https://pan-int.org/de/ernaehrung-im-krankenhaus-offener-brief/>
17. Ministry of Health. Good meal in the hospital. Published September 27, 2023. Accessed November 21, 2023. <https://www.gov.pl/web/zdrowie/dobry-posilek-w-szpitalu>
18. Sińska B, Kucharska A. Dietary guidelines in diabetes – why are they so difficult to follow? Zalecenia żywieniowe w cukrzycy – dlaczego tak trudno ich przestrzegać? *Pediatr Endocrinol Diabetes Metab* 2023; 29: 125-7.
19. ElSayed NA, Aleppo G, Aroda VR, et al. Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes: Standards of Care in Diabetes-2023. *Diabetes Care* 2023; 46 (Suppl 1): S68-96.
20. Asche C, LaFleur J, Conner C. A review of diabetes treatment adherence and the association with clinical and economic outcomes. *Clin Ther* 2011; 33: 74-109.
21. Czapla M, Juárez-Vela R, Łokieć K, Wleklík M, Karniej P, Smereka J. The association between nutritional status and length of hospital stay among patients with hypertension. *Int J Environ Res Public Health* 2022; 19: 5827.
22. Zaremba N, Watson A, Kan C, et al. Multidisciplinary healthcare teams' challenges and strategies in supporting people with type 1 diabetes to recover from disordered eating. *Diabet Med* 2020; 37: 1992-2000.
23. American Diabetes Association Professional Practice Committee. 3. Prevention or Delay of Diabetes and Associated Comorbidities: Standards of Care in Diabetes-2024. *Diabetes Care* 2024; 47 (Suppl 1): S43-51.
24. Beverly EA, Ganda OP, Ritholz MD, et al. Look who's (not) talking: diabetic patients' willingness to discuss self-care with physicians. *Diabetes Care* 2012; 35: 1466-72.
25. Heisler M, Cole I, Weir D, Kerr EA, Hayward RA. Does physician communication influence older patients' diabetes self-management and glycemic control? Results from the Health and Retirement Study (HRS). *J Gerontol A Biol Sci Med Sci* 2007; 62: 1435-42.
26. National Center for Nutrition Education. National survey of the diet and nutritional status of the Polish population – report 2021. Published October 26, 2021. Accessed December 6, 2023. Available online: [https://ncez.pzh.gov.pl/sdm\\_downloads/20467/](https://ncez.pzh.gov.pl/sdm_downloads/20467/)
27. Evert AB, Boucher JL, Cypress M, et al. Nutrition therapy recommendations for the management of adults with diabetes. *Diabetes Care* 2014; 37 Suppl 1: S120-43.
28. He M, van Dam RM, Rimm E, Hu FB, Qi L. Whole-grain, cereal fiber, bran, and germ intake and the risks of all-cause and cardiovascular disease-specific mortality among women with type 2 diabetes mellitus. *Circulation* 2010; 121: 2162-8.
29. Burger KN, Beulens JW, van der Schouw YT, et al. Dietary fiber, carbohydrate quality and quantity, and mortality risk of individuals with diabetes mellitus. *PLoS One* 2012; 7: e43127.
30. Zafar MI, Mills KE, Zheng J, et al. Low-glycemic index diets as an intervention for diabetes: a systematic review and meta-analysis. *Am J Clin Nutr* 2019; 110: 891-902.
31. Schwingshackl L, Strasser B, Hoffmann G. Effects of monounsaturated fatty acids on glycaemic control in patients with abnormal glucose metabolism: a systematic review and meta-analysis. *Ann Nutr Metab* 2011; 58: 290-6.
32. Rożdżeński W. improper nutrition as a basis of responsibility of a medical entity. *Med Law Quart* 2019; 2: 71-98. <https://przegladprawamedycznego.wum.edu.pl/index.php/ppm/article/view/95>.