

Effect of single-bundle autologous hamstring tendon reconstruction combined with braided thread on joint stability and clinical efficacy in patients with posterior cruciate ligament rupture

Keywords

clinical efficacy, Autologous hamstring tendon single-bundle reconstruction, braided thread, posterior cruciate ligament rupture, joint stability

Abstract

Introduction

Background: Single-bundle autologous hamstring tendon reconstruction is a surgical procedure used primarily in orthopedics. To investigate the impact of using autologous hamstring tendon single-bundle restoration in conjunction with braided threads on both joint stability and clinical efficacy in patients suffering from posterior cruciate ligament (PCL) rupture. There are relatively few studies on the diagnosis and treatment of posterior cruciate ligament.

Material and methods

methods: Total of 106 patients diagnosed with PCL rupture were randomly assigned to the control group and study group, and each group consisting of 53 instances. The control group received autologous hamstring tendon single-bundle reconstruction, whereas the study group received autologous hamstring tendon single-bundle reconstruction together with braided thread treatment. The comparative rates of treatment success and satisfaction, complication occurrence, pre-and post-surgery joint activity indicators, gait parameters, knee joint function, and joint stability are assessed.

Results

Results: The study group showed a significantly higher rate of excellent and good treatment outcomes compared to the control group, with a statistical significance of $P < 0.05$. After 12 months of surgery, study group showed significantly higher joint activity index, stride length, stride speed, and Rasmussen score compared to the control group.

Conclusions

conclusion: The single-bundle reconstruction combined with braided thread has good clinical efficacy in patients with PCL rupture and more effectively improve the patient's joint stability.

Effect of single-bundle autologous hamstring tendon reconstruction combined with braided thread on joint stability and clinical efficacy in patients with posterior cruciate ligament rupture

Abstract

Background: Single-bundle autologous hamstring tendon reconstruction is a surgical procedure used primarily in orthopedics. **Objective:** To investigate the impact of using autologous hamstring tendon single-bundle restoration in conjunction with braided threads on both joint stability and clinical efficacy in patients suffering from posterior cruciate ligament (PCL) rupture. **Methods:** Total of 106 patients diagnosed with PCL rupture were randomly assigned to the control group and study group, and each group consisting of 53 instances. The control group received autologous hamstring tendon single-bundle reconstruction, whereas the study group received autologous hamstring tendon single-bundle reconstruction together with braided thread treatment. The comparative rates of treatment success and satisfaction, complication occurrence, pre-and post-surgery joint activity indicators, gait parameters, knee joint function, and joint stability are assessed. **Results:** The study group showed a significantly higher rate of excellent and good treatment outcomes compared to the control group, with a statistical significance of $P < 0.05$. After 12 months of surgery, study group showed significantly higher joint activity index, stride length, stride speed, and Rasmussen score compared to the control group. **Conclusion:** The single-bundle reconstruction combined with braided thread has good clinical efficacy in patients with PCL rupture and more effectively improve the patient's joint stability.

Key words: Autologous hamstring tendon single-bundle reconstruction; braided thread;

posterior cruciate ligament rupture; joint stability; clinical efficacy

Background

Cruciate ligament rupture is a common injury around knee joint. Among these type of injuries, anterior cruciate ligament rupture accounts for a high proportion and the incidence of posterior cruciate ligament is relatively low. Therefore, there are relatively few studies on the diagnosis and treatment of posterior cruciate ligament. As the posterior cruciate ligament is an important structure in maintaining the stability of the rear of the knee joint, there is a high demand for its diagnosis and treatment. Surgical treatment is an effective method for cruciate ligament rupture. The effects of different surgical methods vary significantly, and their impact on joint stability also varies [1,2]. Recently, there are an increasing number of clinical studies using single-bundle autologous hamstring tendon reconstruction to treat this type of injury, and most studies have given high recognition to its effectiveness. However, research also generally shows that its early treatment effect on posterior cruciate ligament is better and its stability is acceptable, but as time goes by, its stability is affected. Therefore, finding a treatment method that can effectively control graft stability has become the focus of clinical research [3-5]. As a non-absorbable suture with increasing application rate in orthopedic surgeries, braided sutures have obvious advantages in terms of tissue compatibility and strength. However, there are relatively few studies on its application in patients with posterior cruciate ligament autologous hamstring tendon single-bundle reconstruction, including an extremely lack of research on the impact on patients' joint stability. Therefore, this study now explores the effect of autologous hamstring tendon single-bundle reconstruction combined with braided threads on the joint stability and clinical efficacy of patients with posterior

cruciate ligament rupture to provide a reference for the selection and formulation of treatment methods for patients with this type of surgery. The objective of this study is to examine the effects of utilizing autologous hamstring tendon single-bundle restoration in combination with braided threads on both joint stability and clinical efficacy in patients with posterior cruciate ligament rupture.

Materials and Methods

General Data

Based on the preliminary experiment and the calculation formula, the number of participants in this trial was determined to be 106 cases. Therefore, 106 patients with posterior cruciate ligament rupture during the period from December 2020 to June 2022 were divided into a control group of 53 cases and an observation group of 53 cases according to the random number table approach. **Inclusion Criteria:** Patients of 20-65 years age; patients with posterior cruciate ligament rupture confirmed by drawer test and MRI; patients who meet the indications for surgery; patients, who signed the informed consent form for this study. **Exclusion Criteria:** patients with other ligament injuries, degenerative disease, history of knee joint surgery; bilateral injuries; chronic diseases; pregnancy and lactation are excluded from this study.

Method

Treatment methods

The control group underwent autologous hamstring tendon single-bundle reconstruction treatment. Combined epidural anesthesia was performed with double Endo-Button technique after routine preoperative examination. Routinely establish an anterior medial and lateral approach to the knee, insert an arthroscope, and conduct detailed exploration of the lesion and

surrounding conditions; those with meniscal injuries should be repaired first. Take the patient's autogenous hamstring tendon, braid and suture both ends of the tendon, and fold it into four strands (each with a diameter of 9 mm). Establish tibial and femoral channels, paying attention to the location of the channels. The tibial tunnel uses a reverse drill to drill a bone channel of appropriate length. Then the tendon is first introduced into the femoral channel under the guidance of the steel wire. Then guide the other end of the tendon into the tibial tunnel under the guidance of the wire. With the knee extended, the tendons are effectively tightened. The study group underwent autologous hamstring tendon single-bundle reconstruction combined with braided thread treatment. It adopts double Endo-Button technology and uses Ethicon thread to knit with a flat knot (as shown in Figure 1). Braid to the desired length, wrap and suture the tendon around the braided wire (the total diameter of the graft reaches 9 mm) (Fig. 2). MR During operation and before and after operation are shown in figures 3-5.



FIG. 1 Braid method



FIG. 2 Tendon wrapping suture method



FIG. 3 During operation




FIG. 4 MR before operation

FIG. 5 MR after operation

Postoperative Rehabilitation

During the first week after the operation, the focus is on isometric contractions of the quadriceps femoris, with small-range active flexion and extension activities within the painless range. Starting from the second week, get out of bed wearing a brace and walk with partial weight-bearing while using crutches. Under the non-weight-bearing condition, gradually strengthen the joint flexion and extension function exercises until the full range of motion is achieved. Continue with muscle strength training. The brace and crutches are removed three months after the operation, and normal walking is allowed. Gradually resume low- to moderate-intensity exercises six to seven months after the operation.

Observation Indicators

Assess the comparative rates of successful and satisfactory outcomes in treatment, the occurrence of complications, indicators of joint activity both pre- and post-surgery (including range of motion and maximum flexion), parameters related to walking (such as stride length, walking speed, and gait asymmetry index), knee joint function (Rasmussen score), and joint stability (as determined by the results of the KT2000 test). (1) Effect of treatment: The patients' treatment effects were assessed based on the knee joint function scoring criteria. The evaluation consists of six positive aspects and one negative aspect, with a total score of 100 points. Scores of ≥ 85 , 70-84, 60-69, and ≤ 59 correspond to excellent, good, average, and bad performance, respectively [6]. The treatment satisfaction rate is computed. (2) Complication incidence rate: The occurrence rates of complications, including infection, vascular and nerve injury, and flexion limitation, were computed for both groups. (3) Joint activity index: The joint activity of both groups was assessed prior to surgery and 12 months post-surgery, measuring the range of motion and maximum flexion of the knee joint. These specific indicators were determined using plain radiograph assessment. (4) Gait parameters: A three-dimensional gait analysis system was used to identify and examine the gait characteristics of both patient groups before to surgery and 12 months post-surgery. These parameters include stride length, walking speed, and gait asymmetry index. (5) Knee function: The knee joint function of the two groups was evaluated according to Rasmussen score before surgery and 12 months after surgery. It includes five aspects of assessment including pain, walking ability, knee extension, joint mobility and joint stability. The maximum score for each aspect is 6 points. The higher the score, the better the functional status of the knee joint. (6) Joint stability index: The KT2000 arthrometer was used to

detect the tibial posterior translation of the two groups at 15 pounds, 20 pounds and 30 pounds before surgery and 12 months after surgery, including forward and rear displacement, the average value is measured three times [7].

Statistical analysis

The data testing was conducted using SPSS 23.0 programme. The count data were presented using the notation [n (%)]. The chi-square test was employed to make comparisons between the groups. The measurement data, which followed a normal distribution, were described using the notation ($\bar{x} \pm s$). Group comparisons were conducted using independent sample t-tests. A statistically significant difference is indicated by a P-value of < 0.05 .

Results

Comparison of data of two groups

The two groups did not demonstrate any significant differences in terms of gender, age, BMI value, course of disease, lesion site, and cause of injury ($P > 0.05$) (Table 1).

Table 1 Comparison of general data of the two groups

Group	Cases	Gender [n(%)]		Age (Years Old)	BMI value (kg/m ²)	Course of disease (Month)	Lesion site [n(%)]		Cause of injury [n(%)]	
		Male	Female				Left	Right	Car accident	Movement
Control	53	35 (66.04)	18 (33.96)	35.63±10.2 6	25.12±2.06	5.32±1.91	33 (62.26)	20 (37.74)	35 (66.04)	18 (33.96)
Study	53	37 (69.81)	16 (30.19)	35.39±11.0 2	25.09±2.11	5.26±1.79	34 (64.15)	19 (35.85)	33 (62.26)	20 (37.74)
χ^2/t value		0.173		0.116	0.074	0.166	0.041		0.164	
P value		0.677		0.907	0.971	0.867	0.840		0.685	

Comparison of overall treatment of two groups

The treatment rate in the study group was significantly higher than that in the control group, as indicated by the statistical analysis ($P < 0.05$) presented in Table 2.

Table 2 Comparison of overall treatment of two groups [n(%)]

Group	Cases	Excellent	Good	Medium	Poor	Overall excellent and good rate
Control	53	28 (52.83)	17 (32.08)	7 (13.21)	1 (1.89)	45 (84.91)
Study	53	35 (66.04)	17 (32.08)	1 (1.89)	0 (0.00)	52 (98.11)
χ^2 value		---	---	---	---	5.950
P value		---	---	---	---	0.015

Comparison of complication rates of the two groups

Statistically, there was no significant difference in the incidence of complications between the two groups ($P>0.05$), see Table 3.

Table 3 Comparison of complication rates between the two groups [n(%)]

Group	Cases	Infection	Vascular nerve damage	Limited flexion
Control	53	0 (0.00)	1 (1.89)	2 (3.77)
Study	53	1 (3.77)	0 (0.00)	0 (0.00)
χ^2 value		1.010	1.010	2.038
P value		0.315	0.315	0.153

Comparison of joint activity indicators of the two groups before and after surgery

The preoperative difference in joint activity indices between the two groups was not statistically significant ($P>0.05$). However, 12 months following surgery, the study group exhibited significantly higher joint activity indexes compared to the control group ($P<0.05$) (Table 4).

Table 4 Comparison of joint activity indicators before and after surgery between the two groups

Group	Cases	Knee Range Of Motion		Maximum Knee Flexion	
		Before surgery	12 months after surgery	Before surgery	12 months after surgery
Control	53	85.69±2.96	105.72±3.19	90.03±2.63	106.19±3.06
Study	53	85.75±3.03	108.69±3.36	89.96±2.66	109.35±3.32
T value		0.103	4.666	0.136	5.095
P value		0.918	<0.001	0.891	<0.001

Comparison of gait parameters of two groups before and after surgery

When analyzing the gait metrics of the two groups before surgery, there was no statistically

significant difference detected ($P>0.05$). After 12 months of surgery, the study group demonstrated a substantial increase in both stride length and walking speed, in comparison to the control group. Furthermore, the study group had a significantly decreased gait asymmetry index compared to the control group. The control group exhibited a statistically significant difference at a significance level of $P<0.05$, as indicated in Table 5.

Table 5 Comparison of gait parameters between the two groups before and after surgery

Group	Cases	Stride (m)		Pace (m/s)		Gait asymmetry index	
		Before surgery	12 months after surgery	Before surgery	12 months after surgery	Before surgery	12 months after surgery
Control	53	0.63±0.10	1.16±0.11	1.01±0.12	1.26±0.13	0.32±0.06	0.09±0.03
Study	53	0.65±0.09	1.22±0.13	0.99±0.10	1.33±0.15	0.31±0.07	0.06±0.02
T value		1.082	2.565	0.932	2.567	0.789	6.057
P value		0.281	0.011	0.353	0.011	0.431	<0.001

Comparison of Rasmussen scores between the two groups before and after surgery

The difference in Rasmussen scores between the two groups before surgery did not reach statistical significance ($P>0.05$). The Rasmussen score of the study group 12 months' post-surgery exhibited a notable increase compared to the control group, and this disparity was statistically significant ($P<0.05$). Refer to table 6.

Table 6 Comparison of Rasmussen scores between the two groups before and after surgery (points)

Group	Cases	Pain		Walking ability		Knee extension		Joint range of motion		Joint stability	
		Before surgery	12 months after surgery	Before surgery	12 months after surgery	Before surgery	12 months after surgery	Before surgery	12 months after surgery	Before surgery	12 months after surgery
Control	53	3.10±0	5.50±0.26	2.71±0.53	5.21±0.35	2.90±0.36	5.31±0.32	2.73±0.35	4.35±0.41	2.03±0.35	5.11±0.29
		.61									
Study	53	3.06±0	5.66±0.29	2.67±0.51	5.39±0.36	2.97±0.33	5.49±0.35	2.75±0.36	4.63±0.50	1.93±0.36	5.32±0.35
		.59									
T value		0.343	2.990	0.395	2.609	1.043	2.763	0.289	3.152	1.449	3.363
P value		0.732	0.003	0.692	0.010	0.299	0.006	0.772	0.002	0.150	0.001

Comparison of KT2000 test results between the two groups before and after surgery

The comparison of the KT2000 test results between the two groups before to surgery did not show a statistically significant difference ($P>0.05$). The KT2000 test results of the study group 12 months post-surgery were markedly lower compared to the control group, and this difference was statistically significant ($P<0.05$). Refer to Table 7 and Table 8.

Table 7 Comparison of forward displacement between the two groups before and after surgery (mm)

Group	Cases	15 pounds		20 pounds		30 pounds	
		Before surgery	12 months after surgery	Before surgery	12 months after surgery	Before surgery	12 months after surgery
Control	53	3.35±0.61	2.59±0.53	6.65±1.93	3.59±0.73	9.13±2.26	4.56±1.13
Study	53	3.50±0.59	2.36±0.50	6.59±2.01	3.30±0.65	9.25±2.31	4.32±1.01
T value		1.286	2.298	0.156	2.159	0.270	1.152
P value		0.201	0.023	0.875	0.033	0.787	0.251

Table 8 Comparison of posterior displacement between the two groups before and after surgery (mm)

Group	Cases	15 pounds		20 pounds		30 pounds	
		Before surgery	12 months after surgery	Before surgery	12 months after surgery	Before surgery	12 months after surgery
Control	53	0.75±0.10	0.70±0.06	1.96±0.25	1.35±0.15	3.39±0.39	1.41±0.28
Study	53	0.77±0.11	0.66±0.05	2.01±0.26	1.20±0.13	3.38±0.36	1.30±0.26
T value		0.979	3.728	1.009	5.501	0.137	2.095
P value		0.329	<0.001	0.315	<0.001	0.891	0.038

Discussion

Posterior cruciate ligament rupture can cause local pain, swelling and joint dysfunction, seriously affecting the patient's quality of life. While the occurrence of cruciate ligament rupture is less common compared to anterior cruciate ligament rupture, it is nevertheless a major injury that should not be taken lightly. This type of injury requires accurate clinical diagnosis and appropriate treatment. The primary emphasis of research on the posterior cruciate ligament is the investigation of the impact of surgical interventions. Historically, a

significant number of clinical investigations have employed autologous hamstring tendon single-bundle restoration as a treatment for posterior cruciate ligament rupture. These studies have also confirmed its clinical effectiveness [8,9]. Nevertheless, additional research is required to fully investigate the enduring consequences of this surgical procedure. Furthermore, there is room for improvement in various aspects, such as enhancing knee joint functionality and sustaining joint stability. Specifically, the long-term maintenance of stability requires further improvement. Simultaneously, the stability of the joint is intricately linked to the gait parameters of patients with knee cruciate ligament rupture. consequently, there is an increased requirement for prolonged maintenance of knee joint stability in these individuals [10].

Tinga et al demonstrated that employing high-strength braided wires in animal studies can enhance and sustain joint stability [11]. Considering the presence of these circumstances, it is possible to enhance the autologous hamstring tendon graft in patients with posterior cruciate ligament rupture. This helps to manage the reduction of its looseness and has a positive impact on maintaining long-term joint stability. However, research in this area is still relatively inadequate. The results of this study show that the application effect of autologous hamstring tendon single-bundle reconstruction combined with braided thread is relatively effective in patients with posterior cruciate ligament rupture. The overall rate of therapy effectiveness is rather high. Simultaneously, the joint activity indicators, gait characteristics, Rasmussen score, and KT2000 test results one year after treatment exhibit comparatively superior outcomes in individuals with braided threads compared to those without. This shows that the application of braided wires effectively improves the long-term treatment effect of patients with posterior

cruciate ligament, and has a positive effect on improving the long-term stability and functional status of the joints. At the same time, their gait parameters are also effectively improved. McDonald et al. believed that the ligament-enhanced reconstruction system can help to shorten the recovery time during the reconstruction of isolated posterior cruciate ligament injuries, and has a relatively prominent effect on improving knee joint laxity [12]. In addition, Saragaglia et al. showed that the application of artificial ligament reinforcement system based on the application of autologous hamstring tendon graft is more helpful in improving the laxity in patients with posterior cruciate ligament, and is therefore more helpful to maintain long-term stability. All of the aforementioned factors indicate the need for further stabilization measures for patients who have had autologous hamstring tendon single-bundle repair of the posterior cruciate ligament [13]. Zhang Lei et al have shown that adding high-strength braided wires to autologous hamstring tendon grafts has a positive effect on improving joint function in patients with posterior cruciate ligament rupture [14]. Simultaneously, it also exerts a beneficial impact on enhancing the patient's joint stability after one year. Hence, the utilization of autologous hamstring tendon single-bundle restoration in conjunction with braided thread yields superior outcomes and greater significance for individuals suffering from posterior cruciate ligament rupture.

Upon examination, it has been seen that the use of braided wires in graft applications can cause the graft to collapse and then reconstruct during its proliferation stage. These occurrences can alter the mechanics of the graft, leading to greater looseness and a reduced capacity to withstand stress. The level of joint stability is quite low, as indicated by previous research [15,16]. **The advantages of autologous ligaments are the absence of rejection**

reactions and reliable strength in the later stage. The early strength is insufficient and ligament laxity is prone to occur. Therefore, the main purpose of adding braided suture is to enhance the overall strength of the implant in the early stage without affecting the diameter of the ligament. The high-strength suture is re-braided to have greater strength and an elasticity of 1 - 2 mm at the same time, which can effectively reduce the friction between the braided suture and the tendon. Consequently, the use of braided wires on the graft significantly enhances its ability to withstand stress and strengthens its stability. This is also a crucial factor in improving joint function and gait characteristics. **In addition, whether the single-bundle autologous hamstring tendon restoration is suitable for all ligament injuries remains to be verified by clinical experiments, but it is certain that the single-bundle autologous hamstring tendon restoration has a significant clinical effect on anterior cruciate ligament rupture. Xiaodong Bai's research shows that the one-stage reconstruction of anterior cruciate ligament rupture with autologous hamstring tendon transplantation can effectively restore the stability of the knee joint, and the early postoperative knee joint function is good [17].**

Conclusion

The results indicated that the utilization of single-bundle autologous hamstring tendon restoration in conjunction with braided threads yields superior clinical effectiveness in individuals with posterior cruciate ligament rupture, leading to a more substantial improvement in joint stability.

Ethical statement:

The patients, who signed agreement in the consent form and followed the declaration of the

Helsinki principle. This study was approved by the Zhoushan Dinghai Guanghua hospital medical ethics committee with Ethical Approval Number: 2024 unshan 11.

Conflict of interest

The authors declare that there is no conflict of interest in this work.

References


1. Botao Zhu, Peilong Dong, Xiaobo Tang, et al. Arthroscopic single-tunnel suture bundle fixation for the treatment of tibial insertion fractures of posterior cruciate ligament in adults. *Chinese Journal of Reparative and Reconstructive Surgery*. 2023; 27 (22) : 3503-3507.
2. Xu Liu, Yu Wen, Bin Li. Comparison of the efficacy of autologous hamstring tendon plus peroneus longus tendon anterior bundle and LARS ligament reconstruction of ruptured posterior cruciate ligament. *The Journal of Practical Medicine*. 2019; 354: 593-597.
3. Zhenzhong Yuan, Gangjian Tang, Yexiong Wu, et al. Comparative study of arthroscopic autologous peroneus longus tendon and hamstring tendon reconstruction of anterior cruciate ligament. *Chin J. Endosc.* 2021; 27 (11) : 83-88.
4. Ming Li, Yuqiang Liu, Shiji Gao, et al. Early clinical efficacy of new autologous semitendinosus strip graft reconstruction in the treatment of anterior cruciate ligament rupture. *Chin J. Trauma*. 2022; 38 (4) : 332-339.
5. Xiao Wang, Xu Han, Xiaotao Shi, et al. Early efficacy of posterior cruciate ligament reconstruction using a single bundle of four autogenous semitendinosus tendons and the anterior semitendinosus bundle of the peroneus longus tendon under arthroscopic surgery. *Chinese Journal of Reparative and Reconstructive Surgery*. 2021; 35 (5) : 556-561.
6. Guoning Zhang, You Wang. Evaluation of Knee Scoring Criteria. *Chin J Surg*. 2006; 44 (16) : 1141-1143.
7. Jiashen Shao. Clinical study on Zhang's homeopathic traction minimally invasive reducer in the treatment of posterolateral tibial plateau fractures. Master thesis. Harbin: Harbin Institute of Technology, 2019.
8. Liu H, Yin H, Yi Y, et al. Effects of different rehabilitation training on balance function

in stroke patients: a systematic review and network meta-analysis. *Arch Med Sci.* 2023; 19(6): 1671-1683.

9. Tengyun Yang, Zhujun Zheng, Yanlin Li, et al. Comparative study on the efficacy of posterior cruciate ligament reconstruction using arthroscopic posterior triangulation technique and anterior and posterior approaches. *Chinese Journal of Reparative and Reconstructive Surgery.* 2021; 35 (7) : 823-828.
10. Bingkun Li, Lei Zhang, Zhangrong Xia, et al. Analysis of the clinical efficacy of posterior cruciate ligament reconstruction with autogenous tendon under knee arthroscopy. *Chin J Clin Anat.* 2022; 40 (4) : 466-470.
11. Tinga S, Kim SE, Banks SA, et al. Femorotibial joint kinematics in nine dogs treated with lateral suture stabilization for complete cranial cruciate ligament rupture. *J Am Vet Med Assoc.* 2021; 258(5): 493-501.
12. McDonald LK, Cosic F, Joseph S. The use of the ligament augmentation and reconstruction system for posterior cruciate ligament reconstruction in isolated and multiligament knee injuries: A systematic review. *Knee.* 2021; 30 (6) : 322-336.
13. Saragaglia D, Francony F, Gaillot J, et al. Posterior cruciate ligament reconstruction for chronic lesions: clinical experience with hamstring versus ligament advanced reinforcement system as graft. *Int Orthop.* 2020; 44(1): 179-185.
14. Lei Zhang, Bin Li. A clinical comparative study on whether to add high-strength braided wires to posterior cruciate ligament grafts for single-bundle reconstruction of autologous hamstring tendons. *China Comprehensive Clinical.* 2020; 36 (1) : 71-75.
15. Wei Hu, Xianke Yan Effect of posterior cruciate ligament preserving knee prosthesis on gait and lower extremity venous return in varus knee arthroplasty. *Journal of Clinical Rehabilitative Tissue Engineering Research, J Clin Rehabil Tis Eng Res.* 2023; 27 (22) : 3503-3507.
16. Lu He, Xinyu Liao, Yanlin Li, et al. Arthroscopic repair of multiple ligament injuries after knee joint dislocation: a three-dimensional gait analysis. *Chin. J. Tissue Eng. Res.* 2022; 26 (26) : 4205-4210.
17. Xiaodong Bai, Yoating Wang, Qi Che, et al. One-stage reconstructing the anterior and

posterior cruciate ligaments injuries with autologous hamstring tendon using Tight Rope devices all inside. Chinese Journal of Orthopaedics, 2018,38(17):1045-1054.

Preprint



Braided method and suture the tendon around the braided wire

Preprint