

Research based on nucleotide polymorphism reveals the role of inflammatory cytokines in regulating the influence of blood metabolites on drug-related osteonecrosis

Yazhou Li^{1,2}, Yanbin Wu², Junming Zhang², Xianghui Mei¹, Jiazhou Wu², Zexian Liu², Tao Qian², Junli Wang², Zaodi Mi², Ying He², Xiaohan Sun², Aiyuan Wang², Jiang Peng^{1,2*}

¹Clinical Medical College, Hebei University of Engineering, Handan, China

²Institute of Orthopedics, The Fourth Medical Center of Chinese PLA General Hospital, Beijing Key Lab of Regenerative Medicine in Orthopedics, Key Laboratory of Musculoskeletal Trauma & War Injuries PLA, Beijing, China

Submitted: 22 November 2024; **Accepted:** 31 January 2025

Online publication: 1 April 2025

Arch Med Sci 2026; 22 (1): 488–495

DOI: <https://doi.org/10.5114/aoms/200695>

Copyright © 2025 Termedia & Banach

***Corresponding author:**

Prof. Jiang Peng
Institute of Orthopedics,
The Fourth Medical Center of
Chinese PLA General Hospital
No. 51 Fucheng Road
Beijing, China
E-mail:
pengjiang301@126.com

Abstract

Introduction: Osteonecrosis is a debilitating disease caused by impaired blood supply leading to bone tissue death, and drug-related osteonecrosis is a significant clinical issue. The role of inflammation and metabolic disorders in the pathogenesis of osteonecrosis has garnered widespread attention, but the exact causal relationships remain unclear. This study aims to explore the causal link between inflammatory cytokines and drug-related osteonecrosis, while also investigating how metabolites might mediate this relationship.

Material and methods: We employed two-sample Mendelian randomization (MR) analysis to examine the causal links between 91 inflammatory cytokines, 1,400 blood metabolites, and drug-related osteonecrosis. Single nucleotide polymorphisms (SNPs) associated with inflammatory cytokines and metabolites were used as instrumental variables (IVs) to assess their potential relationship with drug-related osteonecrosis risk. We further conducted mediation MR analysis to explore the role of metabolites in mediating the impact of inflammatory cytokines on drug-related osteonecrosis.

Results: MR analysis demonstrated notable causal relationships between four inflammatory cytokines and drug-related osteonecrosis. Specifically, interleukin-4 (IL-4) and C-X-C motif chemokine 6 (CXCL6) showed a negative correlation with the risk of drug-related osteonecrosis, while interleukin-6 (IL-6) and glial cell line-derived neurotrophic factor (GDNF) exhibited a positive correlation with the risk. Furthermore, mediation analysis revealed that IL-4 affects the development of drug-related osteonecrosis via blood metabolites. Key metabolites identified as significant mediators included mannitol/sorbitol levels, the mannose-to-mannitol-to-sorbitol ratio, and the glucose-to-mannitol-to-sorbitol ratio.

Conclusions: This study presents new evidence connecting inflammatory cytokines to drug-related osteonecrosis and highlights the mediating role of metabolites. These results help us understand the pathogenesis of the disease and provide new insights for its prevention and treatment.

Key words: nucleotide polymorphism, pro-inflammatory mediators, metabolites, drug-related osteonecrosis, Mendelian randomization.

Introduction

Osteonecrosis is a condition marked by the destruction of bone tissue caused by an inadequate blood supply to bone cells [1]. Factors such as trauma, medications, alcohol, or metabolic disorders can trigger an immune response that interferes with bone repair, ultimately resulting in osteonecrosis [2–5]. Although osteonecrosis primarily occurs in the femoral head, it can also involve other regions [6]. Once diagnosed, it leads to a high rate of disability, severely diminishing patients' quality of life and imposing a substantial burden on families and society [7].

Inflammation is a key factor in the development of steroid-induced osteonecrosis, particularly through the macrophage polarization process driven by tumor necrosis factor- α (TNF- α) [8, 9]. Dysregulated inflammation not only hinders bone repair but also aggravates tissue damage by producing reactive oxygen species and proteases, underscoring the complex interplay between inflammation and bone loss [10–12]. Furthermore, recent research has revealed links between osteonecrosis and various metabolic abnormalities, such as lipid metabolism disruptions, coagulation pathway irregularities, and alterations in linoleic acid metabolism, offering new perspectives for diagnosis and treatment [13, 14]. However, the precise interaction between inflammation and metabolic disturbances in the progression of osteonecrosis remains insufficiently understood.

In this study, we adopted a mediation Mendelian randomization (MR) framework to examine the potential role of metabolites in the causal pathway linking inflammatory cytokines to drug-related osteonecrosis. Our objective was to unravel the mechanistic pathways connecting inflammation, metabolism, and drug-related osteonecrosis. Additionally, we aimed to identify innovative targets and strategies for the prevention, early diagnosis, and treatment of this condition. The outcomes of this research may offer fresh insights into the underlying causes of drug-related osteonecrosis and support the development of precision medicine solutions for managing this challenging disease.

Material and methods

Study design

MR analysis relies on three fundamental assumptions: relevance, independence, and exclusivity [15]. This study was conducted in two stages. In the first stage, a two-sample MR method was applied, with inflammatory cytokines as the exposure and drug-related osteonecrosis as the outcome, to examine their causal relationship. Subsequently, reverse MR analysis was performed.

The second stage focused on evaluating the mediating role of metabolites in the causal pathway between inflammatory cytokines and drug-related osteonecrosis. Inflammatory cytokines identified as significant in the first stage were used as exposure variables. Initially, the causal link between these inflammatory cytokines and potential mediators was tested, followed by an evaluation of the causal association between these mediators and drug-related osteonecrosis (Figure 1).

Data sources

All the data utilized in this study were obtained from publicly available genome-wide association study (GWAS) datasets; thus, no new ethics review board approval was necessary. The genetic data for inflammatory cytokines were derived from an earlier GWAS that included 91 plasma inflammatory cytokines, involving 14,824 individuals of European descent [16]. The genetic data for blood metabolites, which encompass 1,091 serum metabolites and 309 metabolite ratios, were sourced from the most recent GWAS summary dataset, which contained whole-genome genotyping data from 8,096 participants [17]. The data on drug-related osteonecrosis were obtained from the FinnGen R11 GWAS summary dataset (GWAS ID: finn-b-OSTEON_DRUGS), which comprised 348 cases and 453,385 control samples, available at this link: <https://www.finngen.fi/en>. To minimize potential bias arising from population heterogeneity, we specifically restricted the genetic composition of the study cohort to individuals of European ancestry.

Selection of tool variables

In MR analysis, SNPs are chosen as instrumental variables due to their strong association with the exposure variable. To address the potential limitation of obtaining too few SNPs with a p -value




Figure 1. Flowchart of the MR analysis process

threshold of 5×10^{-8} , which may hinder further analysis, we relaxed the threshold to $p < 1 \times 10^{-5}$ [18, 19]. To ensure the selected SNPs are independent, a linkage disequilibrium test was performed using an R^2 threshold of 0.001 and a physical distance threshold of 1,000 kb, filtering out SNPs that exhibit linkage disequilibrium [20]. By applying these stringent criteria, we identified a robust set of instrumental variables that are strongly associated with blood metabolites or inflammatory factors and are mutually independent, minimizing the risk of violating the MR assumptions of independence and exclusivity. Furthermore, to validate the strength of the instrumental variables, we calculated the F -statistic for each SNP, excluding those with an F -statistic below 10 [21]. To enhance the reliability and accuracy of the analysis, we also excluded instrumental variables that could potentially affect the results through pleiotropy.

Statistical analysis

This study employed multiple MR approaches, including inverse variance weighted (IVW), MR-Egger regression, weighted median (WM), simple mode, and weighted mode methods. Given the robustness of the IVW approach for inferring causality, it was chosen as the primary method for estimating causal relationships [22]. A p -value below 0.05 was considered indicative of a significant causal association between the exposure and the outcome.

To ensure the robustness and reliability of the findings, several sensitivity analyses were performed. First, heterogeneity and pleiotropy in the

causal estimates were examined using Cochran's Q test and the MR-Egger intercept [23, 24]. Second, the MR-PRESSO method was employed to detect pleiotropic bias, with a p -value exceeding 0.05 indicating no substantial evidence of horizontal pleiotropy [24]. Additionally, a leave-one-out analysis was conducted to assess the influence of individual SNPs on the overall causal estimate [25].

To explore the causal pathways among blood inflammatory cytokines, metabolites, and drug-related osteonecrosis, a two-step MR analysis was conducted to evaluate whether metabolites mediate the relationship between inflammatory factors and osteonecrosis. Initially, a two-sample MR analysis was used to estimate the total causal effect of inflammatory factors on osteonecrosis, primarily relying on the IVW method while incorporating sensitivity analyses with MR-Egger regression and the WM approach [24]. Subsequently, two independent two-sample MR analyses were performed: the first estimated the causal relationship between inflammatory factors and metabolites, yielding the causal effect estimate β_1 [26], while the second assessed the relationship between metabolites and drug-related osteonecrosis, resulting in the causal effect estimate β_2 [26]. The mediating effects were calculated using the two-step MR approach as follows: mediation effect = $\beta_1 * \beta_2$. The direct effect was determined by subtracting the mediation effect from the total effect [26]. All MR analyses were implemented using R software (version 4.4.1) along with relevant packages such as "TwoSampleMR" and "MRInstruments."




Figure 2. Forest plot of MR results for inflammatory cytokines and drug-related osteonecrosis

Results

The causal relationship between inflammatory cytokines and drug-related osteonecrosis

Using the IVW method as the primary analytical tool, the MR results indicated a significant causal relationship between the genetically predicted levels of four specific inflammatory cytokines and the risk of drug-related osteonecrosis. Notably, IL-4 (OR = 0.562; 95% CI: 0.334–0.947; $p = 0.030$) and CXCL6 (OR = 0.696; 95% CI: 0.498–0.973; $p = 0.034$) were identified as protective factors against drug-related osteonecrosis, while IL-6 (OR = 2.212; 95% CI: 1.181–4.143; $p = 0.013$) and GDNF (OR = 1.561; 95% CI: 1.019–2.389; $p = 0.040$) were found to be risk factors (Figure 2). Further sensitivity analysis showed no evidence of heterogeneity or pleiotropic bias among these four factors (Supplementary Table SI). Additionally, leave-one-out analysis further confirmed the reliability and stability of the results (Supplementary Figure S1). Subsequently, a reverse MR analysis was conducted, with drug-related osteonecrosis as the exposure variable and these four inflammatory cytokines as the outcomes. The results of the IVW method in the reverse analysis showed no causal relationship between drug-related osteonecrosis and these inflammatory cytokines, suggesting the feasibility of further research (Supplementary Table SII).

The causal relationship between metabolites and drug-related osteonecrosis

To verify the robustness of the MR analysis results, we performed cross-validation using five statistical approaches: IVW, MR-Egger regression, WM, simple mode, and weighted mode. Metabolites considered significant had to meet the fol-

lowing inclusion criterion: the OR values from all five methods consistently being either greater than 1 or less than 1. Additionally, pleiotropy was assessed ($p > 0.05$) to rule out the possibility of pleiotropic bias. Ultimately, 16 metabolites showing the strongest associations with drug-related osteonecrosis were identified, including 8 individual metabolites and 8 metabolite ratios (Figure 3). Sensitivity analysis revealed no evidence of horizontal pleiotropy or heterogeneity in the results (Supplementary Table SIII).

The causal relationship between inflammatory cytokines and metabolites

To investigate the mechanisms driving the onset and progression of drug-related osteonecrosis, we performed a mediation analysis to uncover pathways through which metabolites mediate the effects of inflammatory cytokines. Specifically, we analyzed the causal links between IL-4 and three metabolites and their subsequent influence on drug-related osteonecrosis. Our findings demonstrated a significant positive causal relationship between IL-4 and mannitol/sorbitol levels (OR = 1.128, 95% CI: 1.013–1.256, $p = 0.026$), indicating that elevated IL-4 levels may lead to increased mannitol/sorbitol levels. Furthermore, mannitol/sorbitol levels were significantly negatively associated with the risk of drug-related osteonecrosis (OR = 0.609, 95% CI: 0.424–0.876, $p = 0.007$), suggesting that these levels mediate IL-4's protective effect on osteonecrosis.

We also identified a significant negative relationship between IL-4 and the mannose-to-mannitol-to-sorbitol ratio (OR = 0.881, 95% CI: 0.791–0.981, $p = 0.021$), while this ratio was positively associated with an increased risk of osteonecrosis (OR = 1.861, 95% CI: 1.172–2.953, $p = 0.008$). These results suggest that elevated IL-4 may lower the mannose-to-mannitol-to-sorbitol ratio, there-




Figure 3. Forest plot of MR analysis for metabolites most strongly associated with drug-related osteonecrosis

by reducing osteonecrosis risk. Similarly, a significant negative causal relationship was observed between IL-4 and the glucose-to-mannitol-to-sorbitol ratio (OR = 0.876, 95% CI: 0.787–0.974, $p = 0.015$). Conversely, this ratio was positively associated with the risk of drug-related osteonecrosis (OR = 1.666, 95% CI: 1.154–2.405, $p = 0.006$). These findings suggest that higher IL-4 levels may reduce the glucose-to-mannitol-to-sorbitol ratio, further mitigating osteonecrosis risk (Figure 4). Sensitivity analyses confirmed the absence of hor-

izontal pleiotropy and heterogeneity (Supplementary Table SIV). Finally, the mediation effects of the three metabolites – mannitol/sorbitol levels, the mannose-to-mannitol-to-sorbitol ratio, and the glucose-to-mannitol-to-sorbitol ratio – were assessed in the causal pathway between IL-4 and drug-related osteonecrosis. The estimated mediation effects were -0.0598 (95% CI: $[-0.1290, 0.0089]$), -0.0785 (95% CI: $[-0.1600, 0.0103]$), and -0.0674 (95% CI: $[-0.1400, 0.0054]$), respectively (Table I).




Figure 4. Forest plot of MR analysis of the causal effects of IL-4 on mannitol/sorbitol levels, mannose to mannitol to sorbitol ratio, and glucose to mannitol to sorbitol ratio in drug-related osteonecrosis

Table I. The mediation effect of IL-4 on drug-related osteonecrosis through mannitol/sorbitol levels, mannose to mannitol to sorbitol ratio, and glucose to mannitol to sorbitol ratio

Exposure	β_1	Mediation	β_2	Outcome	Mediation effect [95% CI]	Total effect
IL-4	0.1210	Mannitol/sorbitol levels	-0.4945	Drug-related osteonecrosis	-0.0598 (-0.1290, 0.0089)	-0.5748
	-0.1263	Mannose to mannitol to sorbitol ratio	0.6212		-0.0785 (-0.1600, 0.0103)	
	-0.1320	Glucose to mannitol to sorbitol ratio	0.5106		-0.0674 (-0.1400, 0.0054)	

Discussion

We explored the causal impact of inflammatory cytokines on drug-related osteonecrosis and examined possible mediators. The findings confirm that metabolites play a mediating role in the inflammatory cytokine-driven pathogenesis of drug-related osteonecrosis.

IL-4 is a versatile cytokine that plays a crucial role in regulating macrophage activity. It primarily promotes the polarization of macrophages to the M2 phenotype, which is vital in processes such as anti-inflammatory responses, tissue repair, and fibrosis. In particular, IL-4 functions by suppressing the expression of CD14 and inhibiting the release of pro-inflammatory cytokines such as IL-6 and tumor necrosis factor (TNF) [27]. We propose that in the context of drug-induced osteonecrosis, IL-4 may reduce tissue damage by limiting the secretion of key pro-inflammatory mediators. CXCL6, a member of the chemokine ligand family, has been found to be overexpressed in tissues affected by diabetic nephropathy. Research indicates that miR-20a targets CXCL6, thereby inhibiting the JAK/STAT3 pathway, promoting the proliferation of HK-2 cells treated with high glucose, and reducing both cell apoptosis and inflammation [28]. These findings have led us to further explore whether CXCL6 may similarly affect drug-related osteonecrosis through comparable molecular mechanisms and pathways.

Cytokines are pivotal in regulating immune responses, and their imbalance or excessive production is often linked to tissue damage and the progression of various diseases [29]. For instance, IL-6, a cytokine involved in inflammation, immune homeostasis, and bone metabolism regulation [30], demonstrates a minimal effect on bone remodeling under normal conditions. However, its expression significantly increases in pathological states, which may contribute to greater bone resorption, aggravated inflammation, and metabolic disturbances [31]. These findings highlight the intricate relationship between IL-6 and bone-related disorders. GDNF, a member of the transforming growth factor- β family, has been found to be elevated in degenerated intervertebral disc tissues, which have a pro-inflammatory microenvironment. This

cytokine may play a pivotal role in the onset and spread of discogenic pain [32]. Building on these findings, we propose that inflammatory factors might regulate GDNF expression in drug-induced osteonecrosis cells. Additionally, the upregulation of GDNF could potentially enhance the transmission of inflammatory pain sensitivity. However, further research is necessary to better understand the underlying pathological mechanisms.

Mannitol and sorbitol are produced through the metabolism of fructose, mannose, and galactose, while glucose is converted to sorbitol through glycolysis and gluconeogenesis, and further oxidized into fructose [33, 34]. Under high glucose conditions, sorbitol dehydrogenase plays a crucial role in converting sorbitol into fructose. These metabolites can then form advanced glycation end products, which trigger liver metabolic changes, enhance de novo lipogenesis, and influence blood lipid levels, increasing cardiovascular disease risk [35, 36]. Our findings suggest that the levels of mannitol/sorbitol and related metabolites mediate IL-4's role in osteonecrosis development. This is a significant discovery, as research on these metabolites is limited, and their mechanisms remain unclear. Future studies should explore how these metabolites mediate the interaction between inflammatory cytokines and drug-related osteonecrosis, offering new insights into disease mechanisms and therapeutic strategies.

This study investigates the causal link between inflammatory cytokines, metabolites, and the risk of drug-related osteonecrosis using MR analysis. However, there are limitations to consider. The GWAS data primarily come from European populations, which raises concerns about whether the findings can be generalized to other ethnic groups. The reliance on European cohorts may introduce population stratification bias, limiting the broader applicability of the results. Future research should include data from diverse ethnic groups to address this issue. Furthermore, while several metabolites linked to drug-related osteonecrosis have been identified, their precise role in disease development remains unclear, which limits the full interpretation of the study's findings.

In conclusion, this study used MR to investigate the causal relationship between inflammatory cytokines and drug-related osteonecrosis, as well as the mediating role of metabolites. Four inflammatory cytokines were identified as causally linked to drug-related osteonecrosis. Mediation analysis further revealed that three blood metabolites, regulated through the IL-4 pathway, affect the risk of this condition. These findings provide valuable insights for prevention and treatment strategies.

Acknowledgments

We sincerely acknowledge the researchers and participants of all the genome-wide association studies whose data we used for this research.

Funding

This study was supported by the Beijing Natural Science Foundation (L242042).

Ethical approval

Not applicable.

Conflict of interest

The authors declare no conflict of interest.

References

1. Petek D, Hannouche D, Suva D. Osteonecrosis of the femoral head: pathophysiology and current concepts of treatment. *EFORT Open Rev* 2019; 4: 85-97.
2. Zhu T, Cui Y, Zhang M, Zhao D, Liu G, Ding J. Engineered three-dimensional scaffolds for enhanced bone regeneration in osteonecrosis. *Bioact Mater* 2020; 5: 584-601.
3. Li Z, Yang B, Weng X, Tse G, Chan MTV, Wu WKK. Emerging roles of MicroRNAs in osteonecrosis of the femoral head. *Cell Prolif* 2018; 51: e12405.
4. Gupta A, Tripathi L, Pandey S, Dwivedi D. Biology of bone morphogenetic proteins in skeleton disease: osteonecrosis in sickle cell disease patients. *Curr Protein Pept Sci* 2022; 23: 264-70.
5. Li L, Zhao S, Leng Z, et al. Pathological mechanisms and related markers of steroid-induced osteonecrosis of the femoral head. *Ann Med* 2024; 56: 2416070.
6. Murphey MD, Foreman KL, Klassen-Fischer MK, Fox MG, Chung EM, Kransdorf MJ. From the radiologic pathology archives imaging of osteonecrosis: radiologic-pathologic correlation. *Radiographics* 2014; 34: 1003-28.
7. Chen H, Xu J, Lv Y, et al. Proanthocyanidins exert a neuroprotective effect via ROS/JNK signaling in MPTP-induced Parkinson's disease models in vitro and in vivo. *Mol Med Rep* 2018; 18: 4913-21.
8. Weinstein RS. Glucocorticoid-induced osteonecrosis. *Endocrine* 2012; 41: 183-90.
9. Wu X, Xu W, Feng X, et al. TNF- α mediated inflammatory macrophage polarization contributes to the pathogenesis of steroid-induced osteonecrosis in mice. *Int J Immunopathol Pharmacol* 2015; 28: 351-61.
10. Loi F, Córdova LA, Pajarinen J, Lin TH, Yao Z, Goodman SB. Inflammation, fracture and bone repair. *Bone* 2016; 6: 119-30.
11. Kolaczowska E, Kuberski P. Neutrophil recruitment and function in health and inflammation. *Nat Rev Immunol* 2013; 13: 159-75.
12. Zhu D, Yu H, Liu P, et al. Calycosin modulates inflammation via suppressing TLR4/NF- κ B pathway and promotes bone formation to ameliorate glucocorticoid-induced osteonecrosis of the femoral head in rat. *Hytother Res* 2021; 35: 2824-2835.
13. Yu X, Zhang S, Zhang B, Dai M. Relationship of idiopathic femoral head necrosis with blood lipid metabolism and coagulation function: a propensity score-based analysis. *Front Surg* 2023; 9: 938565.
14. Zhu W, Wang R, Yang Z, et al. GC-MS based comparative metabolomic analysis of human cancellous bone reveals the critical role of linoleic acid metabolism in femur head necrosis. *Metabolomics* 2023; 19: 86.
15. Emdin CA, Khera AV, Kathiresan S. Mendelian randomization. *JAMA* 2017; 318: 1925-6.
16. Zhao JH, Stacey D, Eriksson N, et al. Genetics of circulating inflammatory proteins identifies drivers of immune-mediated disease risk and therapeutic targets. *Nat Immunol* 2023; 24: 1540-51.
17. Chen Y, Lu T, Pettersson-Kymmer U, et al. Genomic atlas of the plasma metabolome prioritizes metabolites implicated in human diseases. *Nat Genet* 2023; 55: 44-53.
18. Orrù V, Steri M, Sidore C, et al. Complex genetic signatures in immune cells underlie autoimmunity and inform therapy. *Nat Genet* 2020; 52: 1036-45.
19. Chen J, Yu X, Wu X, Chai K, Wang S. Causal relationships between gut microbiota, immune cell, and non-small cell lung cancer: a two-step, two-sample Mendelian randomization study. *J Cancer* 2024; 15: 1890-7.
20. Jiazhen H, Bozhen F. CSF1 mediating 3-hydroxyhexanoate in endometrial cancer: a Mendelian randomization study. *Arch Med Sci* 2024; 20: 1731-5.
21. Burgess S, Thompson SG, CRP CHD Genetics Collaboration. Avoiding bias from weak instruments in Mendelian randomization studies. *Int J Epidemiol* 2011; 40: 755-64.
22. Burgess S, Butterworth A, Thompson SG. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genet Epidemiol* 2013; 37: 658-65.
23. Bowden J, Del Greco MF, Minelli C, Davey Smith G, Sheehan N, Thompson J. A framework for the investigation of pleiotropy in two-sample summary data Mendelian randomization. *Stat Med* 2017; 36: 1783-802.
24. Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet* 2018; 50: 693-8.
25. Sekula P, del Greco MF, Pattaro C, Köttgen A. Mendelian randomization as an approach to assess causality using observational data. *J Am Soc Nephrol* 2016; 27: 3253-65.
26. Carter AR, Sanderson E, Hammerton G, et al. Mendelian randomisation for mediation analysis: current methods and challenges for implementation. *Eur J Epidemiol* 2021; 36: 465-78.
27. Sokol CL, Barton GM, Farr AG, Medzhitov R. A mechanism for the initiation of allergen-induced T helper type 2 responses. *Nat Immunol* 2008; 9: 310-8.
28. Wang SZ, Zhang YL, Shi HB. Potential repressive impact of microRNA-20a on renal tubular damage in diabetic kidney disease by targeting C-X-C motif chemokine ligand 6. *Arch Med Res* 2021; 52: 58-68.
29. Xu L, Yu Y, Sang R, Li J, Ge B, Zhang X. Protective effects of taraxasterol against ethanol-induced liver injury by regulating CYP2E1/Nrf2/HO-1 and NF- κ B signaling

- pathways in mice. *Oxid Med Cell Longev* 2018; 2018: 8284107.
30. Lorenzo JA. The role of interleukin-6 in bone. *J Endocr Soc* 2020; 4: bvaa112.
 31. Sims NA. Cell-specific paracrine actions of IL-6 family cytokines from bone, marrow and muscle that control bone formation and resorption. *Int J Biochem Cell Biol* 2016; 79: 14-23.
 32. Yamada J, Akeda K, Sano T, Iwasaki T, Takegami N, Sudo A. Expression of glial cell line-derived neurotrophic factor in the human intervertebral disc. *Spine* 2020; 45: E768-75.
 33. Brownlee M. Biochemistry and molecular cell biology of diabetic complications. *Nature* 2001; 414: 813-20.
 34. Tran DH, Wang ZV. Glucose metabolism in cardiac hypertrophy and heart failure. *J Am Heart Assoc* 2019; 8: e012673.
 35. Kizer JR, Benkeser D, Arnold AM, et al. Advanced glycation/glycoxidation endproduct carboxymethyl-lysine and incidence of coronary heart disease and stroke in older adults. *Atherosclerosis* 2014; 235: 116-21.
 36. Lamprea-Montealegre JA, Arnold AM, McClelland R, et al. Plasma levels of advanced glycation endproducts and risk of cardiovascular events: findings from 2 prospective cohorts. *J Am Heart Assoc* 2022; 11: e024012.