

# The Importance of TNF- $\alpha$ Signaling – Potential Risk Factor in Neurodegenerative and Cardiovascular Diseases

---

## Keywords

dementia, cytokines, plaque, Parkinson's disease, central nervous system, TNF- $\alpha$ , neuroinflammation, coronary artery disease, multiple sclerosis, intima-media

---

## Abstract

Cytokines are small and membrane-bound protein-based cell signaling molecules that aid cell-to-cell communication in physiological and pathological processes. TNF- $\alpha$  is a pleiotropic cytokine, it is regarded as a principal cytokine in the acute and chronic phases of the immune and inflammatory response. TNF- $\alpha$  has been demonstrated to play a role in neuroplasticity and myelination, as well as excitotoxicity, neuroinflammation, and damage to the blood-brain barrier. The involvement of TNF- $\alpha$  in the etiology of Parkinson's disease, both in its early and late stages, Alzheimer's disease, Multiple sclerosis, coronary artery disease has been substantiated using both experimental models and clinical studies. TNF also plays a key role in cardiovascular disease, initiating a cascade of inflammatory and vascular responses. The aim of this study is to clarify the molecular mechanisms of TNF action and the effects of its secretion in selected diseases.

Preprint

**Abstract:** Cytokines are small and membrane-bound protein-based cell signaling molecules that aid cell-to-cell communication in physiological and pathological processes. TNF- $\alpha$  is a pleiotropic cytokine, it is regarded as a principal cytokine in the acute and chronic phases of the immune and inflammatory response. TNF- $\alpha$  has been demonstrated to play a role in neuroplasticity and myelination, as well as excitotoxicity, neuroinflammation, and damage to the blood-brain barrier. The involvement of TNF- $\alpha$  in the etiology of Parkinson's disease, both in its early and late stages, Alzheimer's disease, Multiple sclerosis, coronary artery disease has been substantiated using both experimental models and clinical studies. TNF also plays a key role in cardiovascular disease, initiating a cascade of inflammatory and vascular responses. The aim of this study is to clarify the molecular mechanisms of TNF action and the effects of its secretion in selected diseases.

**Keywords:** TNF- $\alpha$ ; cytokines; central nervous system; neuroinflammation; multiple sclerosis; Parkinson's disease; dementia; coronary artery disease; plaque, intima-media

## 1. Introduction

Cytokines are molecules with the ability to regulate numerous physiological and pathological processes in living organisms. The number of newly discovered cytokines continues to grow, and the need to systematize them has led to their classification into several basic groups, including interferons, chemokines, hematopoiesis, tumor necrosis factor (TNF) superfamily cytokines and some growth factors. The basic properties of cytokines are their ability to affect different cell populations in multiple directions (pleiotropism) and redundancy – described as the ability of different cytokines to exert the same biological effect [1]. The complex pathway of interactions has led to the concept of a "net-work of cytokines" within which individual elements can act antagonistically, or – by simultaneously affecting effector cells – produce a synergistic effect.

Civilization-related diseases, such as cardiovascular disease and neurodegenerative disorders, present significant therapeutic challenges in the 21<sup>st</sup> century. Despite numerous clinical observations and experimental studies, our understanding of the pathogenesis of Parkinson's disease, stroke, and dementia and consequently their treatment, remains inadequate. The inflammatory process, along with pro-inflammatory cytokines, is thought to be a key factor in both the development of vascular lesions and the ensuing neuronal death. Therefore, this work aims to consolidate existing knowledge about TNF- $\alpha$  in relation to two of the most prevalent civilization diseases: neurodegenerative and cardiovascular conditions. Additionally, the importance of TNF- $\alpha$  will be described in the context of the pathophysiology and treatment of one of the common demyelinating diseases, i.e., multiple sclerosis (MS).

The importance of pro-inflammatory cytokines, including TNF- $\alpha$ , in the pathogenesis of many diseases is unquestionable. For instance, TNF- $\alpha$  overexpression has been associated with the clinical figures of various autoimmune and inflammatory disorders, including rheumatoid arthritis, psoriasis, or inflammatory bowel diseases. From a clinical point of view, blocking TNF signaling is currently the most effective therapeutic approach. Therefore, it is difficult to imagine treatment of the diseases listed above without drugs such as infliximab, etanercept, adalimumab, or golimumab [2]. Inflammation is a recognized residual cardiovascular disease (CVD) risk factor, the knowledge, diagnosis, and

management of inflammation-related CVD risk among healthcare professionals remain scarce. It is crucial before targeted therapies become widely available [1].

In accordance with the above, the aim of this work is to systematize knowledge of the synthesis, structure and molecular regulatory mechanisms mediated by TNF. From the clinical risks involved, we have chosen some of the most common diseases of civilization today: neurodegenerative and cardiovascular diseases. In the case of the tumor necrosis factor superfamily does not seem a completely obvious choice, even though it is known that the background of their pathogenesis is found in inflammatory reactions.

## 2. Molecular basis of TNF

The tumor necrosis factor superfamily (TNFSF) and tumor necrosis factor receptor superfamily (TNFRSF) contain 19 ligands and 29 receptors respectively. The similar chemical structure of the ligands and the resulting similar affinity with different receptors determine the collective involvement of the proteins in question in the regulation of physiological and pathophysiological processes associated with, among other things, proliferation, activation, and maturation of immune cells as well as apoptosis [4]. One of the most frequently studied representatives of the aforementioned superfamily is TNF. The tumor necrosis factor, which corresponds to the TNF- $\alpha$  form in the majority of published literature, is primarily produced by macrophages and monocytes in response to contact with bacterial cell wall lipopolysaccharides (LPS). The intensity of TNF- $\alpha$  synthesis can be illustrated by the following relationship: during LPS stimulation, the level of mRNA for the *TNF* gene in macrophages increases 100-fold, while the amount of protein increases up to 10,000 times [5]. For the sake of clarity, it should be noted that the tumor necrosis factor  $\beta$  (TNF- $\beta$ ), formerly known as lymphotoxin, is a substance produced by activated lymphocytes. The effects of TNF- $\beta$  in both physiological and pathological conditions remain poorly understood.

### 2.1. Regulation of *TNF* gene expression

The *TNF* gene is a rapidly transcribed gene in a variety of cell types following exposure to a wide range of pathogens and signals of stress or inflammation. Regulation of *TNF* gene expression at the transcriptional level is cell type and stimulus specific and involves the recruitment of distinct sets of

transcription factors to a compact and modular promoter region. The 5' UTR of the *TNF* gene contains a proximal promoter region of approximately 200 nucleotides upstream of the mRNA cap site. The *TNF* gene promoter is sufficient to drive transcription in response to a variety of stimuli, including T cell and B cell activation, calcium ionophore, bacterial LPS, viral infection, *Mycobacterium tuberculosis* and osmotic stress. TNF mRNA is transcribed within minutes and is independent of *de novo* protein synthesis. The TNF promoter contains a TATA box and cis-acting elements that are binding motifs for transcription factors: nuclear factor of activated T cells (NFAT), cyclic AMP response element (CRE) and Egr, Ets/Elk, Sp1 factors. The CRE binding site is a critical regulatory element of the TNF gene in all cell types and under all conditions tested so far [6, 7]. The LPS induces *TNF* expression in macrophages. Several studies have identified signaling pathways that are activated by LPS, including activation of nuclear factor- $\kappa$ B (NF- $\kappa$ B) and activation of mitogen-activated protein kinases (MAPKs), including ERK1 and ERK2, c-Jun N-terminal kinase and p38. The importance of extracellular signal-regulated kinase1 (ERK1) and ERK2 activation is important for LPS-induced TNF- $\alpha$  production. The ERK signaling pathway is causally involved in the synthesis of TNF- $\alpha$  by human monocytes stimulated with LPS [8; 9; 10].

TNF- $\alpha$  is a pleiotropic cytokine. From a chemical perspective, the precursor of the soluble form of TNF is a transmembrane protein with a molecular weight of 26 kDa. In response to a multitude of stimulation signals, this protein is subject to the TNF- $\alpha$  converting enzyme (commonly referred to as TRAY or ADAM 17), which ultimately results in the secretion of the soluble form of TNF- $\alpha$  (sTNF- $\alpha$ ) with a molecular weight of 17 kDa [11]. sTNF- $\alpha$  occurs in an active (trimeric) and inactive (monomeric) form.

## 2.2. *TNF* gene

TNF- $\alpha$  synthesis is modified by genetic factors, especially by variation within the *TNF* gene. The *TNF* gene is located on the shorter arm of chromosome 6 at p21.3, in close proximity to the genes of the major histocompatibility complex (MHC) [12]. The *TNF* gene structure is consistent with that observed in other genes, comprising four exons. The final exon accounts for 80% of the sequence encoding TNF- $\alpha$ . Notably, a portion of the *TNF* gene sequence is identical to that of other cytokines, including IL-1 (interleukin-1) and IFN- $\gamma$  (interferon- $\gamma$ ). The promoter region is distinguished by considerable

variability. Consequently, the majority of single nucleotide polymorphisms (SNPs) identified in the *TNF* gene are within its promoter region. Results from experimental model studies and clinical observations suggest that selected SNPs of the *TNF* gene have influence on the susceptibility to various human diseases and/or their severity [13-15]. The most studied functional SNPs in the promoter of the *TNF* gene include -308G >A (rs1800629), -1031T >C (rs799964), -863C >A (rs1800630) and -857C>T (rs1799724) [16].

### 2.3. *TNF receptors and their regulation*

TNF- $\alpha$  interacts with cells through one of the two receptors, currently designated as tumor necrosis factor receptor 1 (TNFR1) or TNFRSF1A (tumor necrosis factor receptor superfamily, member 1a), formerly p55 or p60, and TNFR2 (tumor necrosis factor receptor 2) or TNFRSF1B (tumor necrosis factor receptor superfamily, member 1b, formerly 75 or p 80).

TNFR1, also known as death receptor, in contrast to the TNFR2, which is typical of cells with a nucleus, is expressed on all cell types in the body and has a similar affinity with the membrane and soluble forms of TNF- $\alpha$ . TNFR2, which is mainly found on endothelial cells and immune cells including lymphocytes and macrophages [11], is stimulated by the membrane form of TNF- $\alpha$ . Currently, studies are being carried out to assess the mode of stimulation and functions of each receptor, but our brief characteristics of both receptors suggest that the clinical effects resulting from their stimulation vary. Primarily, TNFR1, in response to the stimulatory signal of TNF- $\alpha$ , promotes cell death and the inflammatory process, whereas binding of TNF to TNFR2 promotes cell survival and tissue regeneration. Interestingly, from a clinical perspective, cell survival is not always beneficial for the patient. Studies have shown that overexpression of *TNFR2* in cancer cells leads to tumor growth [17, 18].

TNFR receptor proteins consist of three domains: extracellular, transmembrane, and cytoplasmic. Each of these domains has a different biological role [19]. Both types of receptors, TNFR1 and TNFR2, have cysteine-rich extracellular domains capable of binding a ligand, but their intracellular domains are strikingly different.

The activation of TNFR1 by TNF- $\alpha$  results in the stimulation of four characteristic molecular complexes. The cytoplasmic 'death domain', which is characteristic of TNFR1, recruits the adaptor molecule



Another complex is divided into two structurally and functionally compatible Iia and Iib systems. The process is analogous and begins with the stimulation of TRADD (TNF- $\alpha$  receptor-associated death domain) and fas-associated protein with death domain (FADD). The remaining components of the aforementioned complex are TNF receptor-associated factor (TRAF2), RIPK1, cIAP1/2 and procaspase-8. Iib is similar to Iia except for the additional presence of RIPK3 (receptor-interacting serine/threonine- protein kinase 3). The formation of this complex is regulated independently by the long isoform of FLICE-like inhibitors of protein (FLIPL) that plays a dual role in the activation of caspases and the process of apoptosis. Stimulation of the Iia and Iib complexes may also block necrosis. However, the primary effect that they exert appears to be apoptosis (Figure 1).

Another complex, Iic, formed by RIPK1 and RIPK3, stimulates FLIPL-mediated mixed lineage kinase domain-like protein (MLKL), which leads to necroptosis. Apoptosis is different from necroptosis. In the latter, there is a rupture of the plasma membrane, release of intracellular contents into the extracellular space, and, ultimately, a local inflammatory response [21, 22]. TNFR2 lacks a cytoplasmic death domain sequence which is present in FADD. Therefore, the stimulation does not directly result in cell death. As with TNFR1, the impact of combining the transmembrane form of TNF- $\alpha$  with the TNFR2 receptor varies depending on the activated pathways. TNFR2 forms a complex with TNFR, TRAF1 and TRAF2, as well as with cIAP1 and cIAP2. This complex has been observed to interact with the NF- $\kappa$ B inducing kinase (NIK). The function of the  $\kappa$ B inhibitor kinase is phosphorylation of IKK. The IKK complex comprises a number of catalytic units that are essential for the correct progression of the phosphorylation process. One potential pathway for NF- $\kappa$ B synthesis involves stimulation of the TNFR2 receptor by the TNF- $\alpha$  and NIK activation, which subsequently affects the IKK complex. The IKK $\beta$  subunit phosphorylates I $\kappa$ B inhibitors, which results in the activation of NF- $\kappa$ B. This leads to translocation of NF- $\kappa$ B to the cell nucleus, where it binds to specific sequences within the promoter regions of target genes, including A20, cIAP-1, cIAP-2, and Bcl-xL [23, 24]. Furthermore, TNFR2 stimulation may lead to the phosphorylation of endothelial/epithelial protein tyrosine kinase (ETK/BMX), which is associated with the vascular endothelial growth factor receptor 2 (VEGFR2). This results in stimulation of the PI3K/AKT pathway – phosphoinositide 3-kinase (PI3K)/protein kinase B (AKT) – which is responsible for regulating cell survival and proliferation.



#### *2.4. TNF signaling goal*

The outcome of TNFR1-TRADD and TNFR2 signaling is cell survival or death, contingent on the stimulated signaling pathway and cell type. However, the above information suggests that TNFR2 is primarily involved in the processes such as proliferation, regeneration and survival of cells and tissues [25, 26]. Importantly, stimulation of TNFR2 increases the activity of regulatory T lymphocytes and stimulates production of IL-2, which reduces potential autoimmunity and the inflammatory process by inhibiting Th17 cell differentiation. In addition to its effect of stimulating cell survival and proliferation, TNFR2 has also been demonstrated to induce cell apoptosis. This is a consequence of the presence of common elements (e.g. receptors) which are part of the two previously described pathways (e.g. TRAF and c-IAF) [26, 27]. This fact restricts the potential for utilizing drugs with non-selective affinity for both receptor types.

#### *2.5. The role of TNF in the inflammatory process*

TNF- $\alpha$  is regarded as a principal cytokine in the acute and chronic phases of the immune and inflammatory response. As previously stated, the primary stimulus for the synthesis of TNF- $\alpha$  in macrophages is the lipopolysaccharides present in the bacterial cell wall. Furthermore, the factors responsible for the observed increase in TNF- $\alpha$  concentration include INF- $\gamma$  and IL-1, while IL-4, IL-10, PGE2 (prostaglandin E2) and glucocorticoids have been identified as inhibitors of secretion.

The fundamental effect that TNF- $\alpha$  exerts on the immune system is to increase the cytotoxicity of eosinophils, macrophages, and monocytes as well as the phagocytic capacity of neutrophils [2, 28]. Furthermore, the cytokine in question enhances proliferation and differentiation of T lymphocytes, B lymphocytes and NK cells. Given its integral role within the network of cytokines, TNF- $\alpha$  has the ability to stimulate synthesis of INF- $\gamma$ , IL-1, and IL-6. Additionally, as a result of autocrine stimulation, TNF- $\alpha$  itself undergoes an increase in concentration [29]. Furthermore, TNF- $\alpha$  exerts a direct effect on endothelial cells, leading to the increased expression of adhesion molecules and chemokines that are essential for the accumulation of white blood cells at the site of inflammation. It also induces expression of the molecules in the major histocompatibility complex (MHC I and MHC II).

The onset of acute inflammation gives rise to a cascade of cytokine and chemokine synthesis, which attract immune cells, predominantly neutrophils, to the site of damage. It is now well established that the process of acute inflammation is typically self-limiting, due to the inhibition of pro-inflammatory cytokine synthesis and stimulation of the anti-inflammatory protein activity. Inappropriate regulation within the network of cytokines may result in chronicisation of the inflammatory process, which may in turn give rise to neurodegenerative processes within the central nervous system (CNS), an elevated risk of carcinogenesis or cardiovascular complications [16, 30] (Figure 2).




Figure 2. Diseases associated with increased levels of TNF- $\alpha$ .

The pro-inflammatory effect of TNF- $\alpha$  is associated with the activation of NF- $\kappa$ B, which is synthesized in almost all cell types. The NF- $\kappa$ B factor is currently the subject of considerable interest due to the possibility of pharmacological modification of its synthesis. For the sake of clarity and coherence, it is necessary to delineate various pathways of synthesis, including the canonical and non-canonical NF- $\kappa$ B pathway [31]. It is beyond dispute that NF- $\kappa$ B plays a role in both physiological and pathological processes. Its activity can result in both protection (increased survival, proliferation, synaptic plasticity) and cell damage. Considering mainly the pathophysiological dimension, TNF- $\alpha$  activates NF- $\kappa$ B, which leads to the expression of inflammatory genes responsible for the synthesis of proteins such as

cyclooxygenase-2 (COX-2), lipoxygenase-2 (LOX-2), cell adhesion molecules, inflammatory cytokines, chemokines, and inducible nitric oxide synthase (iNOS) [31] (Figure 3).




Figure 3. TNF- $\alpha$  pro-inflammatory effect. TNF- $\alpha$  activates NF- $\kappa$ B, which leads to the expression of inflammatory genes: cyclooxygenase-2 (COX-2), lipoxygenase-2 (LOX-2), cell adhesion molecules, inflammatory cytokines, chemokines and inducible nitric oxide synthase (iNOS).

### 3. The association of TNF with diseases of the central nervous system

The central nervous system is the site at which TNF- $\alpha$  exerts cytoprotective effects as well as stimulates apoptosis. A significant number of studies currently being conducted using the experimental model and clinical observations are primarily focused on the efficacy of drug treatments that aim at reducing the biological effects of TNF- $\alpha$ . TNF- $\alpha$  has been demonstrated to play a role in neuroplasticity and myelination, as well as excitotoxicity, neuroinflammation, and damage to the blood-brain barrier [32-34]. The wide-spread use of drugs that block the synthesis of TNF- $\alpha$  limits the pleiotropic effects of this cytokine. Although anti-TNF- $\alpha$  therapy has been successfully introduced in the treatment of neurosarcoidosis, it has been observed that blocking its synthesis in patients with MS may, paradoxically, result in exacerbation of the disease. There are evidences, that that intravenous treatment of MS patients with the anti-TNF antibody may lead to intrathecal immune activation and may therefore be harmful for MS patients [35, 36]. Similarly, administration of TNF- $\alpha$  in experimental autoimmune encephalomyelitis (EAE) mice has shown to exacerbate the course of the disease [20].

Table 1. The biological effect of TNF- $\alpha$  on the central nervous system

## TNF $\alpha$ in Neurodegenerative Diseases

Biological effect		
Physiological effect (TNFR1 stimulation)	Synaptic plasticity	<ul style="list-style-type: none"> <li>• Increase in the expression of AMPA and NMDA receptors</li> <li>• Increase in the release of glutamates and S100B protein from astrocytes</li> <li>• Decrease in GABA expression</li> </ul>
Physiological effect (TNFR2 stimulation)	Oligodendrocytes	<ul style="list-style-type: none"> <li>• Proliferation</li> <li>• Maturation</li> <li>• Remyelination</li> </ul>
Pathophysiological effect (TNFR1 stimulation)	<p>Microglia</p> <p>Excitotoxicity stimulated by an imbalance between excitatory neurotransmitters (glutamate) and inhibitory neurotransmitters (GABA)</p> <p>Disorders of the blood-brain barrier permeability.</p>	<ul style="list-style-type: none"> <li>• Increase in the secretion of pro-inflammatory cytokines</li> <li>• Increased synthesis of TNF-<math>\alpha</math> and glutamate</li> </ul>

AMPA – amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor

NMDA – N-methyl-D-aspartate receptor

GABA A – gamma-aminobutyric acid A receptor

TNF- $\alpha$  should be treated as a cause of neurological diseases. The results indicate that TNF- $\alpha$  induces neurological dysfunction. TNF- $\alpha$ -induced brain dysfunction and disease are caused by necroptosis and microglial activation in hippocampal neurons. TNF- $\alpha$  is an important cytokine involved in various homeostatic and pathogenic bioactivities [37]. The dualistic nature of TNF- $\alpha$  is particularly evident in the central nervous system. As detailed in Table 1, this cytokine is involved in both inflammatory processes and cell death. It also plays a key role in maintaining homeostasis of the central nervous system by affecting glial cell proliferation and neuronal function. TNF- $\alpha$  plays a special role in regulating survival of oligodendrocyte cells as well as formation and repair of myelin. This explains the limited use of drugs that reduce synthesis or the biological activity of TNF- $\alpha$  in patients with multiple sclerosis.

### *3.1. Parkinson's Disease*

Parkinson's disease (PD) is the second most common neurodegenerative disease after Alzheimer's disease. The estimated prevalence of PD is 0.3% in the general population and 1.0% or 3.0%, respectively, in people older than 60 and 80. A special group of patients is those with earlier onset of PD symptoms (Early-onset Parkinson's disease; EOPD), which is defined as Parkinson's disease with age at onset (AAO) after age 21 years but before the usual AAO for PD. Consensus is lacking, and the reported maximal age for EOPD has varied from 40 to 60 years [38]. The aetiology of Parkinson's disease is not well established. For several decades, environmental factors have been considered primarily responsible for disease development. However, recent studies have demonstrated that Parkinson's disease has a substantial genetic component, probably causing or influencing at least 10 to 25% of cases, more commonly in younger patients. Therefore, the new concept of PD as an "eco-genetic" disease assumes an influence of environmental and genetic factors coinciding with the aging process [39]. Indeed, a family history of PD is reported in 20% of patients with EOPD versus 6.9% of patients with later onset PD, and the age-specific risk of PD is 7.8-fold higher in the relatives of patients with EOPD compared with 2.9-fold among the relatives of patients with later onset PD [38]. Recent

findings have pointed out inflammation as a potential factor contributing to the PD development. It was hypothesized that genetically determined differences in the immune system function, especially anti- and pro-inflammatory cytokines production, might influence the risk of sporadic PD development and/or onset [40, 41]. Of particular interest within the group of pro-inflammatory cytokines is TNF- $\alpha$ . The involvement of TNF- $\alpha$  in the aetiology of PD, both in its early and late stages, has been substantiated using both experimental models and clinical studies [42]. TNF- $\alpha$  and TNFR receptors are responsible for the activation of two independently running pathways. As previously stated, each can result in apoptosis or an increase in the expression of numerous anti-apoptotic genes [43]. In Parkinson's disease, the negative effect of the cytokine on the survival and function of neurons appears to be predominant. The results of studies using experimental models showed a significant increase in the concentration of mRNA for TNF- $\alpha$  in the substantia nigra (SN) of rats treated *in vivo* with the most commonly used experimental neurotoxin MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) [44]. TNF- $\alpha$  not only induced inflammatory responses but was also involved in the maintenance of the inflammatory process that had already been initiated. This is demonstrated by the sustained elevation of the cytokine in question one year following the administration of the neurotoxin. One of the mechanisms proposed as a potential cause of the neurodegeneration characteristic of parkinsonism is the impact of TNF- $\alpha$  on the functions of microglia. TNF- $\alpha$  activates microglia, which leads to a progressive loss of dopaminergic neurons in the SN [45]. As shown by clinical studies, low concentrations of TNF- $\alpha$  in the brains of healthy adults are observed in contrast to high concentrations of TNF- $\alpha$  and the soluble form of the TNFR1 receptor in the cerebrospinal fluid as well as the SN obtained from the brains of patients diagnosed with PD [44-46]. Dopamine neurons in the substantia nigra of the brain exhibit a high degree of sensitivity to the effects of TNF- $\alpha$ . TNF- $\alpha$  can activate microglia in the interbrain, stimulate the inflammatory response and exacerbate oxidative stress through ROS- and NO-dependent mechanisms. Elevated levels of TNF- $\alpha$  have been shown in the substantia nigra of patients with PD and atypical parkinsonism. Elevated levels of TNF- $\alpha$  have also been found in the cerebrospinal fluid of people with REM sleep behaviour disorder [45,47, 48]. This is a particularly interesting observation given that RBD (Rapid eye movement sleep behaviour disorder) is considered a non-motor manifestation of prodromal synucleinopathy [49]. TNF- $\alpha$  levels may also influence the progression of the disease. As shown by

Menza et al. TNF- $\alpha$  levels were positively correlated with cognitive impairment, depression, and disability in patients with PD [50]. Furthermore, Wang et al. investigated the correlation between plasma levels of inflammatory cytokines as well as alpha-synuclein ( $\alpha$ -syn) and fatigue in patients with Parkinson's disease. The study included 63 patients with Parkinson's disease, 35 patients with symptoms of fatigue and 28 patients without clinical symptoms of fatigue. In addition to factors related to the progression of the disease, the degree of advancement and older age, the authors demonstrated that plasma levels of cytokines IL-1 $\beta$ , IL-18, TNF- $\alpha$  and phosphorylated  $\alpha$ -syn (p- $\alpha$ -syn) exhibited a positive correlation with the results on the Fatigue Severity Scale 9 (FSS-9). Furthermore, a higher score on the Parkinson's Fatigue Scale 16 (PFS-16) was associated with elevated levels of TNF- $\alpha$  and p- $\alpha$ -syn in plasma. The findings suggest a potential correlation between elevated TNF- $\alpha$  and p- $\alpha$ -syn levels in plasma and the onset of fatigue in individuals with Parkinson's disease [51].

The confirmation of the relationship between TNF- $\alpha$  and the risk of PD, as well as the progression of the disease, presents an opportunity for pharmacotherapy utilizing drugs that block the activity of the cytokine in question. The findings of the research studies that have been published to date are encouraging. For example, Peter et al. demonstrated that the utilization of TNF inhibitors was linked to a 78% reduction in the likelihood of PD (incidence rate 0.22; 95% CI 0.05, 0.88) [52]. The study was conducted on a group of 144,018 individuals diagnosed with inflammatory bowel disease (IBD), matched for age and sex with a control group of individuals unaffected by IBD, who were treated with adalimumab, certolizumab, golimumab, and infliximab.

The pathogenesis of diseases of the central nervous system, including Parkinson's disease, is a complex process. In addition to environmental factors, there is evidence to suggest that genetic predisposition also affects the risk and progression of Parkinson's disease. The results of genetic tests conducted on the variability of genes encoding the *TNF- $\alpha$*  and *TNFR* activity are inconclusive [53, 54]. Although there are isolated reports of neuroprotective effects associated with high concentrations of TNF- $\alpha$ , the cytokine generally exerts an opposing effect to that of IL-10 in the majority of processes occurring in the human body. Genetically determined inter-individual differences in cytokine production may be at the root of differences in the progression of the inflammatory process [55, 56]. The *TNF* gene polymorphism, specifically the -308G >A variant, which has been most extensively studied in the

Caucasian population, is associated with differences in the expression of the encoded protein. The -308A variant allele increases the transcriptional activity of the gene compared to the more common G allele, which may clinically correspond to the severity or dysregulation of inflammatory responses. The study by Nishimura et al. on a group of patients with sporadic PD showed that SNP -1031T>C, -863C>A and -857C>T may influence the risk and age of developing PD in oriental populations [56]. In patients with early-onset Parkinson's disease (EOPD), the frequency of the -1031C allele, which is associated with increased transcriptional activity and TNF- $\alpha$  concentration, was significantly higher compared to patients with late-onset Parkinson's disease (LOPD) and healthy subjects. Furthermore, the -1031C allele and the -1031CC genotype were identified as risk factors for Parkinson's disease [5]. It is noteworthy that the -1031C allele, which is associated with increased TNF- $\alpha$  synthesis in oriental population patients, corresponds phenotypically to the -308A variant of the -308G >A polymorphism, which is prevalent in the Caucasian population.

Subsequent studies by Kang et al. investigated similar issues, focusing on the genetic variability of *TNFRSF1A*, which is responsible for the expression of the *TNFR1* receptor [57]. The aim of the study was to assess the association between selected genetic variants of the gene encoding the *TNFR1* receptor and the development of Parkinson's disease. Previous studies linked the effect of pharmacological blockade of TNFR1 signaling during anti-TNF therapy to genetically determined *TNFRSF1A* variability. Genetic associations with lower levels of circulating C-reactive protein, the effects of TNF-TNFR1 inhibition were estimated for PD risk and age at the onset of PD using GWAS (Genome-Wide Association Studies) data from the International Parkinson's Disease Genomics Consortium and 23andMe, Inc. Despite the large number of control groups studied, the importance of inhibiting TNF-TNFR1 signaling in preventing or delaying the onset of PD symptoms has not been confirmed.

### 3.2. Alzheimer's dementia

Alzheimer's disease (AD) is one of the most common forms of dementia. A typical pathological feature of AD is the accumulation of  $\beta$ -amyloid ( $A\beta$ ) plaques and neurofibrillary tangles. The pathogenetic model under consideration fails to account for a substantial proportion of the clinical and pathological variability observed in the disease. Consequently, further hypotheses have been formulated, of which



one suggests that chronic inflammation within the CNS may be a significant contributing factor to the development and progression of the disease [58]. Hypotheses of the AD pathophysiology include also amyloid cascade, inflammation, vascular, and infection factors [59, 60].

In the CNS, TNF- $\alpha$  is secreted by various cell types, mainly microglia. Due to the complex interactions of TNF- $\alpha$  with other cytokines, this factor plays a key role in the regulation of numerous physiological and pathophysiological changes in the brain. Primary, TNF- $\alpha$  exerts an effect of increasing synaptic transmission by both direct and indirect means, including the additional release of glutamate from astrocytes. By activating pathways associated with NF- $\kappa$ B, it is involved in regulating the neuronal function in the key areas of the brain that regulate sleep [61]. TNF- $\alpha$  is a major pro-inflammatory cytokine that stimulates production of IL-1, IL-6 and IL-8, promoting chronic inflammation, especially when the production of pro-inflammatory cytokines is not balanced by anti-inflammatory cytokines such as IL-10 [62].

The importance of TNF- $\alpha$  in the pathogenesis of AD has been confirmed in clinical studies and experimental models, both in the context of beta-amyloid and tau protein [63]. Experimental studies have shown that TNF- $\alpha$  increases the expression of myeloid precursor protein (APP) and  $\beta$ -site amyloid precursor protein cleaving enzyme-1 (BACE1) in mouse astrocytes. Furthermore, it has been demonstrated that it stimulates the activity of  $\gamma$ -secretase in HEK cells, thus releasing A $\beta$  peptides [64]. The splitting of microglia, a phenomenon observed in dementia, has been shown to facilitate production of TNF- $\alpha$ . This, in turn, sustains the persistence of chronic inflammation, thereby stimulating the synthesis of A $\beta$  and, subsequently, contributing to the loss of neurons. Recently, another concept associated with triggering the receptor expressed on myeloid cells 2 (TREM2) has emerged regarding the role of TNF- $\alpha$  in the pathology of Alzheimer's disease. The TREM2 is found in various tissue macrophages, such as CNS microglia, bone osteoclasts, alveolar, peritoneal, and intestinal macrophages. TREM2 is also present on cultured bone marrow-derived macrophages and monocyte-derived dendritic cells. Some researchers have identified a rare variant of TREM2 that is a risk factor for Alzheimer's disease [65]. TREM2 has been reported to bind to proteins with a strong tendency to aggregate and accumulate in neurodegenerative diseases, such as  $\beta$ -amyloid and supports the microglial response to A $\beta$  plaques [66]. The Guerreiro et al. and Jonsson et al. showed the association between variability in

the gene encoding the *TREM2* expression and the risk of late-onset sporadic Alzheimer's disease. The authors indicated that heterozygous hypomorphic variants of *TREM2*, mainly *TREM2R47H*, increase the risk of AD [67,68]. Impaired receptor function is responsible for the limited ability of microglia to encase A $\beta$  plaques, leading to the spread of  $\beta$ -amyloid accumulation, severity of neurotoxicity and the progression of the disease. At present, studies are being carried out to assess the efficacy of *TREM2* agonists in the treatment of AD [69].

Clinical studies of patients with mild cognitive impairment (MCI) showed higher levels of TNF- $\alpha$  and tau protein and lower levels of A $\beta$ 42 and the anti-inflammatory cytokine TGF $\beta$  (Transforming Growth Factor  $\beta$ ) in patients with MCI compared to the control group [70]. In addition, the authors confirmed the relationship between elevated levels of TNF- $\alpha$  in the cerebrospinal fluid and the conversion of MCI to Alzheimer's disease. However, this observation was not confirmed in recent studies by Fu et al. [71]. Clinical evidence for the importance of TNF- $\alpha$  in the pathogenesis of CNS diseases, including AD, is provided by observational studies of patients with systemic inflammatory diseases treated with TNF- $\alpha$  inhibitors [72,73]. Zhou et al. published the results of a retrospective clinical study based on the medical records of 56 million adult patients [72]. The aim of the study was to investigate the link between inflammatory diseases and AD dementia as well as answer the question of whether the drugs that block tumor necrosis factor functions, used in patients with rheumatoid arthritis (RA), psoriasis and other inflammatory diseases, affect the risk of AD. The study demonstrated an elevated susceptibility to AD development among the study group of patients, with those diagnosed with rheumatoid arthritis, inflammatory bowel disease, or Crohn's disease exhibiting the highest risk ( $p < 0.0001$ ). Importantly, the use of drugs that target TNF- $\alpha$  may influence the development of dementia. Etanercept, adalimumab or infliximab were associated with a significantly lower risk of AD in patients with RA. In patients with psoriasis, this association has only been confirmed for etanercept and adalimumab. This study may provide a basis for further clinical trials of drugs in this class, with a potentially protective mechanism of action, particularly in patients with inflammatory diseases who are predisposed to developing dementia.

The current evidence suggests that the clinical efficacy of drugs in this class is more significant in patients with inflammatory conditions. For example, Chen et al. found cognitive improvement in elderly

patients with RA following the administration of etanercept and adalimumab. Conversely, etanercept demonstrated minimal impact on the core symptoms in patients with mild and moderate AD [74, 75]. There is little interest in the use of this group of drugs, and currently no clinical trials are being conducted [<https://clinicaltrials.gov/>]. This may be related to the pharmacokinetic properties of this class of drugs. The limited efficacy of subcutaneous anti-TNF- $\alpha$  therapies for AD may be attributed to the high molecular weight of anti-TNF- $\alpha$  monoclonal antibodies, which presents a challenge in their crossing the blood-brain barrier under physiological conditions.

#### **4. Neuroinflammatory diseases.**

Neuroinflammation is a common pathological characteristic of several CNS disorders. It is characterized by the activation of glial cells, demyelination, and neurodegeneration. In autoimmune diseases of the CNS, TNF- $\alpha$  is a key inflammatory cytokine that regulates peripheral and local immune responses. Its involvement in excitotoxicity and neuroinflammation highlights its impact on CNS pathophysiology and emphasizes its multifaceted roles in regulating cellular responses and inflammatory processes within the CNS. Neuroinflammatory diseases encompass a range of conditions, including neurosarcoidosis, neuro-Beçet's disease — a systemic inflammatory vasculitis of unknown origin — and MS. However, given the breadth of this subject, this article will focus solely on multiple sclerosis.

##### *4.1. Multiple sclerosis*

MS is an autoimmune demyelinating and neurodegenerative disease of the CNS. The most well-known concept of MS is based on the co-occurrence of genetic susceptibility and a specific immune system response against its own antigens, which occurs in response to environmental factors. The signal that initiates demyelination is the activation of CD4+ T lymphocytes in response to myelin antigens. Myelin proteins are presented by antigen-presenting cells (APC) in combination with the MHC class II antigens. Subsequent steps involve the passage of autoreactive T lymphocytes to the CNS and differentiation into subclasses showing pro-inflammatory (Th1 and Th17) or anti-inflammatory (Th2) activity [76,77]. Proinflammatory cytokines, including TNF- $\alpha$ , IFN- $\gamma$ , IL-2, IL-17, are involved in the activation of microglia, B lymphocytes, as well as the influx and stimulation of peripheral monocytes and

macrophages. This briefly outlined pathway of pathophysiological changes indicates the importance of cytokines, including TNF- $\alpha$ , in the initiation and propagation of the autoimmune process that characterizes demyelination in the progression of MS.

Clinical studies of patients with multiple sclerosis have shown a correlation between both the severity of symptoms as well as the progression of the disease and elevated levels of TNF- $\alpha$  in the cerebrospinal fluid [78]. However, a double-blind, placebo-controlled Phase 2 study of 168 patients with relapsing-remitting MS treated with lenercept showed exacerbation of the disease. The study revealed that patients treated with lenercept (p55 tumor necrosis factor receptor fusion protein) exhibited a significantly higher incidence of disease exacerbation at the 48-week mark compared to those receiving placebo ( $P = 0.007$ ). However, treatment had no influence on the Expanded Disability Status Scale (EDSS) score in either group after 24 weeks of treatment and at the final clinical assessment [79]. The explanation for the above observation should be sought in the results of studies conducted using animal models. It has been suggested that chronic inhibition of TNF- $\alpha$  may enhance the T lymphocyte response to a specific antigen. Paradoxically, drugs that inhibit TNF- $\alpha$  activity may enhance the pro-inflammatory activity in the CNS as well as increase survival of peripheral myelin-autoreactive T lymphocytes. Furthermore, drugs based on this mechanism of action may result in a reduction of IL-10 levels and an increase in IL-12 and IFN- $\gamma$ , thereby creating a profile comparable to that observed in patients with multiple sclerosis [80,81]. Another theory is that the distribution of TNF- $\alpha$  levels between the CNS and peripheral tissues is dysregulated (higher levels in the CNS due to the impermeability of the blood-brain barrier). The non-selective inhibition of the TNF- $\alpha$  activity by affecting two receptors (TNFR1 and 2) may be considered potentially harmful for patients with multiple sclerosis. This is also confirmed by the study conducted on a group of patients with rheumatic diseases in whom anti-TNF- $\alpha$  therapy resulted in the appearance of demyelinating lesions [82].

Drugs that non-selectively inhibit TNF- $\alpha$  binding to the receptor and inactivate its effect limit transmembrane interaction with both TNFR1 and TNFR2. As mentioned earlier in this article, TNFR2 plays an important role in the proliferation of immature oligodendrocytes, remyelination and inhibition of autoreactive CD4<sup>+</sup> T lymphocytes. Although some studies have questioned the role of anti-TNF- $\alpha$  drugs in the demyelinating process [82, 83], until drugs that selectively block the TNFR1 activity are

developed, drugs in this group should be used with great caution, especially in the case of MS comorbidity with rheumatological diseases. Another question is the risk of inflammatory central nervous system diseases in patients receiving anti-TNF therapies. The results of a systematic review and meta-analysis of 18 studies, including more than 1 million patients with autoimmune diseases, were published by Xie et al. last year [84]. The incidence rates of new-onset inflammatory CNS events after initiating anti-TNF therapies ranged from 2.0 to 13.4 per 10,000 person-years, and overall, exposure to this treatment was associated with a 36% increased risk of any inflammatory CNS disease compared to conventional therapies (RR, 1.36; 95% CI, 1.01-1.84; I<sup>2</sup>, 49%) with similar risk across different types of autoimmune diseases and TNF inhibitors.

Table 2. Summary of selected study presenting the role of TNF-  $\alpha$  in neurology.

Disease	Experimental model/clinical trials	Reference
<b>Parkinson's disease</b>	<b>Experimental studies</b>  A significant increase in the concentration of mRNA for TNF- $\alpha$ in the substantia nigra (SN) of rats treated <i>in vivo</i> with experimental neurotoxin MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine)	<b>44</b>
<b>Parkinson's disease</b>	<b>Clinical trials</b>  Low concentrations of TNF- $\alpha$ in control group in contrast to high concentrations of TNF- $\alpha$ and the soluble form of the TNFR1 receptor in the cerebrospinal fluid as well as the SN obtained from the brains patients diagnosed with PD.	<b>46</b>

	Increased serum levels of sTNFR1 in PD patients, no differences in TNF- $\alpha$ and sTNFR2 levels between PD and control.	47
<b>Parkinson's disease</b>	<b>Clinical trials</b>  TNF- $\alpha$ serum levels positively correlated with cognitive impairment, depression, and disability in patients with PD	50
<b>Parkinson's disease</b>	<b>Clinical trials</b>  A higher incidence of PD was observed among patients with inflammatory bowel disease than among individuals without IBD. Early exposure to anti-inflammatory anti-TNF therapy was associated with substantially reduced PD incidence.	52
<b>Parkinson's disease</b>	<b>Clinical trials (study using genome-wide association studies (GWAS))</b>  The findings do not imply that TNF-TNFR1 signaling inhibition will prevent or delay PD onset.	57
<b>Alzheimer's disease</b>	<b>Clinical trials</b>  Increased production of the proinflammatory cytokine, TNF- $\alpha$ and decreased production of the anti-inflammatory cytokine TGF- $\beta$ in	70

	<p>cerebrospinal fluid patients with mild cognitive impairment (MCI) at risk to develop AD</p>	
<p><b>Alzheimer's disease</b></p>	<p><b>Clinical trials</b></p> <p>The study showed an elevated susceptibility to AD development among patients diagnosed with rheumatoid arthritis, inflammatory bowel disease, or Crohn's disease. For RA and psoriasis, treatment with a biologic drug that targets TNF correspondingly is associated with decreased risk for AD.</p>	<p>72</p>
<p><b>Multiple sclerosis</b></p>	<p><b>Clinical trial</b></p> <p>Mean serum TNF-<math>\alpha</math> levels were significantly higher in patients with MS than control group. TNF-<math>\alpha</math> levels were found to be negatively correlated with the Expanded Disability Status Scale (EDSS) score</p>	<p>78</p>
<p><b>Multiple sclerosis</b></p>	<p><b>Clinical trials</b></p> <p>The study revealed that patients treated with lenercept (p55 tumor necrosis factor receptor fusion protein) exhibited a significantly higher incidence of disease</p>	<p>79</p>

	exacerbation at the 48-week mark compared to placebo group	
<b>CNS inflammatory diseases</b>	<b>Clinical trials</b> The anti-TNF therapies was associated with a 36% increased risk of any CNS inflammatory disease compared to conventional therapies (RR, 1.36; 95% CI, 1.01-1.84; I <sup>2</sup> , 49%) with similar risk across different types of autoimmune diseases and TNF inhibitors	<b>84</b>

## 5. The association of TNF with cardio-vascular diseases

The pathogenesis of cardiovascular diseases involves many conventional predictors, as dyslipidemia, hypertension, obesity, diabetes, smoking and new risk factors including genetics factors. Studies have shown that inflammation plays a role in the pathophysiology of atherosclerosis and CAD and is increasingly seen as a new modifiable cardiovascular risk factor. Inflammatory markers are important mediators in the multi-step cascade of atherosclerosis, which finally leads to the rupture of the atherosclerotic plaque [85].

### 5.1. Coronary artery disease

In the western civilization coronary artery disease (CAD) is associated with high morbidity and mortality and has therefore become a major public health challenge [86]. The pathogenesis of atherosclerosis is multifactorial and diverse. It is associated with the activation of several signaling pathways through inflammatory or mechanical damage to the vascular endothelium, thus mobilizing the body's immune processes and involving pro-inflammatory cytokines [87, 88].

Furthermore, there is an increasing recognition of inflammation as a novel modifiable risk factor for cardiovascular disease. TNF is a well-established pro-atherosclerotic factor [89]. Binding of TNF to its



receptor (TNF-R1 or TNF-R2) on the surface of the cell membrane leads to the activation of cytokine production and initiation of the arachidonic acid cascade. This increases the intracellular concentration of free radicals and stimulates the liver to produce acute phase proteins and adipose tissue to produce adiponectin [90]. As a pro-inflammatory cytokine, TNF can be produced by cardiomyocytes. This occurs under unfavorable haemodynamic conditions, excessive pressure, or volume load. Consequently, it can lead to myocardial metabolic dysfunction, reduced peripheral blood flow, left ventricular dysfunction or its reconstruction [91]. The precise mechanism through which TNF exerts this effect remains unclear. It was observed that there is an inverse correlation between TNF and the E/A ratio [92]. In contrast, in patients with atrial fibrillation, the level of TNF correlates with the volume of the left atrium, which is a reliable indicator of diastolic function. Higher plasma TNF concentrations are associated with a higher risk of left ventricular hypertrophy and therefore may be a potential risk factor for diastolic dysfunction in patients with early CAD [93, 94]. TNF levels are also inversely correlated with a percentage of ejection fraction [95] and QT prolongation, even in patients without heart failure. TNF is responsible for this phenomenon through the stimulation of reactive oxygen species (ROS). The "electric storm" of recurrent ventricular arrhythmias observed in heart diseases is also associated with elevated levels of this cytokine [96]. TNF levels measured 48 hours after a cardiovascular episode have been shown to have a sensitivity of 78% and a specificity of 72.5% in predicting the occurrence of ischaemic cardiovascular incidents, including angina, re-infarction, heart failure and death [97, 98]. It has been demonstrated that the likelihood of death can be predicted by the degree of TNF elevation, which reduces myocardial contractility in patients with heart failure by disrupting intra-cellular calcium homeostasis [99]. Studies have also shown an association between elevated levels of inflammatory markers, including TNF- $\alpha$ , with the severity of CAD and the likelihood of future cardiovascular occurrences. This phenomenon can be attributed to the adaptive immune response of the body, which is caused by the prolonged retention of plasma lipoproteins and subsequent oxidative modification. This ultimately leads to endothelial damage. Recruited immune cells coordinate early atherosclerotic lesions by releasing pro-inflammatory cytokines, thereby accelerating foam cell formation, and ultimately promoting coronary syndrome through various inflammatory cascades [100]. Pericardial adipose tissue represents a small yet distinctive depot of fat, situated in close proximity to the cardiac muscle and

coronary arteries. This proximity allows for the penetration of cells and proteins into the surrounding tissues [101]. TNF- $\alpha$ , derived from epicardial adipocytes, is the major mediator of local inflammation in several areas surrounding coronary arteries in patients with advanced CAD [102]. Coronary artery bypass graft (CABG) surgery, which is used to save the life of people CAD, can also induce an inflammatory response, release cytokines, including TNF, and lead to complications [103]. It is established that mean TNF levels in blood serum are significantly higher in patients with arterial hypertension and patients with acute myocardial infarction or stroke [104, 105]. Increases in circulating pro-inflammatory cytokines in patients with heart failure, particularly TNF, may reduce NO (nitric oxide) synthesis by decreasing the expression of endothelial synthase of nitric oxide – a key enzyme involved in NO production. NO, however, acts as a regulator of muscle tension in the vascular endothelium/muscularis mucosae system. It is also an inhibitor of adhesion, platelet aggregation and angiogenesis [106]. TNF promotes hypercholesterolemia and hypertriglyceridemia by reducing the lipoprotein lipase activity in adipocytes, prolonging LDL (Low Density Lipoprotein) retention in vascular walls and activating NF- $\kappa$ B and PPAR- $\gamma$  (Peroxisome proliferator-activated receptor gamma) [107, 108]. The atherosclerotic effect of TNF inhibits HDL (High Density Lipoprotein) and reduces cytokine release [109, 110]. In general, inverse associations have been observed between ApoA1, HDL-C, and ApoA1/HDL-C ratios and levels of inflammatory markers, including TNF- $\alpha$ . Furthermore, statins have been demonstrated to reduce TNF levels in plasma [111-113]. This is of particular importance in the case of patients with coronary artery disease. An inverse relationship between HDL and TNF- $\alpha$  levels is also observed in patients with cerebrovascular ischaemia [114, 115].

There is a dearth of publications documenting the strong statistical relationship between cytokines and measurements of atherosclerotic vascular lesions in the Doppler study. It is established that the concentration of soluble TNF receptors in serum is significantly correlated with the thickness of atherosclerotic plaques [116-118] and that the ex-pression of TNF receptor on monocytes is positively related to the thickness of the inner and middle carotid artery [119]. In elderly patients, the length, thickness, and number of carotid atherosclerotic plaques, as well as TNF levels in plasma, decrease after they are treated with statins [120]. In contrast, in cultured endothelial cells, the neuropeptide catestatin (an endogenous nicotine cholinergic antagonist) inhibited the TNF- $\alpha$ -induced expression of adhesion

molecules and inflammatory cytokines by activating angiotensin-2 converting enzyme (ACE2). Administration of the drug reduces interaction of leukocytes with the endothelium and acts as a pleiotropic cardioprotective hormone [121].

Inflammation plays an important role in atherosclerosis and CAD. Cytokines produced by immune cells, such as TNF- $\alpha$ , can be the target of therapeutic and anti-inflammatory molecules. The main anti-inflammatory therapies are aspirin, statins, colchicine, IL-1 $\beta$  and IL-6 inhibitors, and new therapies such as TNF- $\alpha$  inhibitors and IL-1 receptor antagonists. Aspirin and statins reduce TNF levels, but TNF inhibitors are not yet used in clinical practice. The effect of TNF- $\alpha$  inhibitors such as infliximab, etanercept and adalimumab on atherosclerosis has been investigated in preclinical studies. Infliximab reversed TNF- $\alpha$ -induced endothelial dysfunction. Etanercept may prevent TNF-induced endothelial cell apoptosis, while adalimumab may improve endothelial function and inflammation in patients. A meta-analysis of epidemiological and randomized control trials found that patients on anti-TNF therapy had a reduced risk of cardiovascular events, but this effect was not significant [100, 122-127].

CAD is a multifactorial disease in which many predictors are known, but new research is emerging on their dependence on signaling pathways. The introduction of new drugs, such as TNF inhibitors, must therefore consider the complexity of metabolic interactions [128]. Increased *PCSK9* expression in the liver is positively associated with TNF- $\alpha$ , but a recent study reported no significant effects on leukocytes, monocytes or neutrophils and no significant reduction in TNF- $\alpha$  after treatment with PCSK9 antagonists [129]. Different classes of drugs commonly used in the treatment of CAD have been shown to have different effects on inflammatory biomarkers. However, the current knowledge on inflammation and cardiovascular risk points to the importance of diagnostic criteria and evaluation of the anti-inflammatory and pro-inflammatory profile of the main treatment used in CAD therapy for clinicians [123, 130]. Knowledge of the inflammatory basis of CAD has not yet been supported by several large, controlled intervention trials. Selective TNF inhibitors may have led to unknown complications. A follow-up study is needed to improve their safety. A further trial investigating the efficacy of anti-inflammatory treatment in CAD patients' needs to evaluate the effects of low-dose treatment versus placebo and assess the effects on the lipid profile.

## 5.2. Stroke

The redox reaction and stress are aetiologies of vascular lesions [131]. The major reactive oxygen species (ROS) produced in response to several stimuli, such as hyperglycemia, hyperlipidemia, and hypertension is superoxide anion ( $O_2^{\cdot-}$ ). It combines with NO to produce peroxynitrite leading to endothelial dysfunction by decreasing NO bioavailability. In response, endothelial cells become activated, produce vasoconstrictor agents: endothelin-1, thromboxane A2 or prostaglandin H2. That way an inflammatory response is initiated. The endothelium starts expressing adhesion molecules and secretes chemokines to attract immune cells. The increased expression of chemokines and proteases in endothelial cells perpetuates the inflammatory response [132]. In stroke, the lack of oxygen and nutrients can cause neurovascular units to become dysfunctional [133], affecting the production of ROS and the secretion of pro-inflammatory mediators [134]. Blood vessel occlusion or underperfusion affects the immune response near the ischemic parenchyma and then extends to the ischemic zone. Microglia are activated and promote the release of TNF- $\alpha$  [135]. TNF levels in brain tissue increase up to one day after ischaemic injury and correlate with its severity [136]. The effects of TNF on the vascular endothelium include: oxidative stress through stimulation of xanthine oxidase expression, expression of tissue factors, production of leukocyte adhesion molecules and activation of matrix metalloproteinases. These actions trigger local vascular inflammation, bleeding, and thrombosis [137]. TNF- $\alpha$  also disrupts the protective barrier between the brain and blood circulation, affects the activation and proliferation of astrocytes and microglia, and regulates apoptosis factors such as cysteine secretion and the transcription of metalloproteinases and inflammatory cytokines [138]. TNF- $\alpha$  has neurotoxic or neuroprotective effects in stroke [139]. In several studies, TNF- $\alpha$  has been associated with movement disorders, aphasia, epileptic seizures, pain, and cognitive impairment [140-142]. TNF is used as a potential therapeutic target in stroke, and its blockade reduces focal ischaemic injury and improves clinical outcomes [143]. Therefore, detection of high levels of TNF may be useful as an inflammatory marker to predict stroke outcome. There are three effective ways to interfere with TNF: blocking TNF receptors, interfering with TNF signaling, and removing TNF protein [144]. TNF- $\alpha$  inhibitors, including infliximab, enanercib, adalimumab, golimumab and pertuzumab, are currently used to treat inflammatory or autoimmune

diseases, but may also reduce the risk of stroke [145]. The safety of anti-TNF agents and their lower efficacy in penetrating the blood-brain barrier need further investigation [146].

Table 3. Summary of selected study presenting the role of TNF-  $\alpha$  in cardiovascular diseases.

<b>Disease</b>	<b>Experimental model/clinical trials</b>	<b>Reference</b>
<b>CAD</b>	<p><b>Experimental studies</b></p> <p>The aim of the study was to assess the relationship between plasma levels of the TNF cytokine with clinical risk factors for cardiovascular disease, along with the parameters of cardiac morphology in patients diagnosed with coronary artery disease (CAD) at a young age. Results suggest an association between higher plasma TNF concentrations and components of metabolic syndrome, including dyslipidemia. TNF may be a potential risk factor for impaired diastolic function.</p>	<b>94</b>
<b>CAD</b>	<p><b>Review</b></p> <p>Aim of publication was to provide indication on the anti-inflammatory profile of agents of general use in cardiovascular prevention, <i>i.e.</i> affecting lipids, blood pressure, diabetes as well nutraceuticals such as n-3 fatty acids. A crucial issue in the evaluation of the benefit of the anti-inflammatory activity is the frequent discordance between a beneficial activity on a major risk factor and associated changes of others, as in the case of statins.</p>	<b>126</b>
<b>cardiovascular diseases</b>	<b>Randomized trials and meta-analysis</b>	<b>127</b>

	Twelve randomized controlled trials with a total of 13 073 patients, showed that colchicine reduced cardiovascular events and inflammatory markers in patients with CAD.	
<b>stroke</b>	<p><b>Experimental studies</b></p> <p>The study was conducted to identify independent predictors of severity and short-term outcome of first-ever ischemic stroke in young adults. In two years, observation retrospectively enrolled 325 consecutive patients aged 18-49 years with first-ever ischemic stroke.</p>	<b>135</b>

## 6. Conclusions

In response to harmful stimuli or dysregulation in the CNS and cardio-vascular system, pro-inflammatory cytokines, such as TNF- $\alpha$ , are released by endothelial cells, neurons, and glial and infiltrating immune cells. As mentioned earlier, the inflammation may be harmful or protective depending on different factors that include the activation causes, targets, transcriptional changes, and resulting intercellular activities. For this reason, overexpression of TNF- $\alpha$  should be considered an effect of deleterious stimuli. The elevated level of this pro-inflammatory cytokine may cause neuronal excitotoxicity, synapse loss, and the development of neurological and cardiological diseases, however the importance of TNF- $\alpha$  as a biomarker is ambiguous. For instance, although some current studies do not support TNF- $\alpha$  as a clear marker of stroke, it seems to be a promising research direction to use TNF- $\alpha$  as a biomarker of the stroke development process or prognosis [142]. The average cost of a TNF assay in the laboratory is around 30 euros. The recognition of TNF-alpha as a potentially valuable biomarker in neurodegenerative diseases is increasing. However, monitoring its levels and applying anti- TNF- $\alpha$  medications have not yet demonstrated significant clinical relevance. Another interesting aspect of TNF research is the importance of this pro-inflammatory cytokine in the aging process. Lindbergh et al. demonstrated that plasma TNF- $\alpha$  concentrations significantly relate to changes in white matter hyperintensities (WMH), gray matter volume (GMV), and global cognition. Authors suggest that systemic inflammation, as indexed by plasma TNF- $\alpha$ , holds potential as a biomarker for age-related

declines in brain health [147]. Inflammation is increasingly recognized as a new modifiable cardiovascular risk factor. As a pro-inflammatory cytokine, TNF can be produced by cardiomyocytes, which typically occurs under unfavorable haemodynamic conditions with excessive pressure or volume overload. It can lead to myocardial metabolic disorders, left ventricular dysfunction, reduced peripheral flow and left ventricular remodelling. The association CAD to subclinical inflammation, quantified by values of circulating markers associated with inflammation remains debatable. There is data supporting associations between circulating hs-CRP and cardiovascular outcomes. However, there is insufficient evidence showing treatments to target inflammation and TNF level results in improvements in patients. Therefore, the determination of plasma TNF levels is not a reliable marker of vascular status to replace the classic radiological examination of the vessels [94, 118, 148].

In conclusion, the basis for assessing the importance of TNF- $\alpha$  signaling in diseases of affluence, including neurodegenerative and cardiovascular diseases, is an understanding of the complex nature of the cytokine in question, as well as the involvement of TNF- $\alpha$  in regulating the release of other factors involved in physiological, inflammatory, or autoimmune processes. As our understanding of TNF- $\alpha$  deepens, we are closer to developing innovative prevention and treatment strategies that take genetic predisposition into account. It is a promising direction of research to use TNF as a biomarker for the development or prognosis of the described diseases. **The conclusions need verification by clinical trials.** To make TNF a reliable marker, the specific role and mechanism it plays, the protective effect and mechanism of anti-TNF- $\alpha$  treatment, and how to reduce the side effects of antibodies are the main issues that need to be further studied and solved by researchers. On the other hand, anti-TNF therapy is also worth exploring as a therapeutic target. The identification of intrinsic cell survival pathways should provide more direct opportunities for trials in neuroprotection or cardiovascular disease.

**Author Contributions:** investigation, conceptualization, literature overview, MB<sup>1</sup>, MB<sup>2</sup>, MER, AM-M; illustrations, DM; manuscript preparation, MB<sup>1</sup>, MB<sup>2</sup>, MER, AM-M, DM. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Lee YB, Nagai A, Kim SU. Cytokines, chemokines, and cytokine receptors in human microglia. *J Neurosci Res.* 2002;69(1):94-103.
2. Brenner D, Blaser H, Mak TW. Regulation of tumour necrosis factor signalling: live or let die. *Nat Rev Immunol.* 2015;15(6):362-374.
3. Banach M, Surma S, Alnouri F, et al. Abstracts From the American Heart Association's 2024 Scientific Sessions and the American Heart Association's 2024 Resuscitation Science Symposium, *Circulation*, 2024;150:4141978.
4. Hehlhans T, Pfeffer K. The intriguing biology of the tumour necrosis factor/tumour necrosis factor receptor superfamily: players, rules and the games. *Immunology.* 2005;115(1):1-20
5. Wu YR, Feng IH, Lyu RK, et al. Tumor necrosis factor- $\alpha$  promoter polymorphism is associated with the risk of Parkinson's disease. *Am J Med Genet B Neuropsychiatr Genet.* 2007;144B(3):300-304.
6. Falvo JV, Tsytsykova AV, Goldfeld AE. Transcriptional control of the TNF gene. *Curr Dir Autoimmun.* 2010;11:27-60.
7. Falvo JV, Lin CH, Tsytsykova AV, Hwang PK, Thanos D, Goldfeld AE, Maniatis T. A dimer-specific function of the transcription factor NFATp. *Proc Natl Acad Sci U S A.* 2008 Dec 16;105(50):19637-42.
8. van Loo G, Bertrand MJM. Death by TNF: a road to inflammation. *Nat Rev Immunol.* 2023 May;23(5):289-303.
9. van der Bruggen T, Nijenhuis S, van Raaij E, Verhoef J, van Asbeck BS. Lipopolysaccharide-induced tumor necrosis factor  $\alpha$  production by human monocytes involves the raf-1/MEK1-MEK2/ERK1-ERK2 pathway. *Infect Immun.* 1999 Aug;67(8):3824-9.
10. Guo YJ, Pan WW, Liu SB, Shen ZF, Xu Y, Hu LL. ERK/MAPK signalling pathway and tumorigenesis. *Exp Ther Med.* 2020 Mar;19(3):1997-2007. doi: 10.3892/etm.2020.8454.
11. Aggarwal BB. Signalling pathways of the TNF superfamily: a double-edged sword. *Nat Rev Immunol.* 2003;3(9):745-756.
12. El-Tahan RR, Ghoneim AM, El-Mashad N. TNF- $\alpha$  gene polymorphisms and expression. *Springerplus.* 2016;5(1):1508.
13. Asgharzadeh M, Najafi-Ghalehlou N, Poor BM, et al. IFN- $\gamma$  and TNF- $\alpha$  Gene Polymorphisms in Multiple Sclerosis Patients in Northwest Iran. *Endocr Metab Immune Disord Drug Targets.* 2021;21(3):520-525.
14. Jabłońska A, Zagrapan B, Neumayer C, et al. Polymorphisms in the IL-6 and TNF- $\alpha$  gene are associated with an increased risk of abdominal aortic aneurysm. *Int J Cardiol.* 2021;329:192-197.
15. Bagheri-Hosseinabadi Z, Mirzaei MR, Aliakbari M, Abbasifard M. Association of interleukin 33 gene polymorphisms with susceptibility and regulation of inflammatory mediators in Systemic lupus erythematosus patients. *Clin Rheumatol.* 2023;42(8):2187-2197.
16. Qidwai T, Khan F. Tumour necrosis factor gene polymorphism and disease prevalence. *Scand J Immunol.* 2011;74(6):522-547.
17. Takahashi H, Yoshimatsu G, Faustman DL. The Roles of TNFR2 Signaling in Cancer Cells and the Tumor Microenvironment and the Potency of TNFR2 Targeted Therapy. *Cells.* 2022;11(12):1952.
18. Bai J, Ding B, Li H. Targeting TNFR2 in Cancer: All Roads Lead to Rome. *Front Immunol.* 2022;13:844931.




19. McMillan D, Martinez-Fleites C, Porter J, et al. Structural insights into the disruption of TNF-TNFR1 signalling by small molecules stabilising a distorted TNF. *Nat Commun.* 2021;12(1):582.
20. Fresegna D, Bullitta S, Musella A, et al. Re-Examining the Role of TNF in MS Pathogenesis and Therapy. *Cells.* 2020;9(10):2290.
21. Kalliolias GD, Ivashkiv LB. TNF biology, pathogenic mechanisms and emerging therapeutic strategies. *Nat Rev Rheumatol.* 2016;12(1):49-62.
22. Dostert C, Grusdat M, Letellier E, Brenner D. The TNF Family of Ligands and Receptors: Communication Modules in the Immune System and Beyond. *Physiol Rev.* 2019;99(1):115-160.
23. Piotrowska A, Izykowska I, Podhorska-Okołów M, Zabel M, Dziegiel P. The structure of NF- $\kappa$ B family proteins and their role in apoptosis. *Postepy Hig Med Dosw (Online).* 2008;62:64-74.
24. Hayden MS, Ghosh S. Regulation of NF- $\kappa$ B by TNF family cytokines. *Semin Immunol.* 2014;26(3):253-266. doi:10.1016/j.smim.2014.05.004
25. Medler J, Kucka K, Wajant H. Tumor Necrosis Factor Receptor 2 (TNFR2): An Emerging Target in Cancer Therapy. *Cancers (Basel).* 2022;14(11):2603.
26. Moatti A, Cohen JL. The TNF- $\alpha$ /TNFR2 Pathway: Targeting a Brake to Release the Anti-tumor Immune Response. *Front Cell Dev Biol.* 2021;9:725473.
27. Naudé PJ, den Boer JA, Luiten PG, Eisel UL. Tumor necrosis factor receptor cross-talk. *FEBS J.* 2011;278(6):888-898.
28. Holbrook J, Lara-Reyna S, Jarosz-Griffiths H, McDermott M. Tumour necrosis factor signalling in health and disease. *F1000Res.* 2019;8:F1000 Faculty Rev-111.
29. Sethi G, Sung B, Aggarwal BB. TNF: a master switch for inflammation to cancer. *Front Biosci.* 2008;13:5094-5107.
30. Leone GM, Mangano K, Petralia MC, Nicoletti F, Fagone P. Past, Present and (Foreseeable) Future of Biological Anti-TNF Alpha Therapy. *J Clin Med.* 2023;12(4):1630.
31. Guo Q, Jin Y, Chen X, et al. NF- $\kappa$ B in biology and targeted therapy: new insights and translational implications. *Signal Transduct Target Ther.* 2024;9(1):53.
32. Zahid M, Busmail A, Penumetcha SS, et al. Tumor Necrosis Factor Alpha Blockade and Multiple Sclerosis: Exploring New Avenues. *Cureus.* 2021;13(10):e18847.
33. Richter F, Williams SK, John K, et al. The TNFR1 Antagonist Atrosimab Is Therapeutic in Mouse Models of Acute and Chronic Inflammation. *Front Immunol.* 2021;12:705485.
34. Gonzalez Caldito N. Role of tumor necrosis factor-alpha in the central nervous system: a focus on autoimmune disorders. *Front Immunol.* 2023;14:1213448.
35. van Oosten BW, Barkhof F, Truyen L, et al. Increased MRI activity and immune activation in two multiple sclerosis patients treated with the monoclonal anti-tumor necrosis factor antibody cA2. *Neurology.* 1996;47(6):1531-1534.
36. Li L, Aviña-Zubieta JA, Bernstein CN, et al. Risk of Multiple Sclerosis Among Users of Antitumor Necrosis Factor  $\alpha$  in 4 Canadian Provinces: A Population-Based Study. *Neurology.* 2023;100(6):e558-e567.
37. Wang L, Deng B, Yan P, Wu H, Li C, Zhu H, Du J, Hou L. Neuroprotective effect of ketamine against TNF- $\alpha$ -induced necroptosis in hippocampal neurons. *J Cell Mol Med.* 2021 Apr;25(7):3449-3459.
38. Mehanna R, Smilowska K, Fleisher J, et al. International Parkinson and Movement Disorder Society Task Force on Early Onset Parkinson's Disease. Age Cutoff for Early-Onset Parkinson's Disease: Recommendations from the International Parkinson and Movement Disorder Society Task Force on Early Onset Parkinson's Disease. *Mov Disord Clin Pract.* 2022 Sep 10;9(7):869-878.
39. Balestrino R, Schapira AHV. Parkinson disease. *Eur J Neurol.* 2020;27(1):27-42.
40. Tansey MG, Wallings RL, Houser MC, Herrick MK, Keating CE, Joers V. Inflammation and immune dysfunction in Parkinson disease. *Nat Rev Immunol.* 2022;22(11):657-673.
41. Pajares M, I Rojo A, Manda G, Boscá L, Cuadrado A. Inflammation in Parkinson's Disease: Mechanisms and Therapeutic Implications. *Cells.* 2020;9(7):1687.
42. Liu TW, Chen CM, Chang KH. Biomarker of Neuroinflammation in Parkinson's Disease. *Int J Mol Sci.* 2022;23(8):4148.
43. Mosley RL, Benner EJ, Kadiu I, et al. Neuroinflammation, Oxidative Stress and the Pathogenesis of Parkinson's Disease. *Clin Neurosci Res.* 2006;6(5):261-281.
44. Mogi M, Kondo T, Mizuno Y, Nagatsu T. p53 protein, interferon-gamma, and NF- $\kappa$ B levels are elevated in the parkinsonian brain. *Neurosci Lett.* 2007;414(1):94-97.
45. Boka G, Anglade P, Wallach D, Javoy-Agid F, Agid Y, Hirsch EC. Immunocytochemical analysis of tumor necrosis factor and its receptors in Parkinson's disease. *Neurosci Lett.* 1994;172(1-2):151-154.
46. Barcia C, de Pablos V, Bautista-Hernández V, et al. Increased plasma levels of TNF-alpha but not of IL1-beta in MPTP-treated monkeys one year after the MPTP administration. *Parkinsonism Relat Disord.* 2005;11(7):435-439.

47. Mogi M, Harada M, Riederer P, Narabayashi H, Fujita K, Nagatsu T. Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) increases both in the brain and in the cerebrospinal fluid from parkinsonian patients. *Neurosci Lett*. 1994;165(1-2):208-210.
48. Hu Y, Yu SY, Zuo LJ, et al. Parkinson disease with REM sleep behavior disorder: features,  $\alpha$ -synuclein, and inflammation. *Neurology*. 2015;84(9):888-894.
49. Joza S, Hu MT, Jung KY, et al. Progression of clinical markers in prodromal Parkinson's disease and dementia with Lewy bodies: a multicentre study. *Brain*. 2023;146(8):3258-3272.
50. Menza M, Dobkin RD, Marin H, et al. The role of inflammatory cytokines in cognition and other non-motor symptoms of Parkinson's disease. *Psychosomatics*. 2010;51(6):474-479.
51. Wang L, Yi H, Liang X, et al. Plasma TNF- $\alpha$  and phosphorylated  $\alpha$ -syn are associated with fatigue in patients with Parkinson's disease. *J Neuroimmunol*. 2023;385:578222.
52. Peter I, Dubinsky M, Bressman S, et al. Anti-Tumor Necrosis Factor Therapy and Incidence of Parkinson Disease Among Patients With Inflammatory Bowel Disease. *JAMA Neurol*. 2018;75(8):939-946.
53. Pascale E, Passarelli E, Purcaro C, et al. Lack of association between IL-1 $\beta$ , TNF- $\alpha$ , and IL-10 gene polymorphisms and sporadic Parkinson's disease in an Italian cohort. *Acta Neurol Scand*. 2011;124(3):176-181.
54. Bialecka M, Klodowska-Duda G, Kurzawski M, et al. Interleukin-10 (IL10) and tumor necrosis factor alpha (TNF) gene polymorphisms in Parkinson's disease patients. *Parkinsonism Relat Disord*. 2008;14(8):636-640.
55. Håkansson A, Westberg L, Nilsson S, et al. Investigation of genes coding for inflammatory components in Parkinson's disease. *Mov Disord*. 2005;20(5):569-573.
56. Nishimura M, Mizuta I, Mizuta E, et al. Tumor necrosis factor gene polymorphisms in patients with sporadic Parkinson's disease. *Neurosci Lett*. 2001;311(1):1-4.
57. Kang X, Ploner A, Pedersen NL, et al. Tumor Necrosis Factor Inhibition and Parkinson Disease: A Mendelian Randomization Study. *Neurology*. 2021;96(12):e1672-e1679.
58. Scheltens P, De Strooper B, Kivipelto M, et al. Alzheimer's disease. *Lancet*. 2021;397(10284):1577-1590.
59. Grobler C, van Tongeren M, Gettemans J, Kell DB, Pretorius E. Alzheimer's Disease: A Systems View Provides a Unifying Explanation of Its Development. *J Alzheimers Dis*. 2023;91(1):43-70.
60. McGeer PL, McGeer EG. The amyloid cascade-inflammatory hypothesis of Alzheimer disease: implications for therapy. *Acta Neuropathol*. 2013 Oct;126(4):479-97.
61. Kaushal N, Ramesh V, Gozal D. TNF- $\alpha$  and temporal changes in sleep architecture in mice exposed to sleep fragmentation. *PLoS One*. 2012;7(9):e45610.
62. Neta R, Sayers TJ, Oppenheim JJ. Relationship of TNF to interleukins. *Immunol Ser*. 1992;56:499-566.
63. Ou W, Yang J, Simanaukaite J, et al. Biologic TNF- $\alpha$  inhibitors reduce microgliosis, neuronal loss, and tau phosphorylation in a transgenic mouse model of tauopathy. *J Neuroinflammation*. 2021;18(1):312.
64. Liao YF, Wang BJ, Cheng HT, Kuo LH, Wolfe MS. Tumor necrosis factor- $\alpha$ , interleukin-1 $\beta$ , and interferon- $\gamma$  stimulate gamma-secretase-mediated cleavage of amyloid precursor protein through a JNK-dependent MAPK pathway. *J Biol Chem*. 2004;279(47):49523-49532.
65. Gratuzze M, Leyns CEG, Holtzman DM. New insights into the role of TREM2 in Alzheimer's disease. *Mol Neurodegener*. 2018;13(1):66.
66. Zhao Y, Wu X, Li X, et al. TREM2 Is a Receptor for  $\beta$ -Amyloid that Mediates Microglial Function. *Neuron*. 2018;97(5):1023-1031.e7.
67. Guerreiro R, Wojtas A, Bras J, et al. TREM2 variants in Alzheimer's disease. *N Engl J Med*. 2013;368(2):117-127.
68. Jonsson T, Stefansson H, Steinberg S, et al. Variant of TREM2 associated with the risk of Alzheimer's disease. *N Engl J Med*. 2013;368(2):107-116.
69. Colonna M. The biology of TREM receptors. *Nat Rev Immunol*. 2023;23(9):580-594.
70. Tarkowski E, Andreasen N, Tarkowski A, Blennow K. Intrathecal inflammation precedes development of Alzheimer's disease. *J Neurol Neurosurg Psychiatry*. 2003;74(9):1200-1205.
71. Fu P, Peng F; Alzheimer's Disease Neuroimaging Initiative. CSF TNF  $\alpha$  levels were associated with conversion from mild cognitive impairment to dementia. *PLoS One*. 2022;17(10):e0274503.
72. Zhou M, Xu R, Kaelber DC, Gurney ME. Tumor Necrosis Factor (TNF) blocking agents are associated with lower risk for Alzheimer's disease in patients with rheumatoid arthritis and psoriasis. *PLoS One*. 2020;15(3):e0229819.
73. Chang R, Yee KL, Sumbria RK. Tumor necrosis factor  $\alpha$  Inhibition for Alzheimer's Disease. *J Cent Nerv Syst Dis*. 2017;9:1179573517709278.
74. Chen YM, Chen HH, Lan JL, Chen DY. Improvement of cognition, a potential benefit of anti-TNF therapy in elderly patients with rheumatoid arthritis. *Joint Bone Spine*. 2010;77(4):366-367.
75. Butchart J, Brook L, Hopkins V, et al. Etanercept in Alzheimer disease: A randomized, placebo-controlled, double-blind, phase 2 trial [published correction appears in *Neurology*. 2015 Dec 8;85(23):2084].
76. Garg N, Smith TW. An update on immunopathogenesis, diagnosis, and treatment of multiple sclerosis. *Brain Behav*. 2015;5(9):e00362.

77. Abbadessa G, Lavorgna L, Treaba CA, Bonavita S, Mainero C. Hemostatic factors in the pathogenesis of neuroinflammation in multiple sclerosis. *Mult Scler.* 2022;28(12):1834-1842.
78. Uysal S, Meriç Yılmaz F, Boğdaycioğlu N, Mungan Öztürk S, Ak F. Increased serum levels of some inflammatory markers in patients with multiple sclerosis. *Minerva Med.* 2014;105(3):229-235.
79. TNF neutralization in MS: results of a randomized, placebo-controlled multicenter study. The Lenercept Multiple Sclerosis Study Group and The University of British Columbia MS/MRI Analysis Group. *Neurology.* 1999;53(3):457-465.
80. van Boxel-Dezaire AH, Hoff SC, van Oosten BW, et al. Decreased interleukin-10 and increased interleukin-12p40 mRNA are associated with disease activity and characterize different disease stages in multiple sclerosis. *Ann Neurol.* 1999;45(6):695-703.
81. Kemanetzoglou E, Andreadou E. CNS Demyelination with TNF- $\alpha$  Blockers. *Curr Neurol Neurosci Rep.* 2017;17(4):36.
82. Bernatsky S, Renoux C, Suissa S. Demyelinating events in rheumatoid arthritis after drug exposures. *Ann Rheum Dis.* 2010;69(9):1691-1693.
83. Kalén E, Piehl F, Andersson M. Demyelinating events following anti-tumor necrosis factor alpha therapy: Rare but challenging to treat. *Eur J Neurol.* 2022;29(7):2047-2055.
84. Xie W, Sun Y, Zhang W, Zhu N, Xiao S. Risk of Inflammatory Central Nervous System Diseases After Tumor Necrosis Factor-Inhibitor Treatment for Autoimmune Diseases: A Systematic Review and Meta-Analysis. *JAMA Neurol.* 2024 Dec 1;81(12):1284-1294.
85. Pauli N, Puchałowicz K, Kuligowska A, et al. Associations between IL-6 and Echo-Parameters in Patients with Early Onset Coronary Artery Disease. *Diagnostics (Basel).* 2019;9(4):189.
86. Wainstein MV, Mossmann M, Araujo GN, et al. Elevated serum interleukin-6 is predictive of coronary artery disease in intermediate risk overweight patients referred for coronary angiography. *Diabetol Metab Syndr.* 2017;9:67.
87. Mourouzis K, Oikonomou E, Siasos G, et al. Pro-inflammatory Cytokines in Acute Coronary Syndromes. *Curr Pharm Des.* 2020;26(36):4624-4647.
88. Vukašinović A, Ostanek B, Klisic A, et al. Telomere-telomerase system status in patients with acute myocardial infarction with ST-segment elevation - relationship with oxidative stress. *Arch Med Sci.* 2021;19(2):313-323.
89. Moller MJ, Qin Z, Toursarkissian B. Tissue markers in human atherosclerotic carotid artery plaque. *Ann Vasc Surg.* 2012;26(8):1160-1165.
90. Akash MSH, Rehman K, Liaqat A. Tumor Necrosis Factor-Alpha: Role in Development of Insulin Resistance and Pathogenesis of Type 2 Diabetes Mellitus. *J Cell Biochem.* 2018;119(1):105-110.
91. Puzkarska A, Niklas A, Głuszek J, Lipski D, Niklas K. The concentration of tumor necrosis factor in the blood serum and in the urine and selected early organ damages in patients with primary systemic arterial hypertension. *Medicine (Baltimore).* 2019;98(22):e15773.
92. Abernethy A, Raza S, Sun JL, et al. Pro-Inflammatory Biomarkers in Stable Versus Acutely Decompensated Heart Failure With Preserved Ejection Fraction. *J Am Heart Assoc.* 2018;7(8):e007385.
93. Li JY, He Y, Ke HH, Jin Y, Jiang ZY, Zhong GQ. Plasma oxidative stress and inflammatory biomarkers are associated with the sizes of the left atrium and pulmonary vein in atrial fibrillation patients. *Clin Cardiol.* 2017;40(2):89-94.
94. Białecka M, Dziejewski V, Safranow K, et al. Could Tumor Necrosis Factor Serve as a Marker for Cardiovascular Risk Factors and Left Ventricular Hypertrophy in Patients with Early-Onset Coronary Artery Disease?. *Diagnostics (Basel).* 2024;14(4):449.
95. Solaro RJ, Rosas PC, Langa P, Warren CM, Wolska BM, Goldspink PH. Mechanisms of troponin release into serum in cardiac injury associated with COVID-19 patients. *Int J Cardiol Cardiovasc Dis.* 2021;1(2):41-47.
96. Sordillo PP, Sordillo DC, Helson L. The Prolonged QT Interval: Role of Pro-inflammatory Cytokines, Reactive Oxygen Species and the Ceramide and Sphingosine-1 Phosphate Pathways. *In Vivo.* 2015;29(6):619-636.
97. González M, Ruiz-Ros JA, Pérez-Paredes M, et al. Prognostic value of tumor necrosis factor-alpha in patients with ST-segment elevation acute myocardial infarction. *Rev Esp Cardiol.* 2007;60(12):1233-1241.
98. Medenwald D, Swenne CA, Loppnow H, et al. Prognostic relevance of the interaction between short-term, metronome-paced heart rate variability, and inflammation: results from the population-based CARLA cohort study. *Europace.* 2017;19(1):110-118.
99. Galeone A, Grano M, Brunetti G. Tumor Necrosis Factor Family Members and Myocardial Ischemia-Reperfusion Injury: State of the Art and Therapeutic Implications. *Int J Mol Sci.* 2023;24(5):4606.
100. Attiq A, Afzal S, Ahmad W, Kandeel M. Hegemony of inflammation in atherosclerosis and coronary artery disease. *Eur J Pharmacol.* 2024;966:176338.
101. Toczyłowski K, Hirnle T, Harasiuk D, et al. Plasma concentration and expression of adipokines in epicardial and subcutaneous adipose tissue are associated with impaired left ventricular filling pattern. *J Transl Med.* 2019;17(1):310.

102. Karampetsou N, Tzani A, Doulamis IP, et al. Epicardial Adipocyte-derived TNF- $\alpha$  Modulates Local Inflammation in Patients with Advanced Coronary Artery Disease. *Curr Vasc Pharmacol*. 2022;20(1):87-93.
103. Mirhafez SR, Khadem SH, Sahebkar A, et al. Comparative effects of on-pump versus off-pump coronary artery bypass grafting surgery on serum cytokine and chemokine levels. *IUBMB Life*. 2021;73(12):1423-1431.
104. Jortveit J, Pripp AH, Langørgen J, Halvorsen S. Incidence, risk factors and outcome of young patients with myocardial infarction. *Heart*. 2020;106(18):1420-1426.
105. Liu M, Zhang P, Chen M, et al. Aging might increase myocardial ischemia / reperfusion-induced apoptosis in humans and rats. *Age (Dordr)*. 2012;34(3):621-632.
106. Charakida M, Halcox JP. Tumor necrosis factor-alpha in heart failure: more questions than answers. *Rev Esp Cardiol*. 2005;58(5):470-472.
107. Suzuki T, Ogita H, Sato A, Minamide N, Hachiro K. Differences Between Patients with and without Atherosclerosis in Expression Levels of Inflammatory Mediators in the Adipose Tissue Around the Coronary Artery. *Int Heart J*. 2021;62(2):390-395.
108. Zhang Y, Yang X, Bian F, et al. TNF- $\alpha$  promotes early atherosclerosis by increasing transcytosis of LDL across endothelial cells: crosstalk between NF- $\kappa$ B and PPAR- $\gamma$ . *J Mol Cell Cardiol*. 2014;72:85-94.
109. Franco AG, Sandri S, Campa A. High-density lipoprotein prevents SAA-induced production of TNF- $\alpha$  in THP-1 monocytic cells and peripheral blood mononuclear cells. *Mem Inst Oswaldo Cruz*. 2011;106(8):986-992.
110. Patel VK, Williams H, Li SCH, Fletcher JP, Medbury HJ. Monocyte inflammatory profile is specific for individuals and associated with altered blood lipid levels. *Atherosclerosis*. 2017;263:15-23.
111. Dong H, Wang J, Hu P, Lu N. Association of Apolipoprotein A1, High Density Lipoprotein Cholesterol, and Their Ratio with Inflammatory Marker in Chinese Adults with Coronary Artery Disease. *Angiology*. 2023;74(8):765-773.
112. Abbasifard M, Kandelouei T, Aslani S, et al. Effect of statins on the plasma/serum levels of inflammatory markers in patients with cardiovascular disease; a systematic review and meta-analysis of randomized clinical trials. *Inflammopharmacology*. 2022;30(2):369-383.
113. Vangaveti V, Shashidhar V, Collier F, Hodge J, Rush C, Malabu U, Baune B, Kennedy RL. 9- and 13-HODE regulate fatty acid binding protein-4 in human macrophages, but does not involve HODE/GPR132 axis in PPAR- $\gamma$  regulation of FABP4. *Ther Adv Endocrinol Metab*. 2018 May;9(5):137-150.
114. Gilani STA, Khan DA, Rauf A, Haroon ZH, Khan KA, Hassan FU. Early Diagnosis of Coronary Artery Disease by Inflammatory Biomarkers of Atherosclerosis in Patients with Angina. *J Interferon Cytokine Res*. 2022;42(9):493-500.
115. Dong L, Hou R, Xu Y, et al. Analyzing the Correlation between the Level of Serum Markers and Ischemic Cerebral Vascular Disease by Multiple Parameters. *Comput Math Methods Med*. 2015;2015:972851.
116. Elkind MS, Cheng J, Boden-Albala B, et al. Tumor necrosis factor receptor levels are associated with carotid atherosclerosis. *Stroke*. 2002;33(1):31-37.
117. Andersson J, Sundström J, Kurland L, et al. The carotid artery plaque size and echogenicity are related to different cardiovascular risk factors in the elderly: the Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS) study. *Lipids*. 2009;44(5):397-403.
118. Bialecka M, Rac M, Dziedziejko V, Safranow K, Chlubek D, Rać ME. An Evaluation of Plasma TNF, VEGF-A, and IL-6 Determination as a Risk Marker of Atherosclerotic Vascular Damage in Early-Onset CAD Patients. *J Clin Med*. 2024;13(6):1742.
119. Cortez-Cooper M, Meaders E, Stallings J, et al. Soluble TNF and IL-6 receptors: indicators of vascular health in women without cardiovascular disease. *Vasc Med*. 2013;18(5):282-289.
120. Zhang H, Jiang M, Hou H, Li Q. Efficacy of simvastatin on carotid atherosclerotic plaque and its effects on serum inflammatory factors and cardiocerebrovascular events in elderly patients. *Exp Ther Med*. 2021;22(2):819.
121. Chen Y, Wang X, Yang C, et al. Decreased circulating catestatin levels are associated with coronary artery disease: The emerging anti-inflammatory role. *Atherosclerosis*. 2019;281:78-88.
122. Nedeljkovic M, Mihajlovic M, Mujovic N, et al. Assessment of patient-reported treatment burden in patients with coronary artery disease. *Arch Med Sci*. 2024;20(3):1006-1010.
123. Dimitroglou Y, Aggeli C, Theofilis P, Tsioufis P, et al. Novel Anti-Inflammatory Therapies in Coronary Artery Disease and Acute Coronary Syndromes. *Life (Basel)*. 2023;13(8):1669.
124. Ogrič M, Poljšak KM, Lakota K, et al. Neutralizing effects of anti-infliximab antibodies on synergistically-stimulated human coronary artery endothelial cells. *Atherosclerosis*. 2019 291:1-8.
125. Barbati C, Colasanti T, Vomero M, et al. Up-regulation of autophagy by etanercept treatment results in TNF-induced apoptosis reduction in EA.hy926 endothelial cell line. *Clin Exp Rheumatol*. 2021;39(3):606-611.
126. Mazurek A, Gryga K, Bugala K, et al. Influence of statin therapy on antiphospholipid antibodies and endothelial dysfunction in young patients with coronary artery disease and systemic lupus erythematosus. *Arch Med Sci*. 2020;18(1):18-24.

127. Bytyçi I, Bajraktari G, Penson PE, et al. Efficacy and safety of colchicine in patients with coronary artery disease: A systematic review and meta-analysis of randomized controlled trials. *Br J Clin Pharmacol.* 2022;88(4):1520-1528.
128. Ruscica M, Corsini A, Ferri N, et al. Clinical approach to the inflammatory etiology of cardiovascular diseases. *Pharmacol Res.* 2020;159:104916.
129. Landmesser U, Haghikia A, Leiter LA, et al. Effect of inclisiran, the small-interfering RNA against proprotein convertase subtilisin/kexin type 9, on platelets, immune cells, and immunological biomarkers: a pre-specified analysis from ORION-1. *Cardiovasc Res.* 2021;117(1):284-291.
130. Rakocevic J, Dobric M, Borovic ML, et al. Anti-Inflammatory Therapy in Coronary Artery Disease: Where Do We Stand? *Rev Cardiovasc Med.* 2023;24(1):10.
131. Maciejczyk M, Bielas M, Zalewska A, Gerreth K. Salivary Biomarkers of Oxidative Stress and Inflammation in Stroke Patients: From Basic Research to Clinical Practice. *Oxid Med Cell Longev.* 2021;2021:5545330.
132. Sena CM, Leandro A, Azul L, et al. Vascular Oxidative Stress: Impact and Therapeutic Approaches. *Front Physiol.* 2018;9:1668.
133. del Zoppo GJ. Stroke and neurovascular protection. *N Engl J Med.* 2006;354(6):553-555.
134. Dirnagl U, Iadecola C, Moskowitz MA. Pathobiology of ischaemic stroke: an integrated view. *Trends Neurosci.* 1999;22(9):391-397.
135. Zhou X, Yu F, Feng X, et al. Immunity and inflammation predictors for short-term outcome of stroke in young adults. *Int J Neurosci.* 2018;128(7):634-639.
136. Zaremba J, Losy J. Early TNF-alpha levels correlate with ischaemic stroke severity. *Acta Neurol Scand.* 2001;104(5):288-295.
137. Badiola N, Malagelada C, Llecha N, et al. Activation of caspase-8 by tumour necrosis factor receptor 1 is necessary for caspase-3 activation and apoptosis in oxygen-glucose deprived cultured cortical cells. *Neurobiol Dis.* 2009;35(3):438-447.
138. Yong VW, Power C, Forsyth P, Edwards DR. Metalloproteinases in biology and pathology of the nervous system. *Nat Rev Neurosci.* 2001;2(7):502-511.
139. Doll DN, Barr TL, Simpkins JW. Cytokines: their role in stroke and potential use as biomarkers and therapeutic targets. *Aging Dis.* 2014;5(5):294-306.
140. Liang M, Zhang L, Geng Z. Advances in the Development of Biomarkers for Poststroke Epilepsy. *Biomed Res Int.* 2021;2021:5567046.
141. Chen Y, Pu J, Liu Y, et al. Pro-inflammatory cytokines are associated with the development of post-stroke depression in the acute stage of stroke: A meta-analysis. *Top Stroke Rehabil.* 2020;27(8):620-629.
142. Xue Y, Zeng X, Tu WJ, Zhao J. Tumor Necrosis Factor- $\alpha$ : The Next Marker of Stroke. *Dis Markers.* 2022;2022:2395269.
143. Arango-Dávila CA, Vera A, Londoño AC, et al. Soluble or soluble/membrane TNF- $\alpha$  inhibitors protect the brain from focal ischemic injury in rats. *Int J Neurosci.* 2015;125(12):936-940.
144. Frankola KA, Greig NH, Luo W, Tweedie D. Targeting TNF- $\alpha$  to elucidate and ameliorate neuroinflammation in neurodegenerative diseases. *CNS Neurol Disord Drug Targets.* 2011;10(3):391-403.
145. Gerriets V, Goyal A, Khaddour K. Tumor Necrosis Factor Inhibitors. In: *StatPearls.* Treasure Island (FL): StatPearls Publishing; July 3, 2023.
146. Wang X, Feuerstein GZ, Xu L, et al. Inhibition of tumor necrosis factor-alpha-converting enzyme by a selective antagonist protects brain from focal ischemic injury in rats. *Mol Pharmacol.* 2004;65(4):890-896.
147. Lindbergh CA, Casaletto KB, Staffaroni AM, et al. Systemic Tumor Necrosis Factor-Alpha Trajectories Relate to Brain Health in Typically Aging Older Adults. *J Gerontol A Biol Sci Med Sci.* 2020;75(8):1558-1565.
148. Maiorean S, Webb R, Banach M, Mazidi M. The role of inflammation and the possibilities of inflammation reduction to prevent cardiovascular events. *Eur Heart J Open.* 2022;2(4):oeac039.



Preprint