

2024: The year in cardiovascular disease – the year of lipoprotein(a). Research advances and new findings

Bożena Sosnowska^{1*}, Peter P. Toth², Alexander C. Razavi³, Alan T. Remaley⁴, Roger S. Blumenthal⁵, Maciej Banach^{1,6,7} on behalf of the International Lipid Expert Panel (ILEP)

¹Department of Preventive Cardiology and Lipidology, Medical University of Lodz (MUL), Lodz, Poland

²CGH Medical Center, Sterling, IL, and Ciccarone Center for the Prevention of Cardiovascular Disease, Johns Hopkins University School of Medicine, Baltimore, MD, USA

³Emory Clinical Cardiovascular Research Institute, Emory University School of Medicine, Atlanta, GA, USA

⁴Lipoprotein Metabolism Laboratory, Translational Vascular Medicine Branch, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD, USA

⁵Ciccarone Center for the Prevention of Cardiovascular Disease, Johns Hopkins University School of Medicine, Baltimore, MD, USA

⁶Faculty of Medicine, the John Paul II Catholic University of Lublin (KUL), Lublin, Poland

⁷Department of Cardiology and Adult Congenital Heart Diseases, Polish Mother's Memorial Hospital Research Institute (PMMHRI), Lodz, Poland

***Corresponding author:**

Dr Bożena Sosnowska
Department of Preventive Cardiology and Lipidology
Medical University of Lodz
Rzgowska 281/289
93-338 Lodz, Poland
Phone: +48 42 2711124
E-mail: bozena.sosnowska@umed.lodz.pl

Submitted: 7 December 2024; **Accepted:** 20 February 2025

Online publication: 22 February 2025

Arch Med Sci 2025; 21 (2): 355–373

DOI: <https://doi.org/10.5114/aoms/202213>

Copyright © 2025 Termedia & Banach

Abstract

Elevated plasma lipoprotein(a) [Lp(a)] levels, which occur in as many as 1.5 billion people worldwide, are an independent and causal risk factor for atherosclerotic cardiovascular disease and calcific aortic valve disease. Unlike low-density lipoprotein cholesterol, Lp(a) levels are approximately 70–90% genetically determined. Currently, no approved pharmacological therapies specifically target lowering Lp(a) concentrations. Several drugs, mainly RNA-based therapies, that specifically and potently lower Lp(a), are under investigation. Three of these new therapeutic agents are advancing through clinical development to evaluate whether reducing Lp(a) levels can decrease cardiovascular risk. The outcomes of these trials could potentially transform cardiovascular disease prevention strategies; however, once approved, the drugs will likely be used for secondary prevention, and ongoing strategies for managing elevated Lp(a) in primary prevention will be important. Lipoprotein(a) research is a rapidly evolving field, but unanswered questions remain concerning the physiological function of Lp(a) and its true pathogenic mechanisms. This review of Lp(a) research focuses on new findings and clinical trial results that appeared in 2024.

Key words: aortic valve calcification, atherosclerosis, cardiovascular disease, lipoprotein(a), apo(a) isoforms, RNA therapeutics, outcomes, variability.

Introduction

Elevated lipoprotein(a) [Lp(a)] accounts for a significant component of residual cardiovascular disease risk (CVD) [1]. Globally, elevated Lp(a) levels (≥ 50 mg/dl or ≥ 125 nmol/l) are estimated to affect over 1.5 billion

people [2]. The contribution of Lp(a) to cardiovascular risk varies, ranging from 13.5% in relatively healthy patients to 20% in patients at high and very high CVD risk, which suggests that Lp(a) may contribute to the baseline risk [3–5]. Individuals with the highest Lp(a) levels face as much as a 31% increased risk of CVD and a 42% greater likelihood of experiencing an atherosclerotic cardiovascular disease (ASCVD) event [6]. Moreover, Lp(a) concentrations exceeding 180 mg/dl are associated with a cardiovascular risk comparable to that in patients with heterozygous familial hypercholesterolemia (heFH) [7].

Apolipoprotein B100 (apoB), which is present as a single copy per particle in both Lp(a) and low-density lipoprotein (LDL), has been used to directly compare the relative atherogenicity of these two kinds of lipoproteins [1]. In the UK Biobank cohort, it was found that the Lp(a) particle has a more than 6-fold stronger association with CVD risk than the LDL particle [8]. The results of this genetic analysis were similar to the epidemiological study of Marston *et al.* [9]. It is important to note, however, that in most people LDL particles are present in greater amounts than Lp(a) particles; thus, the overall CVD risk associated with LDL is higher [10]. Another important issue is that exposure to high levels of Lp(a) starts in very early life – according to the Baby Copenhagen substudy, as early as 15 months of age [11]. On the other hand, Lp(a) appears to play a more pronounced role in promoting atheroma development and progression, particularly during the advanced stages of the disease, whereas LDL-C contributes consistently throughout the entire time course of atherogenesis [12, 13].

Unlike LDL-C, Lp(a) levels are not generally modifiable through lifestyle changes (although there is ongoing discussion on Lp(a) level variability), as they are primarily determined by genetic factors [14]. Thus, developing novel therapies is crucial for addressing residual cardiovascular risk, with Lp(a) emerging as a key target for advancing cardiovascular disease prevention. Earlier estimates indicated that lowering Lp(a) by more than 100 mg/dl was necessary to reduce the associated risk of CVD [15]. More recent analyses suggest that a reduction of just 65.7 mg/dl may achieve a comparable decrease in CVD risk to that seen with a 38.7 mg/dl reduction in LDL-C [16].

This review highlights the physiology and pathophysiology of Lp(a), its association with cardiovascular risk, challenges in its measurement, and emerging therapeutics aimed at lowering Lp(a) levels, with a particular focus on the latest findings from 2024. We also explore some of long lingering questions about Lp(a) and its role in cardiovascular disease.

Lipoprotein(A) structure, physiology, and pathophysiology

Lipoprotein(a) is similar to an LDL particle, but it has apolipoprotein(a) [apo(a)] attached to it by both covalent and non-covalent bonding. Apo(a) is a large protein that is primarily synthesized in the liver [17]. A non-covalent bond attaches apo(a) kringle IV domains 7 and 8 to apoB lysine residues; in addition, apo(a) is attached covalently to apoB via a cysteine residue (Cys4057) in the KIV9 domain [18]. Apo(a) is structurally similar to plasminogen, a protein involved in fibrinolysis. Due to this structural similarity, apo(a) may compete with plasminogen when binding fibrin to inhibit fibrinolysis and increase thrombotic risk [18–20]. Plasminogen has 5 kringle domains (KI, KII, KIII, KIV, and KV) and one protease domain at the end, whereas apo(a) has only 10 KIV subtypes (KIV1–KIV10), KV, and a non-reactive serine protease-like domain [19, 20].

Various copies of KIV2 lead to size polymorphism of apo(a), resulting in different levels of Lp(a) in the plasma. Apo(a) isoforms contain between 3 and over 50 KIV2 repeats and have polypeptide molecular masses between ~200 and ~800 kDa [19, 20]. A low number of KIV2 copies (≤ 22) leads to the production of small apo(a) isoforms, which is associated with higher Lp(a) concentrations compared to large apo(a) isoforms (> 22 KIV2 repeats) [21]. When the number of KIV2 repeats is high, a significant portion of apo(a) molecules undergo degradation within hepatocytes before secretion. In contrast, with a low number of KIV2 repeats, the molecules are efficiently secreted and bind to LDL particles outside hepatocytes to form Lp(a) [22]; this may help to understand why Lp(a) is the fifth most prevalent CVD risk factor.

In addition to apo(a) isoform size, which is encoded by the *LPA* gene, other genetic variants, such as single nucleotide polymorphisms (SNPs), also determine Lp(a) serum levels [23, 24]. Over 2000 SNPs genome-wide are significantly associated with Lp(a) concentrations [23]. Certain SNPs, such as rs1800769 and rs1853021, are associated with lower Lp(a) levels, whereas others, such as rs10455872 and rs3798220, are usually observed with small apo(a) isoforms and are associated with elevated Lp(a) levels [24].

Besides carrying cholesterol esters, free cholesterol, triglycerides, and phospholipids, Lp(a) is the main carrier of oxidized phospholipids (OxPL) among all the apoB-containing lipoproteins [25]. The physiological role of Lp(a) in the body is not fully understood, and its function remains a subject of intense research [26]. However, there are hypotheses regarding its possible roles. Since Lp(a) has no recognized physiological function, it may be associated with a reduced lifespan; this, however, still needs to be confirmed [26].

The high homology between apo(a) and plasminogen (75–99%) suggests a possible role for Lp(a) in fibrinolysis [27]. The latest *ex vivo* and some human studies do not support the view that Lp(a) is an antifibrinolytic factor [28]. In contrast, *in vitro* studies have shown that apo(a) potentially affects fibrinolysis in several ways. Apo(a) inhibits (1) plasminogen activation by tissue plasminogen activator (tPA) and (2) the binding of plasminogen to fibrin surfaces. Apo(a) interrupts plasmin-mediated conversion of native Glu1-plasminogen to Lys77-plasminogen. Lys77-plasminogen is a better substrate for tPA, so it is more efficiently converted to plasmin. Inhibition of this conversion by apo(a) leads to a reduction in the amount of plasmin and a weakening of fibrinolysis [27].

The inhibition of fibrinolysis improves wound healing, supporting the theory that Lp(a) may have evolved to have a protective role in primates for certain conditions, such as injuries and infections, particularly during early life [2, 29]. The potential role of Lp(a) in wound healing aligns with immunohistochemical analyses that demonstrated positive staining for apo(a)/apoB in wounds at various stages of healing [30, 31]. Furthermore, a proteomics study identified a correlation between Lp(a) and numerous proteins involved in wound healing [30, 32].

The pathophysiological mechanisms by which Lp(a) contributes to cardiovascular disease risk may involve several interconnected pathways. These include the proatherogenic effects mediated by apoB [33], a pro-inflammatory response driven by OxPL [34], and pro-thrombotic effects resulting from the antifibrinolytic properties of apo(a) [27] and its interactions with platelets [35]. Specifically, processes such as endothelial dysfunction, inflammation, lipid accumulation, and calcification – partly driven by oxidized phospholipids present on the Lp(a) particle – may play a central role in the development of ASCVD and calcific aortic valve stenosis (CAVS) [36, 37].

Lp(a) measurement, nmol/l vs. mg/dl unit

Measurement of Lp(a) is challenging mainly due to the different apo(a) isoforms resulting from the variable number of KIV2 repeats, which can alter its quantification by immunoassays [27]. Additionally, every individual typically inherits and expresses two copies of the *LPA* gene, one from each parent. Consequently, unless an individual is homozygous for two *LPA* genes with the same KIV2 repeat count, most people possess two distinct apo(a) isoforms, with their levels representing the combined contributions of both apo(a) isoform sizes [38]. Accurate Lp(a) measurement methods are crucial for properly assessing the impact of Lp(a) on various CVDs and for advancing

the clinical development of therapies targeting Lp(a). The optimal diagnostic test for Lp(a) should remain unaffected by Lp(a) isoform variability, specifically detect Lp(a) particles, provide results in nanomoles per liter (nmol/l), and rely on standards traceable to globally recognized reference materials [39].

While numerous methods, including immunoassays, fluorescence-based techniques, and electrophoresis, have been developed to measure Lp(a), achieving standardization has proven difficult due to variations in antibody reactivity to different Lp(a) phenotypes [40]. Heydari *et al.* [40] did not find the assay to be 100% insensitive to apo(a) size. Tests exhibit varying degrees of sensitivity to apo(a) isoforms, which can result in the measured Lp(a) concentration being underestimated in the presence of small isoforms or overestimated when large isoforms are present. However, those assays using 5 to 6 independent calibrators, covering a wide range of Lp(a) levels and a balanced distribution of apo(a) isoforms, are minimally influenced by apo(a) size [36, 41, 42].

The new challenge for Lp(a) measurement arises as a result of the development of muvalaplin – new drug targeting Lp(a) that disrupts the initial non-covalent interaction between apo(a) and apoB100, preventing formation of the disulfide bond and leading to an increase in free apo(a) [43]. Current commercial Lp(a) assays that measure total apo(a) may be insufficient to accurately measure Lp(a) concentrations in patients on this drug; thus, a novel immunoassay that measures only intact Lp(a) particles has been developed [43]. It has been confirmed that this new test is insensitive to apo(a) isoform size and correlates with a liquid chromatography–tandem mass spectrometry method [43].

Measuring Lp(a) in molar units is optimal but poses considerable difficulties. Measuring in mass units (mg/dl) is equally effective, however, for clinical applications [44, 45]. Converting the unit from mg/dl to nmol/l is not recommended, as there is no consistent conversion factor between mass and molar scales due to the varying isoform dependency of each immunoassay-based analytical method [45, 46]. Although the available assays are not yet perfect, most of them can be used for risk stratification of patients [44]. Despite the limitations of mass-based measurement, Lp(a) testing with the most readily available assay (mass- or particle-based) is favored over no testing to facilitate CVD risk stratification, especially in persons with high baseline risk.

This is reinforced by a sub-analysis of the ODYSSEY Outcome trial (Evaluation of Cardiovascular Outcomes After an Acute Coronary Syndrome During Treatment With Alirocumab), which

Table I. The most burning questions on lipoprotein(a) based on research results published in 2023–2024

No.	Burning question	Current state of knowledge
1.	Does lipoprotein(a) have any physiological function in humans?	There is lack of data for humans or data are inconsistent (role in wound healing?). It might be one of the factors associated with a reduction of lifespan.
2.	What should be the best test for Lp(a) measurement?	Lp(a) measurement is challenging mainly due to the different apo(a) isoforms depending on the number of KIV2 repeats. The optimal diagnostic test should remain unaffected by Lp(a) isoform variability, specifically detect Lp(a) particles, provide results in nanomoles per liter (nmol/L), and rely on standards traceable to globally recognized reference materials.
3.	Should lipoprotein(a) be measured in children and adolescents?	The Polish 2024 guidelines, as well as the updated NLA recommendations on the use of Lp(a) in clinical practice, recommended screening of Lp(a) in children < 18 years of age in certain cases. Guidelines also recommend cascade screening of immediate family members of a child or adult with elevated Lp(a).
4.	How often should we measure lipoprotein(a)?	For most patients it is enough to have one Lp(a) measurement, especially in those with normal values. However, in the case of intermediate and elevated Lp(a) levels, in those with risk factors and conditions that may affect Lp(a) levels, in those with a previous test in mg/dl, and those with visit-to-visit variability, more than one measurement should be considered.
5.	How to effectively increase awareness on the role of Lp(a) in CVD risk and the number of patients with Lp(a) measurement?	Lipoprotein(a) is the 4 th most prevalent risk factor (after lipid disorders, hypertension, and smoking), and elevated levels are observed in 15–25% of the population (depending on baseline CVD risk). However, knowledge about this among physicians and especially patients is very low. Therefore, the frequency of Lp(a) measurement remains low, just a few percent.
6.	What are the non-genetic risk factors that may affect Lp(a) level?	There is still an ongoing discussion on the non-genetic risk factors and conditions (including drugs) that may significantly affect Lp(a) levels. It is, among other things, related to the fact that in a relatively high percentage of patients, Lp(a) level variability is observed without any recognized variables that may have influenced this.
7.	What is the role of apo(a) isoforms?	The apo(a) isoforms affect the Lp(a) level and response to some lipid-lowering drugs (statins, niacin, PCSK9 inhibitors). However, we still do not know their exact role in the response to some risk factors/conditions that may affect Lp(a) level, visit-to-visit Lp(a) variability, or CVD risk.
8.	Are patients from the gray zone those at increased CVD risk?	Although Mendelian randomization studies suggest that the risk increases starting from the levels of Lp(a) over 30 mg/dl (75 nmol/l), imaging data on atherosclerosis progression and outcomes data still give inconsistent results in comparison to patients with higher Lp(a) levels (> 50 mg/dl).
9.	Is Lp(a) a significant CVD risk factor for all patients?	From the point of view of clinical practice, the question constantly arises: how is it possible that we encounter patients aged 65 and over with elevated Lp(a) levels without significant signs of atherosclerosis? Is this related to a healthy lifestyle and high adherence to therapy against established CVD risk factors, or are there any protective factors that may reduce the harmful effects of elevated Lp(a) that we have yet to recognize?
10.	Should patients with elevated Lp(a) be treated with aspirin?	Available data recently published indicate that in patients with elevated Lp(a) > 50 mg/dl, low-dose aspirin might significantly reduce CVD outcomes without significantly increasing the risk of major bleeding events. The new PoLA Expert Opinion paper recommends low-dose aspirin in high-risk patients in primary prevention with Lp(a) > 50 mg/dl. Further confirmation of the clinical net benefit of this intervention is needed.
11.	Will new targeted therapies that significantly reduce Lp(a) by over 70% also reduce cardiovascular outcomes and mortality?	This is a critical question on the role of future targeted Lp(a) therapies, because the data from HORIZON-Lp(a) with pelacarsen will shed light on how a significant Lp(a) reduction will affect cardiovascular outcomes and mortality reduction. This will help to position these drugs alongside other cardiovascular therapies to significantly reduce Lp(a)-related residual CVD risk.
12.	Are low and extremely low Lp(a) levels safe?	Observational studies have suggested that extremely low Lp(a) levels might be associated with an elevated risk of type 2 diabetes mellitus. MR analyses did not confirm this association, indicating the link with hyperinsulinemia, but without reverse causality. Thus, aggressive Lp(a)-lowering therapy does not substantiate any concerns about exacerbated T2DM risk, although this still needs to be confirmed in long-term studies with new targeted therapies.

highlighted that mass and molar lipoprotein(a) immunoassays were similar in their prognostic risk for a major adverse cardiovascular event (MACE) or for MACE reduction with alirocumab. The study included the Siemens N-latex nephelometric immunoassay (measured in mass units – mg/dl), the Roche Tina-Quant turbidimetric immunoassay (measured in molar units – nmol/l), and a non-commercial mass spectrometry (nmol/l) assay [47] (Table I).

Lp(a) concentrations

Plasma Lp(a) levels are approximately 90% genetically determined through the *LPA* gene [14]. Lipoprotein(a) levels are low at birth, and measurements from umbilical cord blood can be a reliable indicator of neonatal venous blood levels. By 15 months of age, Lp(a) levels reach those typically observed in adults, and the 90th percentile of levels measured at birth (whether from cord or venous blood) serves as a strong predictor for later high-risk levels [11].

Lp(a) concentrations are significantly influenced by race and gender, though the proportional CVD risk due to Lp(a) is roughly similar across ethnicities after considering traditional risk factors [48]. Lower Lp(a) concentrations are found in Chinese, Caucasians, and South Asians, and the highest in Black individuals [2]. The level of Lp(a) is about 17% higher in women than in men after age 50, which is usually associated with the onset of menopause [49]. In women, Lp(a) levels are more variable during their lifetime [50]. Lp(a) levels remain stable from menarche through the reproductive years until the perimenopause. However, during pregnancy, Lp(a) levels approximately double, though the exact mechanisms behind this increase are not understood [51]. It is hypothesized that estrogen may influence Lp(a) synthesis and clearance and that Lp(a) could act as an acute-phase protein in response to endothelial damage, or that it may play a role in placental development [52, 53]. Elevated Lp(a) levels during pregnancy might impact outcomes, potentially raising the risk of complications, such as gestational diabetes, preterm delivery, and low birth weight [54]. Recent data did not confirm the causal relationship between Lp(a) levels and the risk of preeclampsia [55].

Growing evidence has emerged regarding environmental risk factors, conditions, and therapies that can influence Lp(a) levels, potentially contributing to the observed individual variability in Lp(a) [56–58]. Among patients in the placebo group in the OCEAN(a)-DOSE Trial with stable ASCVD and elevated baseline Lp(a) concentrations, notable intraindividual variability of about 10% in Lp(a) levels was observed across all visits [59]. Data

from the Nashville Biosciences database showed that baseline and follow-up paired values were significantly different, with an absolute change of ≥ 10 mg/dl in 38.1% and a $> 25\%$ change in 40.5% of individuals. Black individuals exhibited greater variability than White individuals; likewise, women exhibited greater variability than men. A positive correlation between the baseline Lp(a) levels and the absolute changes in Lp(a) was also observed; 53% of those in the intermediate “gray zone” category transitioned to either the low-risk (20%) or high-risk (33%) category [60]. In the PMMHRI-Lp(a) Registry, the Lp(a) visit-to-visit variability (mean time distance: 7 ± 5 months) was insignificant and only 3.25 mg/dl, but as many as every fourth patient had a difference greater than 10 mg/dl [3]. This issue needs to be further investigated, including the effect of apo(a) isoforms on the level’s variability. This may provide invaluable information for identifying patients who should have Lp(a) measurement more than once [56].

Lp(a) is an acute phase reactant, and is often elevated in conditions such as sepsis, post-surgical states, viral infections, and myocardial infarction [61, 62]. However, the role of Lp(a) as an acute phase reactant remains a subject of ongoing debate [63]. A prospective observational study examined changes in Lp(a) levels in individuals with ST-elevation myocardial infarction (STEMI) at four time points. Median Lp(a) levels increased from 7.9 mg/dl upon hospital admission to 8.4 mg/dl the following day, then to 9.3 mg/dl on the second day ($p < 0.001$), and further increased to 11.2 mg/dl at 3 months after MI ($p < 0.001$) [64]. Another study obtained similar results: Lp(a) levels in patients with acute myocardial infarction (AMI) were significantly elevated 6 months after the event, with an increase of at least 25 nmol/l (~ 10 mg/dl) observed in over 20% of participants. This pattern contrasted with that of high-sensitivity C-reactive protein (hsCRP), suggesting that Lp(a) does not act as a conventional acute phase reactant during AMI [64]. Both studies, however, lack baseline Lp(a) data prior to the occurrence of AMI [64, 65]. This may suggest that repeat testing of Lp(a) after MI should be performed. In total, these findings suggest that repeated measurement of Lp(a) after AMI may be warranted. This concept may be similar to the change in LDL levels after AMI, which in contrast are approximately 10% lower in the peri- and post-MI period [66].

The European Atherosclerosis Society (EAS) consensus statement, the National Lipid Association (NLA), and cardiovascular prevention guidelines from France, Poland, Italy, Canada, India, and China only advise measuring Lp(a) once in a lifetime [45, 56, 67–72]. Polish guidelines do suggest repeating Lp(a) measurements in those patients

whose initial Lp(a) levels fall within 30–50 mg/dl (75–125 nmol/l), a range considered the “gray zone” or near the threshold for cardiovascular risk categories. Repeated measurements may be considered for women over the age of 50 years and for patients with chronic kidney disease, particularly nephrotic syndrome, since these conditions can significantly elevate Lp(a) levels [56]. The Polish guidelines [56], as well as the updated NLA recommendations on the use of Lp(a) in clinical practice, recommended selective screening of Lp(a) in high-risk children < 18 years of age in certain cases (based on NLA recommendations in the presence of suspected FH, a first-degree relative with premature ASCVD or elevated Lp(a), or a history of ischemic stroke) [72]. Guidelines also recommend cascade screening of immediate family members of a child or adult with elevated Lp(a) [45, 56, 72]. Recent studies proved that cascade testing for elevated Lp(a) was effective in identifying new cases of elevated Lp(a) [73] (Table I).

Concentrations of Lp(a) and CV risk

CVD risk rises progressively with increasing Lp(a) levels [74, 75], indicating a direct relationship between higher Lp(a) concentrations and elevated cardiovascular risk. However, specific thresholds have been defined and implemented in practice to guide clinical management. Therefore, an Lp(a) level requiring clinical intervention is defined as ≥ 30 mg/dl (≥ 75 nmol/l). Patients with Lp(a) levels between 30 and 50 mg/dl (75–125 nmol/l) fall into a gray zone (also called intermediate risk), which is influenced by both the test’s accuracy and the individual’s overall risk profile [45, 56]. Nonetheless, such individuals already face an elevated risk of adverse events and should be categorized as having moderate risk.

High risk is associated with Lp(a) concentrations above 50 mg/dl (> 125 nmol/l), particularly within the range of 50–180 mg/dl (125–450 nmol/l), while concentrations exceeding 180 mg/dl (> 450 nmol/l) indicate very high risk [45, 56]. There is still an ongoing discussion on the importance of the gray zone in CVD risk prediction. Recent data from the STAR-Lp(a) study showed that the mean CAC-Score in those at low risk (Lp(a) < 30 mg/dl) and patients in the gray zone was almost the same (203.1 ± 414.3 vs. 216.6 ± 469.4) and essentially increased only for Lp(a) > 50 mg/dl (125 nmol/l) (335.6 ± 784.9) [76].

Elevated Lp(a) levels of ≥ 50 mg/dl (≥ 125 nmol/l) are estimated to affect over 1.5 billion individuals globally [2]. Despite this high prevalence, testing for Lp(a) is conducted far less frequently than needed; in most countries, only a few to several percent of patients are tested [77, 78]. A multi-center cross-sectional epidemiological study of

48,135 patients with a history of ASCVD revealed that Lp(a) levels were assessed in only a small proportion of cases (14%) [7]. The prevalence of ASCVD is nearly three times greater in adults with Lp(a) levels above the 99th percentile compared to those with Lp(a) levels at or below the 20th percentile. Among individuals with very high Lp(a), including Lp(a) as a factor leads to the reclassification of one-third of patients in primary prevention and more than half in secondary prevention [79]. Findings from Lp(a) registries highlight the importance of routine Lp(a) measurement, as elevated levels are prevalent among individuals at risk for CVD in primary and secondary prevention settings. This underscores the need for risk re-stratification and treatment optimization in such patients [3–5].

CVD risk and Lp(a)

Elevated Lp(a) levels are associated with increased risk for several cardiovascular diseases including ASCVD [27, 75], aortic stenosis/calcific aortic valve disease (CAVD) [80], ischemic stroke [81], peripheral arterial disease (PAD) [82], heart failure [83, 84], and atrial fibrillation [85]. Based on the data from the Copenhagen General Population Study, Lp(a) levels required to achieve a hazard ratio of 1.5 are 154 nmol/l (~ 62 mg/dl) for CAVD, 193 nmol/l (~ 77 mg/dl) for MI, 261 nmol/l (~ 104 mg/dl) for ischemic stroke, and 323 nmol/l (~ 129 mg/dl) for HF [72]. Although average or median Lp(a) levels differ globally and across ancestries, the relative risk associated with baseline Lp(a) concentrations appears to be broadly consistent among the populations studied to date [2]. However, the impact of ASCVD attributable to elevated Lp(a) levels may be over twice as high in individuals of African descent compared to those of Caucasian descent [86]. In an analysis of the Women’s Health Study (WHS), the influence of Lp(a), LDL-C, and hsCRP levels on the 30-year risk of CVD events in women was quantified. Elevated Lp(a) levels at baseline were a significant predictor of cardiovascular events over 30 years of follow-up (HR = 1.33, 95% CI: 1.21–1.47). Each of these three biomarkers contributed independently to risk assessment. Considering Lp(a) measurement in addition to traditional markers, such as LDL-C and hsCRP, improves long-term assessment of cardiovascular risk in women [87]. A meta-analysis of randomized controlled trials (RCTs) involving statin therapy including those with and without clinical ASCVD demonstrated that Lp(a) is an independent risk factor across the spectrum of LDL-C, including those with LDL-C < 77 mg/dl [88].

HsCRP is widely recognized as a marker for systemic inflammation and is frequently used in clinical settings to assess inflammatory status and the risk of inflammation-related ASCVD [89]. Previous

research showed that elevated Lp(a) was associated with a higher level of hsCRP [3]. Recent studies have investigated the association between Lp(a) levels and the risk of major adverse cardiovascular events (MACE) concerning hsCRP levels. Small *et al.* found that higher levels of Lp(a) were associated with MACE, MI, and PAD in both primary and secondary prevention populations regardless of baseline hsCRP [90, 91]. Interestingly, Arnold *et al.* observed that while among individuals without coronary heart disease (CHD) Lp(a) was significantly associated with incident CHD regardless of hsCRP, in participants with CHD at baseline, Lp(a) was related to recurrent CHD events only in those with residual inflammatory risk [92]. Similar results were observed in the analysis from the MESA (Multi-Ethnic Study of Atherosclerosis) study, which identified Lp(a)-associated ASCVD risk only with concomitant elevation of hsCRP [93]. Due to the inconsistency of these studies, this issue merits further investigation [70, 94].

Recently, Alebna *et al.* [95] conducted a meta-analysis of 562,301 individuals from 11 large cohort studies to clarify this issue. They found that elevated Lp(a) was significantly associated with an increased risk of MACE, regardless of hsCRP levels, in both primary and secondary prevention settings [95]. This discovery highlights that Lp(a) contributes to atherogenesis through various mechanisms and holds significant clinical implications for Lp(a)-targeted therapies, indicating their potential effectiveness in all patients with elevated Lp(a), not just those with concurrently high Lp(a) and elevated hsCRP levels [95].

Coronary artery calcification (CAC) is a hallmark of atherosclerosis and is strongly linked to the overall burden of atherosclerotic plaques [96]. The CAC score is a well-validated metric used to assess atherosclerotic disease burden and guide primary ASCVD management decisions. Findings from the MESA (Multi-Ethnic Study for Atherosclerosis) study revealed that Lp(a) levels and CAC score were independently associated with ASCVD risk and may help guide primary prevention strategies. Participants with elevated Lp(a) and a CAC score > 100 experienced the highest risk (HR = 4.71; 95% CI: 3.01–7.40) compared to those with non-elevated Lp(a) and a CAC score of 0, while individuals with elevated Lp(a) and a CAC score of 0 exhibited a modestly increased risk (HR = 1.31; 95% CI: 0.73–2.35) [97]. The above-mentioned Burzyńska *et al.* [76] study showed that for each 10 mg/dl (25 nmol/l) increase in Lp(a), the CAC score rose by 15.7 ± 0.57 ($p = 0.006$). In patients with advanced stable coronary artery disease monitored via coronary computed tomography angiography (CCTA) at baseline and after 12 months, elevated Lp(a) levels (> 70 mg/dl) were associated with acceler-

ated progression of the necrotic core, including changes in total, calcific, noncalcific, and low-attenuation plaque [98].

A meta-analysis including 40,073 individuals from 17 studies found that elevated Lp(a) levels were significantly associated with a higher prevalence of CAC (OR = 1.31, 95% CI 1.06–1.61, $p = 0.01$). When analyzed as a continuous variable, higher Lp(a) levels positively correlated with CAC prevalence (OR = 1.05, 95% CI: 1.02–1.08, $p = 0.003$). Additionally, elevated Lp(a) was associated with increased progression of CAC over time (OR = 1.54, 95% CI: 1.23–1.92, $p = 0.0002$) [99].

The relationship between inflammation, Lp(a), and the risk of aortic valve calcification (AVC) was analyzed in a subgroup of patients from the MESA study. The study included 6,676 participants and assessed baseline levels of Lp(a), hsCRP, and AVC using prior non-contrast cardiac computed tomography. Elevated Lp(a) levels were independently associated with AVC, and individuals with both high Lp(a) and elevated hsCRP (> 2 mg/dl) had the highest risk for developing AVC [100]. Lp(a) and its associated molecules – OxPL, autotaxin (ATX), and lysophosphatidic acid (LysoPA) – play critical roles in the development of ASCVD and AVS [101].

Measuring Lp(a) levels in patients with calcific aortic valve disease can help to predict substantially faster disease progression and the likelihood of requiring aortic valve replacement (AVR) [102, 103]. A recent cohort study involving 44,742 patients from a Korean center with Lp(a) level measured from 2000 to 2020, with a mean follow-up of 6.8 years, indicated that AVR due to severe degenerative aortic stenosis was significantly associated with higher levels of Lp(a) (> 100 mg/dl) (adjusted HR = 2.05; 95% CI: 1.31–3.19; $p = 0.002$) [104]. The authors present compelling evidence reinforcing the association between Lp(a) and advanced CAS, and the necessity for AVR [105].

Aortic valve stenosis, characterized by valvular calcification and stiffness, can lead to heart failure (HF) [106]. Lp(a) has been identified as a possible risk factor for developing heart failure [83, 84]. A meta-analysis of seven Mendelian randomization studies with 300,255 individuals was conducted to explore the causal relationship between Lp(a) and its role in HF. It was demonstrated that increasing Lp(a) levels were significantly associated with increased risk of HF (OR = 1.064, 95% CI: 1.043–1.086, $I^2 = 97.59\%$, $p < 0.001$) [107]. Moreover, Lp(a) could have a greater impact on HF patients compared to other lipid parameters [89]. The CASABLANCA (Catheter Sampled Blood Archive in Cardiovascular Diseases) study with a total of 1251 individuals indicated that Lp(a) and associated OxPLs may independently contribute to heart injury, leading to HF. Further research,

including clinical trials, focused on reducing Lp(a) levels, is needed to determine whether such interventions can prevent the progression to symptomatic HF or mitigate its complications [108].

One of the controversial issues associated with the estimation of CV risk is that cholesterol carried by Lp(a) (Lp(a)-C) is included in both calculated and directly measured LDL-C, as well as in calculated non-HDL-C [109, 110]. Earlier studies suggested that Lp(a) particles contain about 30% cholesterol by mass [109]. However, a recent study involving 68,748 ASCVD-free individuals followed for a median of 9.7 years to track CHD events found that adjusting LDL-C for its Lp(a)-C content did not significantly enhance CHD risk estimation at the population level [111]. The updated NLA guidelines did not endorse the previously proposed correction factor for Lp(a)-C in LDL-C calculations, citing concerns about potentially undertreating high-risk patients [72]. Despite this recommendation, the debate continues. Tsimikas *et al.* [112] pointed out that the mean Lp(a) level in the Arnold *et al.* [111] study was only 9.3 mg/dl, and the top decile reached just 43.5 mg/dl – substantially lower than levels associated with elevated ASCVD risk. They emphasized the need for empirically measuring Lp(a)-C and incorporating this adjustment in LDL-C to better understand its impact in observational studies and clinical trials across diverse populations [112] (Table I).

Available lipid-lowering therapies and Lp(a)

Currently, no medications specifically designed to reduce Lp(a) levels have been approved. Consequently, managing elevated Lp(a) involves focusing on reducing overall cardiovascular risk through lifestyle modifications and the intensive management or optimization of other treatable risk factors, by following current clinical guidelines [45, 56, 113, 114].

The findings from studies investigating the impact of statins on Lp(a) levels are inconsistent [25], and the effect seems to be related to apo(a) isoforms. In patients with the low molecular weight apo(a) phenotype, Lp(a) levels increased significantly from 66.4 to 97.4 mg/dl (by 47%; $p = 0.026$), but not in patients characterized by the high molecular weight apo(a) phenotype [115]. However, even a minor Lp(a) increase of approximately 6–10% following statin use is not clinically significant [116, 117]. Pitavastatin, in contrast to other statins, seems to have a neutral impact on serum Lp(a) levels and may even slightly reduce them; however, this still needs to be confirmed [118]. Ezetimibe and bempedoic acid do not affect Lp(a) levels, as a comprehensive analysis recently found [119–121]. Niacin can lower lipoprotein(a) levels by about 20–30% (depending on the base-

line Lp(a) level) due to a decreased *LPA* mRNA and apo(a) production rate [122]. Again, the amount of reduction seems to be dependent on the size of the apo(a) isoforms [56]. However, no clinical benefit was noted with this therapy, so its use is no longer currently recommended [123, 124]. The prior CVD outcome trials involving niacin were also not enriched for individuals with elevated Lp(a).

Proprotein convertase subtilisin/kexin type 9 inhibitors (PCSK9is) and small interfering RNAs (inclisiran) were found to decrease circulating Lp(a) by ~30%, as demonstrated in various studies including the FOURIER, ODYSSEY Outcomes, and ORION-11 trials [125–129]. *Post-hoc* analyses of the FOURIER and ODYSSEY Outcomes trials demonstrated that the 3-year number needed to treat (NNT) to prevent one recurrent ASCVD event with PCSK9 monoclonal antibodies is at least 2.5-fold lower for individuals with higher versus lower levels of Lp(a) [125, 126]. These drugs, however, are not approved for this purpose, making it difficult to obtain reimbursement for their use in lowering Lp(a). Significant variability in response to these drugs is also observed, with participants who had higher baseline plasma Lp(a) levels experiencing greater absolute reductions in Lp(a) [127–130]. The size of apo(a) seems to be an independent determinant of the response to PCSK9is, with each additional kringle domain being associated with a 3% additional reduction in Lp(a) (the larger the isoforms, the lower the Lp(a) level, the better the response) [130].

A significant number of patients on PCSK9 inhibitors are concurrently treated with statins and/or ezetimibe. However, the combined impact of these therapies on Lp(a) levels, as well as the relationship between apo(a) isoform size and the Lp(a) response, remains unclear. A recent prospective study evaluated lipid lowering in participants with an LDL-C > 100 mg/dl who received evolocumab 140 mg combined with either atorvastatin 80 mg or ezetimibe 10 mg daily. The findings revealed variability in Lp(a) reduction, with changes in Lp(a) levels being strongly associated with apo(a) isoform size [131].

Lipoprotein apheresis is currently the only method capable of significantly lowering Lp(a) levels [132, 133]. A single apheresis session can reduce Lp(a) concentrations by approximately 60–75%, while regular treatments every 1–2 weeks result in a sustained reduction of around 25–40% from baseline levels [134, 135]. Lipoprotein apheresis in patients with ASCVD and elevated Lp(a) leads to a notable decrease in cardiovascular events [136]. The US Food and Drug Administration (FDA), as well as European guidelines [56], approved apheresis for patients with Lp(a) levels exceeding 60 mg/dl (> 150 nmol/l), regardless of

baseline LDL-C levels [137]. However, this treatment approach is invasive, expensive, time-consuming for the patient, has limited availability, and is often associated with a diminished quality of life [138] (Table I).

Aspirin and Lp(a)

Aspirin is one of the most well-established therapies for secondary prevention of ASCVD related events, and based on recent data it may also be beneficial in primary prevention for patients with elevated plasma Lp(a). Large RCTs of aspirin use for primary prevention, including ARRIVE (Aspirin to Reduce Risk of Initial Vascular Events), ASCEND (A Study of Cardiovascular Events in Diabetes), and ASPREE (Aspirin in Reducing Events in the Elderly), conducted in different populations, found no significant net beneficial effect of aspirin on CVD events [139–141]. A meta-analysis of 13 RCTs with 164,225 participants and 1,050,511 participant-years of follow-up indicated a small benefit of aspirin use for CVD risk in primary prevention [142].

Post-hoc analyses of the ASPREE trial have shown that individuals with elevated Lp(a) genotypes may derive a net benefit with aspirin therapy [143]. In a recent propensity-matched cohort study (MESA), aspirin use was associated with a significant reduction (46%) in risk for cardiovascular events among individuals with Lp(a) > 50 mg/dl and without baseline cardiovascular disease [144]. The patients with Lp(a) > 50 mg/dl and aspirin use had similar CHD risk as those with Lp(a) ≤ 50 mg/dl, regardless of aspirin use [144]. This was the first study to focus on Lp(a) level measurements, contrary to earlier studies that used the SNP of *LPA*. These results align with earlier evidence showing a 45–55% reduction in the risk of initial MACE among regular aspirin users who carry the rs3789220 *LPA* gene variant, while no such benefit was observed in non-carriers [143, 145].

Moreover, the results from the MESA study were further supported by data from the National Health and Nutrition Examination Survey (NHANES III) analysis, which gathered baseline data from 1988 to 1994 in a nationally representative cohort of US adults without clinical ASCVD. Their findings revealed that regular aspirin use was independently associated with a 52% reduction in ASCVD mortality risk among individuals with elevated Lp(a), but this association was not observed in those without elevated Lp(a) during a median follow-up period of 26 years [146]. Interestingly, in the *post-hoc* analysis of the PEGASUS-TIMI 54 trial, individuals with elevated Lp(a) ≥ 200 nmol/l (80 mg/dl) and a history of MI (within 1 to 3 years) may possibly more strongly benefit from dual antiplatelet therapy (DAPT) with ticagrelor in the setting of background aspirin [147].

While bleeding events were not evaluated in that study, the MESA study reported a higher bleeding rate among aspirin users (17.5% vs. 12.5%, $p < 0.01$). However, after multivariable adjustments, no correlation was observed between bleeding rates and Lp(a) levels. Consequently, additional research is necessary to assess the overall (net) clinical benefit in this context [148].

There is a lack of effective therapies specifically aimed at reducing cardiovascular disease risk in individuals with elevated lipoprotein(a), particularly for primary prevention. In the just-published recommendations of Polish Experts endorsed by the Polish Lipid Association (PoLA), aspirin is not recommended for primary prevention in patients with low or moderate cardiovascular risk, regardless of the co-occurrence of elevated Lp(a) concentration. However, the use of aspirin in primary prevention should be considered (class IIa recommendation) in patients with at least high cardiovascular risk with elevated Lp(a) levels > 50 mg/dl (> 125 nmol/l), which may help optimize the risk of ASCVD associated with it [149] (Table I).

New therapies for lowering Lp(a)

Although there is a pressing demand for therapies specifically designed to lower Lp(a) levels, the sole available targeted treatment option at present is apheresis. No FDA-approved medications exist for this purpose yet, but several innovative treatments to lower Lp(a) are under investigation in late-stage randomized controlled trials [150].

Several RNA-based therapies against Lp(a) are under development. The success of the first agent – pelacarsen, an antisense oligonucleotide (ASO) – paved the way for the development of further small interfering RNAs (siRNAs) agents, such as olpasiran, zerlasiran, and lepodisiran [41, 151–153] (Table II). The mechanism of action of ASOs and siRNAs is to reduce Lp(a) levels by inhibiting apo(a) protein synthesis [154]. Maximum reported Lp(a) reduction for this drug was 80% for pelacarsen and 98% for olpasiran, zerlasiran, and lepodisiran [150].

Pelacarsen, given as an 80 mg subcutaneous injection once a month, is currently undergoing a phase 3 clinical trial to evaluate its impact on reducing CVD event risk (Lp(a) HORIZON trial, NCT04023552). The trial has completed enrollment of over 8,000 patients but is expected to conclude in 2026 because of the low event rate [155]. Moreover, pelacarsen is also under investigation in the CAVS trial, a phase 2 randomized controlled study (NCT05646381) designed to evaluate its ability to slow the progression of aortic stenosis (AS) compared to placebo. The trial targets patients with mild to moderate AS and elevated Lp(a) levels, plans to recruit around

Table II. Lp(a)-lowering drugs under clinical development

Drug name	Company	Class	Max Lp(a) reduction	Main inclusion criteria	Administration route, frequency, and dose	Trial phase	NCT	Planned study completion
Pelacarsen	Novartis Pharmaceuticals	ASO	80%	<ul style="list-style-type: none"> Age 18 to 80 years Lp(a) ≥ 70 mg/dl Myocardial infarction Ischemic stroke Clinically significant symptomatic PAD 	Subcutaneous injection, monthly, 80 mg	III	NCT04023552	30 May 2025
Olpasiran	Amgen	siRNA	98%	<ul style="list-style-type: none"> Age 18 to ≤ 85 years Lp(a) ≥ 200 nmol/l History of ASCVD: <ul style="list-style-type: none"> - Myocardial infarction (presumed type 1 event due to plaque rupture/erosion) and/or - Coronary revascularization with percutaneous coronary intervention AND at least 1 additional risk factor 	Subcutaneous injection 75 or 225 mg once every 12 weeks	III	NCT05581303	29 Dec 2026
Lepodisiran	Eli Lilly	siRNA	98%	<ul style="list-style-type: none"> Lp(a) ≥ 175 nmol/l Meet criteria 2a or 2b 2a: age ≥ 18 years and established ASCVD with an event or revascularization 2b: age ≥ 55 years and risk for a first CV event and either: documented CAD or PAD; known FH; or a combination of high-risk factors 	Subcutaneous injection 400 mg	III	NCT06292013	Mar 2029
Zerlasiran	Silence Therapeutics	siRNA	98%	<ul style="list-style-type: none"> Age 18 to 80 years Lp(a) ≥ 125 nmol/l At high risk of ASCVD events BMI 18.0 to 32.0 kg/m² 	Subcutaneous injection, 450 mg every 24 weeks for 2 doses, 300 mg every 16 weeks for 3 doses, or 300 mg every 24 weeks for 2 doses	II	NCT05537571	01 Jul 2024
Muvalaplin	Eli Lilly	Lp(a) inhibitor	85%	<ul style="list-style-type: none"> Age ≥ 40 years Lp(a) ≥ 175 nmol/l High risk for cardiovascular events: documented CAD, stroke, or PAD or ASCVD risk equivalents (FH or T2DM) BMI 18.5 to 40 kg/m² 	Oral, 10 mg/day, 60 mg/day, or 240 mg/day for 12 weeks	II	NCT05563246	14 Mar 2024
CTX320	CRISPR Therapeutics	gene editing	95%	–	One-time treatment 2 mg/kg infusion	Preclinical	–	–
Obicetrapib	NewAmsterdam Pharma	CETP inhibitor	57%	<ul style="list-style-type: none"> Age 18–75 years Lp(a) > 50 mg/dl LDL-C > 70 mg/dl TG < 400 mg/dl 	Oral obicetrapib 10 mg daily for 8 weeks, followed by obicetrapib 10 mg daily plus Repatha 140 mg SC biweekly for 8 weeks	II	NCT06496243	Dec 2025

ASCVD – atherosclerotic cardiovascular disease, ASO – antisense oligonucleotides, CAD – coronary artery disease; CETP – cholesteryl ester transport proteins, CV – cardiovascular, FH – familial hypercholesterolemia, PAD – peripheral artery disease, SC – subcutaneous, siRNA – small interfering RNAs, T2DM – type 2 diabetes mellitus.

500 participants, and is projected to conclude in 2029 [156].

Recently reported results of the phase 2 OCEAN(a)-DOSE off-treatment extension period indicated that olpasiran demonstrates long-lasting effects in reducing Lp(a) levels, with participants who received doses of ≥ 75 mg every 12 weeks maintaining approximately a 40% to 50% decrease in Lp(a) levels nearly 1 year after their final dose [151]. Olpasiran, developed by Amgen, is currently being evaluated in the phase 3 clinical trial known as the Ocean(a) study (NCT05581303), which has enrolled approximately 7,000 participants. The findings, anticipated in 2027, are expected to follow the results of the HORIZON study by about a year and will shed light on olpasiran's impact on cardiovascular events [157]. Also worth noting is a forthcoming RCT with olpasiran in primary prevention: A Double-blind, Randomized, Placebo-controlled, Multi-center Study Assessing Olpasiran Use to Prevent First Major Cardiovascular Events in High-risk Participants with Elevated Lipoprotein(a). It is of critical importance, as most patients with elevated Lp(a) are currently diagnosed in primary CVD prevention, still before the event.

Lepodisiran, developed by Eli Lilly, has completed its phase 1 trial. Its phase 2 trial (the Alpaca Phase 2 trial; NCT05565742) is currently underway and is expected to end in October 2024 [152, 158]. The results of this lepodisiran trial with extended duration up to 540 days are to be presented at the ACC Congress in Chicago in March 2025. Undergoing a phase 3 clinical trial, ACCLAIM Lp(a) (NCT06292013), is projected to be the largest study of Lp(a)-lowering therapies, aiming to enroll 12,500 participants, with an expected completion date in 2029 [159]. The ACCLAIM trial is another study that will allow evaluation of the effect of Lp(a) reduction on MACE both in adults with elevated Lp(a) who have established ASCVD and in those in primary prevention who are at risk for a first CVD event [159].

The last siRNA drug, zerlasiran, has completed its phase 1 clinical trial and a phase 2 trial (NCT05537571). The results were released at the American Heart Association Scientific Sessions in Chicago 2024 and published simultaneously. Zerlasiran achieved over an 80% reduction in time-averaged lipoprotein(a) levels over 36 weeks when administered at 300 mg every 16 weeks or 300 mg and 450 mg every 24 weeks. Sustained reductions in Lp(a) levels were observed up to 60 weeks after the initial dose, and no safety concerns were identified. These results support advancing zerlasiran to a phase 3 trial in its development program [152]. Lepodisiran and zerlasiran have similar effects as olpasiran, with an over 90% Lp(a)

reduction at the highest doses, with comparable tolerability and safety profiles [152, 153].

In 2024, there was a major breakthrough in developing a new strategy for lowering Lp(a) levels. Results of phase II clinical research with the first oral agent, muvalaplin, were published, and the new therapy based on gene editing was revealed in preclinical studies [160, 161]. Muvalaplin, developed by Elly Lilly, is a small molecule inhibitor of Lp(a) synthesis, lowering Lp(a) levels by binding to apo(a) KIV7 and KIV8. This interaction sterically hinders the covalent attachment of apo(a) to apoB, thereby reducing Lp(a) formation [162]. Findings from a phase I study (NCT04472676) demonstrated that the treatment was well tolerated and achieved a maximum reduction in Lp(a) levels ranging from 63% to 65% [160]. The phase II KRAKEN trial with muvalaplin (NCT05563246) enrolled 233 participants (median age 66 years) with lipoprotein(a) ≥ 175 nmol/l and ASCVD, diabetes, or familial hypercholesterolemia. Treatment was for 12 weeks. Muvalaplin was well tolerated and caused placebo-adjusted reductions in lipoprotein(a) of 47.6% (95% CI: 35.1–57.7%), 81.7% (95% CI: 78.1–84.6%), and 85.8% (95% CI: 83.1–88.0%) for the dose of 10 mg/day, 60 mg/day, and 240 mg/day, respectively [163].

Available commercial Lp(a) assays measure total apo(a), apo(a) in the Lp(a) particle, and apo(a) that is not bound to apoB, and may be insufficient to accurately measure Lp(a) concentrations, especially after muvalaplin treatment, taking into consideration its mechanism of action where apo(a)-muvalaplin complexes in circulation might be detected. Swearingen *et al.* [43] introduced an innovative immunoassay designed to measure only Lp(a) particles. This particle-specific electrochemiluminescent (ECL) immunoassay employs an anti-apo(a) capture antibody that targets a common epitope found in KIV7, KIV8, and KIV9, ensuring isoform insensitivity. Detection is achieved using an anti-apoB monoclonal antibody, preventing the assay from identifying unbound apo(a). A recent study evaluated the Lp(a)-lowering effects of two therapeutics with distinct mechanisms of action: lepodisiran and muvalaplin. The results revealed that the commercial assay measuring total apo(a) underestimated the Lp(a)-lowering efficacy of muvalaplin compared to the intact Lp(a) assay, which specifically measures Lp(a) particles. However, the Lp(a)-lowering impact of lepodisiran was found to be clinically comparable between the intact Lp(a) assay and the commercial assay [43].

CTX320 is an experimental CRISPR/Cas9-based gene editing therapy designed to target and disable the apo(a) component of Lp(a) production in the liver. The therapy utilizes lipid nanoparticles to deliver Cas9 mRNA and guide RNA (gRNA) di-

rectly into the body. CTX320 has the potential to permanently reduce Lp(a) after a one-time treatment [161]. Preclinical studies in non-human primates demonstrated that CTX320 reduced Lp(a) levels in a dose-dependent manner, achieving approximately 20%, 80%, and 90% reductions from baseline at doses of 0.5, 1.5, and 3 mg/kg, respectively [155]. In a separate ongoing study, a single infusion of CTX320 at 2 mg/kg resulted in a ~94% reduction in Lp(a) levels by day 14, with this reduction maintained until day 224. The therapy was well tolerated in these studies and showed a durable lowering of plasma Lp(a). As a result, CTX320 is being prepared for advancement to phase I clinical trials [164].

Obicetrapib, a next-generation selective inhibitor of cholesteryl ester transporter protein (CETP),

is currently undergoing clinical trials to lower LDL-C levels and reduce major adverse cardiovascular events. The results of the phase 2 ROSE trial (NCT04753606) demonstrated that obicetrapib, administered orally at doses of 5 mg and 10 mg alongside intensive statin therapy, reduced Lp(a) levels by 33.8% and 56.5%, respectively, compared to placebo [165]. The phase 2 ROSE1 and ROSE2 trials investigated obicetrapib as an add-on therapy to high-intensity statins in individuals without CVD but with LDL cholesterol levels > 70 mg/dl. A pooled analysis revealed that obicetrapib 10 mg, combined with high-intensity statin therapy, significantly reduced Lp(a) levels by 57% compared to placebo. This reduction surpasses those achieved with PCSK9 inhibitors (15–30%), niacin (30%), or other CETP inhibitors (25%) [166]. At the AHA 2024,

New findings about Lipoprotein(a) in 2024




Figure 1. New findings on lipoprotein(a) based on the data from 2024 studies. Based on information from [2, 6, 9, 43, 80, 137, 153, 168, 170]

DM – diabetes mellitus, Lp(a) – lipoprotein(a), LDL – low-density lipoprotein, MACE – major adverse cardiovascular event.

the results of the BROOKLYN trial with obicetrapib in patients with heFH (baseline Lp(a) level in the obicetrapib group was 45.8 nmol/l) were released, showing a 54.3% placebo-adjusted Lp(a) reduction in the intervention arm and 38% of patients with > 50% Lp(a) reduction [167]. Currently, obicetrapib is under investigation in a phase 2 trial (VINCENT). It is an open-label, 16-week trial aimed at evaluating Lp(a) levels for patients with elevated Lp(a) being treated with obicetrapib and obicetrapib/evolocumab. The VINCENT trial started recruiting patients at the end of 2024 and will be completed at the end of 2025 (NCT06496243) [168] (Table II).

Safety of very low levels of Lp(a)

Observational and epidemiological studies have suggested that extremely low Lp(a) levels, typically below 7 mg/dl, are associated with an elevated risk of type 2 diabetes mellitus (T2DM), as has been shown for LDL lowering with statin treatment [169–171]. However, the underlying mechanism and causality of this relationship have not been established [172, 173], leaving open the question of whether novel Lp(a)-lowering therapies might contribute to an increased risk of developing T2DM [174].

The results of a Mendelian randomization (MR) analysis published in 2024 shed more light on this association. Data from a two-sample Mendelian randomization analysis involving data from 563,420 patients from the UK Biobank and FinnGen consortia did not show a correlation between Lp(a) and T2DM [175]. Another study with a two-sample MR analysis of the UK Biobank population cohort also found no evidence for an association between genetically predicted Lp(a) and T2D [176]. Moreover, two-sample MR analysis using summary-level genome wide association data suggested that hyperinsulinemia, often associated with type 2 diabetes mellitus, may partially explain the inverse association observed between low lipoprotein(a) levels and an increased risk of developing type 2 diabetes mellitus [177]. High fasting insulin levels, which contribute to the progression of prediabetes and type 2 diabetes, are actually responsible for the observed reduction in Lp(a) level. Therefore, while a connection between Lp(a) and diabetes does exist, it is unlikely that Lp(a) serves as a risk factor for diabetes (no reverse causality) independent of the presence of underlying hyperinsulinemia and insulin resistance [177]. Thus, aggressive Lp(a)-lowering therapy at this time does not raise any substantial concerns about exacerbated T2DM risk (Table I).

Conclusions and take-home message

In 2024, knowledge about the importance of Lp(a) as a risk factor for cardiovascular diseases

was deepened, and great progress was made in developing therapies to effectively lower its levels, which may contribute to reducing the risk of these diseases in the future [178, 179]. While most clinical trials with new Lp(a)-lowering therapies focus on ASCVD patients or those at CVD risk, only one started in 2024 will evaluate the ability of pelacarsen to slow the progression of aortic stenosis. Moreover, there was a breakthrough in therapy for lowering Lp(a) levels based on gene editing. Currently, Lp(a) is utilized in clinical practice to refine risk stratification and guide patients toward more aggressive risk factor management. In the future, it may be possible to implement targeted treatments specifically aimed at lowering Lp(a) levels. However, the physiological functions of Lp(a) are still largely unknown. Although some safety concerns over lowering Lp(a) to very low levels remain, recent Mendelian randomization studies did not find an association between very low levels of Lp(a) and new onset T2D. Lipoprotein(a) research is a rapidly evolving field, but many questions remain unanswered. We eagerly await the results of the HORIZON, OCEAN and ACCLAIM studies to establish whether or not Lp(a) lowering will transform our approach to the prevention of ASCVD (Figure 1).

Funding

This review did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Ethical approval

Not applicable.

Conflict of interest

Individual declarations of conflict of interest: Maciej Banach has received grant(s)/research support from Amgen, Daiichi-Sankyo, Mylan/Viatriis, and Sanofi, and has served as a consultant and/or received speaker's fees from Adamed, Amgen, Exceed Pharma, Daiichi-Sankyo, Esperion, Kogen, KRKA, Lilly, MSD, Mylan/Viatriis, NewAmsterdam Pharma, Novartis, Novo-Nordisk, Polpharma, Sanofi, Teva and Zentiva; all other authors do not declare any conflict of interest.

References

1. Cybulska B, Kłosiewicz-Latoszek L, Penson PE, Banach M. What do we know about the role of lipoprotein(a) in atherogenesis 57 years after its discovery? *Prog Cardiovasc Dis* 2020; 63: 219-27.
2. Tsimikas S, Marcovina SM. Ancestry, lipoprotein(a), and cardiovascular risk thresholds. *J Am Coll Cardiol* 2022; 80: 934-46.
3. Sosnowska B, Lewek J, Adach W, et al. The prevalence, patients' characteristics, and hyper-Lp(a)-emia risk

- factors in the Polish population. The first results from the PMMHRI-Lp(a) Registry. *Prog Cardiovasc Dis* 2024; 86: 54-61.
- Dyrbuś K, Mędrala Z, Konsek K, et al. Lipoprotein(a) and its impact on cardiovascular disease – the Polish perspective: design and first results of the Zabrze-Lipoprotein(a) Registry. *Arch Med Sci* 2024; 20: 1069-76.
 - Burzyńska M, Jankowski P, Babicki M, Banach M, Chudzik M. Prevalence of hyperlipoproteinemia(a) in individuals of European ancestry treated at outpatient cardiology clinics: results from a cross-sectional STAR-Lp(a) study. *Pol Arch Intern Med* 2024; 134: 16860.
 - Koren MJ, Moriarty PM, Baum SJ, et al. Preclinical development and phase 1 trial of a novel siRNA targeting lipoprotein(a). *Nat Med* 2022; 28: 96-103.
 - Nissen SE, Wolski K, Cho L, et al. Lipoprotein(a) levels in a global population with established atherosclerotic cardiovascular disease. *Open Heart* 2022; 9: e002060.
 - Björnson E, Adiels M, Taskinen MR, et al. Lipoprotein(a) is markedly more atherogenic than LDL. *J Am Coll Cardiol* 2024; 83: 385-95.
 - Marston NA, Melloni GEM, Murphy SA, et al. Per-particle cardiovascular risk of lipoprotein(a) vs non-Lp(a) apolipoprotein B-containing lipoproteins. *J Am Coll Cardiol* 2024; 83: 470-2.
 - Tsimikas S, Bittner V. Particle number and characteristics of lipoprotein(a), LDL, and apoB. *J Am Coll Cardiol* 2024; 83: 396-400.
 - Strandkjær N, Hansen MK, Nielsen ST, et al. Lipoprotein(a) levels at birth and in early childhood: the COMPARE study. *J Clin Endocrinol Metab* 2022; 107: 324-35.
 - Clarke SL, Huang RDL, Hilliard AT, et al. Genetically predicted lipoprotein(a) associates with coronary artery plaque severity independent of low-density lipoprotein cholesterol. *Eur J Prev Cardiol* 2025; 32: 116-27.
 - Rocha VZ, Santos RD. The mysterious lipoprotein(a): moving towards further understanding of its atherogenic role. *Eur J Prev Cardiol* 2025; 32: 128-30.
 - Boerwinkle E, Leffert CC, Lin J, Lackner C, Chiesa G, Hobbs HH. Apolipoprotein(a) gene accounts for greater than 90% of the variation in plasma lipoprotein(a) concentrations. *J Clin Invest* 1992; 90: 52-60.
 - Burgess S, Ference BA, Staley JR, et al. Association of LPA variants with risk of coronary disease and the implications for lipoprotein(a)-lowering therapies: a Mendelian randomization analysis. *JAMA Cardiol* 2018; 3: 619-27.
 - Lamina C, Kronenberg F, for the Lp(a)-GWAS Consortium. Estimation of the required lipoprotein(a)-lowering therapeutic effect size for reduction in coronary heart disease outcomes: a Mendelian randomization analysis. *JAMA Cardiol* 2019; 4: 575-9.
 - Schmidt K, Noureen A, Kronenberg F, Utermann G. Structure, function, and genetics of lipoprotein (a). *J Lipid Res* 2016; 57: 1339-59.
 - Becker L, Cook PM, Wright TG, Koschinsky ML. Quantitative evaluation of the contribution of weak lysine-binding sites present within apolipoprotein(a) kringle IV types 6–8 to lipoprotein(a) assembly. *J Biol Chem* 2004; 279: 2679-88.
 - Koschinsky ML, Beisiegel U, Henne-Bruns D, Eaton DL, Lawn RM. Apolipoprotein(a) size heterogeneity is related to variable number of repeat sequences in its mRNA. *Biochemistry* 1990; 29: 640-4.
 - Kostner KM, Marz W, Kostner GM. When should we measure lipoprotein (a)? *Eur Heart J* 2013; 34: 3268-76.
 - Kronenberg F, Utermann G. Lipoprotein(a): resurrected by genetics. *J Intern Med* 2013; 273: 6-30.
 - Lobentanz E, Dieplinger H. Biogenesis of lipoprotein(a) in human and animal hepatocytes. *Electrophoresis* 1997; 18: 2677-81.
 - Coassin S, Kronenberg F. Lipoprotein(a) beyond the kringle IV repeat polymorphism: the complexity of genetic variation in the LPA gene. *Atherosclerosis* 2022; 349: 17-35.
 - Clarke R, Peden JF, Hopewell JC, et al. Genetic variants associated with Lp(a) lipoprotein level and coronary disease. *N Engl J Med* 2009; 361: 2518-28.
 - Koschinsky ML, Kronenberg F. The long journey of lipoprotein(a) from cardiovascular curiosity to therapeutic target. *Atherosclerosis* 2022; 349: 1-6.
 - Banach M. Lipoprotein(a): the enemy that we still don't know how to defeat. *Eur Heart J Open* 2023; 3: oead080.
 - Boffa MB, Koschinsky ML. Lipoprotein (a): truly a direct prothrombotic factor in cardiovascular disease? *J Lipid Res* 2016; 57: 745-57.
 - Boffa MB. Beyond fibrinolysis: the confounding role of Lp(a) in thrombosis. *Atherosclerosis* 2022; 349: 72-81.
 - Lawn RM, Schwartz K, Patthy L. Convergent evolution of apolipoprotein(a) in primates and hedgehog. *Proc Natl Acad Sci* 1997; 94: 11992-7.
 - Riches K, Porter KE. Lipoprotein(a): cellular effects and molecular mechanisms. *Cholesterol* 2012; 2012: 923289.
 - Yano Y, Shimokawa K, Okada Y, Noma A. Immunolocalization of lipoprotein(a) in wounded tissues. *J Histochem Cytochem* 1997; 45: 559-68.
 - Von Zychlinski A, Kleffmann T, Williams MJA, McCormick SP. Proteomics of lipoprotein(a) identifies a protein complement associated with response to wounding. *J Proteomics* 2011; 74: 2881-91.
 - Tsimikas S. A test in context: lipoprotein(a). *J Am Coll Cardiol* 2017; 69: 692-711.
 - Tsimikas S, Witztum JL. Oxidized phospholipids in cardiovascular disease. *Nat Rev Cardiol* 2024; 21: 170-91.
 - Martínez C, Rivera J, Loyau S, et al. Binding of recombinant apolipoprotein(a) to human platelets and effect on platelet aggregation. *Thromb Haemost* 2001; 85: 686-93.
 - Kamstrup PR, Neely RDG, Nissen S, et al. Lipoprotein(a) and cardiovascular disease: sifting the evidence to guide future research. *Eur J Prev Cardiol* 2024; 31: 903-14.
 - Zimodro JM, Gąsecka A, Arski P, Schwarz J, Banach M, Gouni-Berthold I. Lipoprotein (a) measurement: potential for personalized cardiovascular disease management in a patient with acute myocardial infarction. *Arch Med Sci* 2024; 20: 1043-7.
 - Page MM, Watts GF. Contemporary perspectives on the genetics and clinical use of lipoprotein(a) in preventive cardiology. *Curr Opin Cardiol* 2021; 36: 272-80.
 - Reyes-Soffer G, Ginsberg HN, Berglund L, et al. Lipoprotein(a): a genetically determined, causal, and prevalent risk factor for atherosclerotic cardiovascular disease: a scientific statement from the American Heart Association. *Arterioscler Thromb Vasc Biol* 2022; 42: e48-60.
 - Heydari M, Rezayi M, Ruscica M, Jamialahmadi T, Johnston TP, Sahebkar A. The ins and outs of lipoprotein(a) assay methods. *Arch Med Sci Atheroscler Dis* 2024; 8: 128-39.
 - O'Donoghue ML, Rosenson RS, Gencer B, et al. Small interfering RNA to reduce lipoprotein(a) in cardiovascular disease. *N Engl J Med* 2022; 387: 1855-64.

42. Clouet-Foraison N, Vaisar T, Marcovina SM. Standardization of analytical methods for the measurement of lipoprotein(a): bridging past and future initiatives. In: *Lipoprotein(a)* [Internet]. Kostner K, Kostner GM, Toth PP (eds.). Cham: Springer International Publishing 2023; 297-323. (Contemporary Cardiology).
43. Swearingen CA, Sloan JH, Rhodes GM, et al. Measuring Lp(a) particles with a novel isoform-insensitive immunoassay illustrates efficacy of muvalaplin. *J Lipid Res* 2024; 66: 100723.
44. Kronenberg F. Lipoprotein(a) measurement issues: are we making a mountain out of a molehill? *Atherosclerosis* 2022; 349: 123-35.
45. Kronenberg F, Mora S, Stroes ESG, et al. Lipoprotein(a) in atherosclerotic cardiovascular disease and aortic stenosis: a European Atherosclerosis Society consensus statement. *Eur Heart J* 2022; 43: 3925-46.
46. Ruhaak LR, Cobbaert CM. Quantifying apolipoprotein(a) in the era of proteoforms and precision medicine. *Clin Chim Acta* 2020; 511: 260-8.
47. Szarek M, Reijnders E, Jukema JW, et al. Relating lipoprotein(a) concentrations to cardiovascular event risk after acute coronary syndrome: a comparison of 3 tests. *Circulation* 2024; 149: 192-203.
48. Johansen AK, Bogsrud MP, Thoresen M, et al. Lipoprotein(a) in children and adolescents with genetically confirmed familial hypercholesterolemia followed up at a specialized lipid clinic. *Atheroscler Plus* 2024; 57: 13-8.
49. Simony SB, Mortensen MB, Langsted A, Afzal S, Kamstrup PR, Nordestgaard BG. Sex differences of lipoprotein(a) levels and associated risk of morbidity and mortality by age: The Copenhagen General Population Study. *Atherosclerosis* 2022; 355: 76-82.
50. Corral P, Matta MG, Aguilar-Salinas C, et al. Lipoprotein(a) throughout life in women. *Am J Prev Cardiol* 2024; 20: 100885.
51. Sattar N, Clark P, Greer IA, Shepherd J, Packard CJ. Lipoprotein (a) levels in normal pregnancy and in pregnancy complicated with pre-eclampsia. *Atherosclerosis* 2000; 148: 407-11.
52. Wang Q, Würtz P, Auro K, et al. Metabolic profiling of pregnancy: cross-sectional and longitudinal evidence. *BMC Med* 2016; 14: 205.
53. Mante M, Franx A, Van Der Hoek YY, et al. Changes of plasma lipoprotein(a) during and after normal pregnancy in Caucasians. *J Matern Fetal Neonatal Med* 2003; 14: 91-5.
54. Lewek J, Bielecka-Dąbrowa A, Toth PP, Banach M. Dyslipidaemia management in pregnant patients: a 2024 update. *Eur Heart J Open* 2024; 4: oead032.
55. Golawski M, Lejawa M, Osadnik T, et al. Genetically determined lipoprotein(a) levels do not cause an increased risk of preeclampsia – a two-sample Mendelian randomization study. *Eur Heart J* 2023; 44 (Suppl. 2): ehad655.2728.
56. Sosnowska B, Stepinska J, Mitkowski P, et al. Recommendations of the Experts of the Polish Cardiac Society (PCS) and the Polish Lipid Association (PoLA) on the diagnosis and management of elevated lipoprotein(a) levels. *Arch Med Sci* 2024; 20: 8-27.
57. Enkhmaa B, Berglund L. Non-genetic influences on lipoprotein(a) concentrations. *Atherosclerosis* 2022; 349: 53-62.
58. Surma S, Sahebkar A, Banach M, endorsed by the International Lipid Expert Panel (ILEP). Low carbohydrate/ketogenic diet in the optimization of lipoprotein(a) levels: do we have sufficient evidence for any recommendation? *Eur Heart J* 2023; 44: 4904-6.
59. Gaba P, Rosenson RS, López JAG, et al. Intraindividual variability in serial lipoprotein(a) concentrations among placebo-treated patients in the OCEAN(a)-DOSE trial. *J Am Coll Cardiol* 2025; 85: 550-3.
60. Harb T, Ziogos E, Blumenthal RS, Gerstenblith G, Leucker TM. Intra-individual variability in lipoprotein(a): the value of a repeat measure for reclassifying individuals at intermediate risk. *Eur Heart J Open* 2024; 4: oead064.
61. Moriarty PM, Gorby LK, Stroes ES, Kastelein JP, Davidson M, Tsimikas S. Lipoprotein(a) and its potential association with thrombosis and inflammation in COVID-19: a testable hypothesis. *Curr Atheroscler Rep* 2020; 22: 48.
62. Müller N, Schulte DM, Türk K, et al. IL-6 blockade by monoclonal antibodies inhibits apolipoprotein (a) expression and lipoprotein (a) synthesis in humans. *J Lipid Res* 2015; 56: 1034-42.
63. Surma S, Sosnowska B, Reiner Ž, Banach M. New data allow to better understand the secrets of lipoprotein(a): is that for sure? *Eur Heart J Open* 2024; 4: oead066.
64. Sourij C, Aziz F, Krappinger S, et al. Changes in lipoprotein(a) levels in people after ST elevation myocardial infarction – the STEMI-Lipids Study. *Int J Mol Sci* 2023; 24: 15531.
65. Ziogos E, Vavuranakis MA, Harb T, et al. Lipoprotein(a) concentrations in acute myocardial infarction patients are not indicative of levels at six month follow-up. *Eur Heart J Open* 2023; 3: oead035.
66. Barth JH, Jackson BM, Farrin AJ, et al.; SPACE ROCKET Trial Group. Change in serum lipids after acute coronary syndromes: secondary analysis of SPACE ROCKET study data and a comparative literature review. *Clin Chem* 2010; 56: 1592-8.
67. Chiesa G, Zenti MG, Baragetti A, et al. Consensus document on lipoprotein(a) from the Italian Society for the Study of Atherosclerosis (SISA). *Nutr Metab Cardiovasc Dis* 2023; 33: 1866-77.
68. Durlach V, Bonnefont-Rousselot D, Boccara F, et al. Lipoprotein(a): pathophysiology, measurement, indication and treatment in cardiovascular disease. A consensus statement from the Nouvelle Société Francophone d'Athérosclérose (NSFA). *Arch Cardiovasc Dis* 2021; 114: 828-47.
69. Pearson GJ, Thanassoulis G, Anderson TJ, et al. 2021 Canadian Cardiovascular Society guidelines for the management of dyslipidemia for the prevention of cardiovascular disease in adults. *Can J Cardiol* 2021; 37: 1129-50.
70. Puri R, Nissen SE, Arsenault BJ, et al. Effect of C-reactive protein on lipoprotein(a)-associated cardiovascular risk in optimally treated patients with high-risk vascular disease: a prespecified secondary analysis of the ACCELERATE trial. *JAMA Cardiol* 2020; 5: 1136-43.
71. Li JJ, Ma CS, Zhao D, et al. Lipoprotein(a) and cardiovascular disease in Chinese population. *JACC Asia* 2022; 2: 653-65.
72. Koschinsky ML, Bajaj A, Boffa MB, et al. A focused update to the 2019 NLA scientific statement on use of lipoprotein(a) in clinical practice. *J Clin Lipidol* 2024; 18: e308-19.
73. Loh WJ, Pang J, Chakraborty A, et al. Cascade testing of children and adolescents for elevated Lp(a) in pedigrees with familial hypercholesterolemia. *J Clin Lipidol* 2024; 18: e33-7.

74. Patel AP, Wang M, Pirruccello JP, et al. Lp(a) (lipoprotein[a]) concentrations and incident atherosclerotic cardiovascular disease: new insights from a large National Biobank. *Arterioscler Thromb Vasc Biol* 2021; 41: 465-74.
75. Kamstrup PR. Genetically elevated lipoprotein(a) and increased risk of myocardial infarction. *JAMA* 2009; 301: 2331.
76. Burzynska M, Babicki M, Banach M, Jankowski P, Chudzik M. Lipoprotein(a) as a predictive marker of coronary artery calcification (CAC). The results from the STAR-Lp(a) cohort study. *Eur Heart J* 2024; 45 (Suppl. 1): ehae666.2829.
77. Bhatia HS, Hurst S, Desai P, Zhu W, Yeang C. Lipoprotein(a) testing trends in a large academic health system in the United States. *J Am Heart Assoc* 2023; 12: e031255.
78. Zafrir B, Aker A, Saliba W. Lipoprotein(a) testing in clinical practice: real-life data from a large healthcare provider. *Eur J Prev Cardiol* 2022; 29: e331-3.
79. Nurmohamed NS, Kaiser Y, Schuitema PCE, et al. Finding very high lipoprotein(a): the need for routine assessment. *Eur J Prev Cardiol* 2022; 29: 769-76.
80. Thanassoulis G, Campbell CY, Owens DS, et al. Genetic associations with valvular calcification and aortic stenosis. *N Engl J Med* 2013; 368: 503-12.
81. Langsted A, Nordestgaard BG, Kamstrup PR. Elevated lipoprotein(a) and risk of ischemic stroke. *J Am Coll Cardiol* 2019; 74: 54-66.
82. Masson W, Lobo M, Barbagelata L, Molinero G, Bluro I, Nogueira JP. Elevated lipoprotein (a) levels and risk of peripheral artery disease outcomes: a systematic review. *Vasc Med* 2022; 27: 385-91.
83. Kamstrup PR, Nordestgaard BG. Elevated lipoprotein(a) levels, LPA risk genotypes, and increased risk of heart failure in the general population. *JACC Heart Fail* 2016; 4: 78-87.
84. Lewek J, Sosnowska B, Bielecka-Dąbrowa A, Mierczak K, Adach W, Banach M. Lipoprotein(a) and other risk factors in dyslipidemic patients with and without heart failure. *Arch Med Sci* 2024; 20: 1705-9.
85. Singh S, Baars DP, Desai R, Singh D, Pinto-Sietsma SJ. Association between lipoprotein(a) and risk of atrial fibrillation: a systematic review and meta-analysis of Mendelian randomization studies. *Curr Probl Cardiol* 2024; 49: 102024.
86. Grant JK, Martin SS, Zhang S, et al. Racial differences in the burden of atherosclerotic cardiovascular disease related to elevated lipoprotein(a) levels: the ARIC study. *Circulation* 2024; 150: 250-2.
87. Ridker PM, Moorthy MV, Cook NR, Rifai N, Lee IM, Buring JE. Inflammation, cholesterol, lipoprotein(a), and 30-year cardiovascular outcomes in women. *N Engl J Med* 2024; 391: 2087-97.
88. Bhatia HS, Wandel S, Willeit P, et al. Independence of lipoprotein(a) and low-density lipoprotein cholesterol-mediated cardiovascular risk: a participant-level meta-analysis. *Circulation* 2025; 151: 312-21.
89. Ridker PM. A test in context. *J Am Coll Cardiol* 2016; 67: 712-23.
90. Small AM, Pournamdari A, Melloni GEM, et al. Lipoprotein(a), C-reactive protein, and cardiovascular risk in primary and secondary prevention populations. *JAMA Cardiol* 2024; 9: 385.
91. Thomas PE, Vedel-Krogh S, Kamstrup PR, Nordestgaard BG. Lipoprotein(a) is linked to atherothrombosis and aortic valve stenosis independent of C-reactive protein. *Eur Heart J* 2023; 44: 1449-60.
92. Arnold N, Blaum C, Goßling A, et al. C-reactive protein modifies lipoprotein(a)-related risk for coronary heart disease: the BiomarCaRE project. *Eur Heart J* 2024; 45: 1043-54.
93. Zhang W, Speiser JL, Ye F, et al. High-sensitivity C-reactive protein modifies the cardiovascular risk of lipoprotein(a). *J Am Coll Cardiol* 2021; 78: 1083-94.
94. Yuan D, Wang P, Jia S, et al. Lipoprotein(a), high-sensitivity C-reactive protein, and cardiovascular risk in patients undergoing percutaneous coronary intervention. *Atherosclerosis* 2022; 363: 109-16.
95. Alebna PL, Han CY, Ambrosio M, et al. Association of lipoprotein(a) with major adverse cardiovascular events across hs-CRP. *JACC Adv* 2024; 3: 101409.
96. Chung YH, Lee BK, Kwon HM, et al. Coronary calcification is associated with elevated serum lipoprotein (a) levels in asymptomatic men over the age of 45 years: a cross-sectional study of the Korean national health checkup data. *Medicine (Baltimore)* 2021; 100: e24962.
97. Mehta A, Vasquez N, Ayers CR, et al. Independent association of lipoprotein(a) and coronary artery calcification with atherosclerotic cardiovascular risk. *J Am Coll Cardiol* 2022; 79: 757-68.
98. Kaiser Y, Daghm M, Tzolos E, et al. Association of lipoprotein(a) with atherosclerotic plaque progression. *J Am Coll Cardiol* 2022; 79: 223-33.
99. Qiu Y, Hao W, Guo Y, et al. The association of lipoprotein (a) with coronary artery calcification: a systematic review and meta-analysis. *Atherosclerosis* 2024; 388: 117405.
100. Zheng KH, Tsimikas S, Pawade T, et al. Lipoprotein(a) and oxidized phospholipids promote valve calcification in patients with aortic stenosis. *J Am Coll Cardiol* 2019; 73: 2150-62.
101. Khan MI, Zahir RS, Dominguez AC, Romeo FJ. Role of Lipoprotein (A) in aortic valve stenosis: novel disease mechanisms and emerging pharmacotherapeutic approaches. *IJC Heart Vasc* 2024; 55: 101543.
102. Arsenault BJ, Loganath K, Girard A, et al. Lipoprotein(a) and calcific aortic valve stenosis progression: a systematic review and meta-analysis. *JAMA Cardiol* 2024; 9: 835-42.
103. Surma S, Zembala MO, Okopień B, Banach M. Lipoprotein (a) and lipid-lowering treatment from the perspective of a cardiac surgeon. An impact on the prognosis in patients with aortic valve replacement and after heart transplantation. *Int J Cardiol Cardiovasc Risk Prev* 2024; 22: 200297.
104. Kim AR, Ahn JM, Kang DY, et al. Association of lipoprotein(a) with severe degenerative aortic valve stenosis. *JACC Asia* 2024; 4: 751-60.
105. Tsimikas S. Lp(a) and accelerated progression of aortic stenosis. *JACC Asia* 2024; 4: 761-3.
106. Wu B, Zhang Z, Long J, Zhao H, Zeng F. Association between lipoprotein (a) and heart failure with reduced ejection fraction development. *J Clin Lab Anal* 2022; 36: e24083.
107. Singh S, Baars DP, Aggarwal K, Desai R, Singh D, Pinto-Sietsma SJ. Association between lipoprotein (a) and risk of heart failure: a systematic review and meta-analysis of Mendelian randomization studies. *Curr Probl Cardiol* 2024; 49: 102439.
108. Januzzi JL, Van Kimmenade RRI, Liu Y, et al. Lipoprotein(a), oxidized phospholipids, and progression to

- symptomatic heart failure: the CASABLANCA study. *J Am Heart Assoc* 2024; 13: e034774.
109. Zheng W, Chilazi M, Park J, et al. Assessing the accuracy of estimated lipoprotein(a) cholesterol and lipoprotein(a)-free low-density lipoprotein cholesterol. *J Am Heart Assoc* 2022; 11: e023136.
 110. Solnica B, Sygitowicz G, Sitkiewicz D, et al. 2024 Guidelines of the Polish Society of Laboratory Diagnostics and the Polish Lipid Association on laboratory diagnostics of lipid metabolism disorders. *Arch Med Sci* 2024; 20: 357-74.
 111. Arnold N, Blaum C, Goßling A, et al. Impact of lipoprotein(a) level on low-density lipoprotein cholesterol- or apolipoprotein b-related risk of coronary heart disease. *J Am Coll Cardiol* 2024; 84: 165-77.
 112. Tsimikas S, Yeang C, Kronenberg F. In search of an accurate measurement of LDL-C. *J Am Coll Cardiol* 2024; 84: 178-81.
 113. Banach M, Surma S, Toth PP; endorsed by the International Lipid Expert Panel (ILEP). 2023: The year in cardiovascular disease – the year of new and prospective lipid lowering therapies. Can we render dyslipidemia a rare disease by 2024? *Arch Med Sci* 2023; 19: 1602-15.
 114. Banach M, Burchardt P, Chlebus K, et al. PoLA/CFPiP/PCS/PSLD/PSD/PSH guidelines on diagnosis and therapy of lipid disorders in Poland 2021. *Arch Med Sci* 2021; 17: 1447-547.
 115. Yahya R, Berk K, Verhoeven A, et al. Statin treatment increases lipoprotein(a) levels in subjects with low molecular weight apolipoprotein(a) phenotype. *Atherosclerosis* 2019; 289: 201-5.
 116. Tsimikas S, Gordts PLSM, Nora C, Yeang C, Witztum JL. Statin therapy increases lipoprotein(a) levels. *Eur Heart J* 2020; 41: 2275-84.
 117. Banach M, Penson PE. Statins and Lp(a): do not make perfect the enemy of excellent. *Eur Heart J* 2020; 41: 190-1.
 118. Banach M, Surma S, Kapłon-Cieślicka A, et al. Position paper of the Polish Expert Group on the use of pitavastatin in the treatment of lipid disorders in Poland endorsed by the Polish Lipid Association. *Arch Med Sci* 2024; 20: 28-42.
 119. Sahebkar A, Simental-Mendía LE, Pirro M, et al. Impact of ezetimibe on plasma lipoprotein(a) concentrations as monotherapy or in combination with statins: a systematic review and meta-analysis of randomized controlled trials. *Sci Rep* 2018; 8: 17887.
 120. Awad K, Mikhailidis DP, Katsiki N, Muntner P, Banach M. Effect of ezetimibe monotherapy on plasma lipoprotein(a) concentrations in patients with primary hypercholesterolemia: a systematic review and meta-analysis of randomized controlled trials. *Drugs* 2018; 78: 453-62.
 121. Ridker PM, Lei L, Ray KK, Ballantyne CM, Bradwin G, Rifai N. Effects of bempedoic acid on CRP, IL-6, fibrinogen and lipoprotein(a) in patients with residual inflammatory risk: a secondary analysis of the CLEAR harmony trial. *J Clin Lipidol* 2023; 17: 297-302.
 122. Chemello K, Chan DC, Lambert G, Watts GF. Recent advances in demystifying the metabolism of lipoprotein(a). *Atherosclerosis* 2022; 349: 82-91.
 123. The HPS2-THRIVE Collaborative Group. Effects of extended-release niacin with laropirant in high-risk patients. *N Engl J Med* 2014; 371: 203-12.
 124. Haynes R, Valdes-Marquez E, Hopewell JC, et al. Serious adverse effects of extended-release niacin/laropirant: results from the Heart Protection Study 2–Treatment of HDL to Reduce the Incidence of Vascular Events (HPS2-THRIVE) Trial. *Clin Ther* 2019; 41: 1767-77.
 125. O'Donoghue ML, Fazio S, Giugliano RP, et al. Lipoprotein(a), PCSK9 inhibition, and cardiovascular risk: insights from the FOURIER Trial. *Circulation* 2019; 139: 1483-92.
 126. Bittner VA, Szarek M, Aylward PE, et al. Effect of alirocumab on lipoprotein(a) and cardiovascular risk after acute coronary syndrome. *J Am Coll Cardiol* 2020; 75: 133-44.
 127. Toth PP, Jones SR, Monsalvo ML, Elliott-Davey M, López JAG, Banach M. Effect of evolocumab on non-high-density lipoprotein cholesterol, apolipoprotein b, and lipoprotein(a): a pooled analysis of phase 2 and phase 3 studies. *J Am Heart Assoc* 2020; 9: e014129.
 128. Ray KK, Kallend D, Leiter LA, et al. Effect of inclisiran on lipids in primary prevention: the ORION-11 trial. *Eur Heart J* 2022; 43: 5047-57.
 129. Katsiki N, Vrablik M, Banach M, Gouni-Berthold I. Inclisiran, low-density lipoprotein cholesterol and lipoprotein (a). *Pharmaceuticals (Basel)* 2023; 16: 577.
 130. Blanchard V, Chemello K, Hollstein T, et al. The size of apolipoprotein (a) is an independent determinant of the reduction in lipoprotein (a) induced by PCSK9 inhibitors. *Cardiovasc Res* 2022; 118: 2103-11.
 131. Akinlonu A, Boffa MB, Lyu C, et al. Variation in lipoprotein(a) response to potent lipid lowering: the role of apolipoprotein (a) isoform size. *J Clin Lipidol* 2025; 19: 39-50.
 132. Connelly Smith L, Alquist CR, Aqui NA, et al. Guidelines on the use of therapeutic apheresis in clinical practice – evidence based approach from the writing committee of the American Society for Apheresis: the ninth special issue. *J Clin Apheresis* 2023; 38: 77-278.
 133. Safarova MS, Moriarty PM. Lipoprotein apheresis: current recommendations for treating familial hypercholesterolemia and elevated lipoprotein(a). *Curr Atheroscler Rep* 2023; 25: 391-404.
 134. Waldmann E, Parhofer KG. Lipoprotein apheresis to treat elevated lipoprotein (a). *J Lipid Res* 2016; 57: 1751-7.
 135. Stefanutti C, Mazza F, Pasqualetti D, et al. Lipoprotein apheresis downregulates IL-1 α , IL-6 and TNF- α mRNA expression in severe dyslipidaemia. *Atheroscler Suppl* 2017; 30: 200-8.
 136. Leebmann J, Roeseler E, Julius U, et al. Lipoprotein apheresis in patients with maximally tolerated lipid-lowering therapy, lipoprotein(a)-hyperlipoproteinemia, and progressive cardiovascular disease: prospective observational multicenter study. *Circulation* 2013; 128: 2567-76.
 137. Lipoprotein Apheresis: First FDA Indicated Treatment for Elevated Lipoprotein(a). *J Clin Cardiol [Internet]*. 15 października 2020 [cytowane 8 stycznia 2025];1(1). Available: <https://www.scientificarchives.com/article/lipoprotein-apheresis-first-fda-indicated-treatment-for-elevated-lipoprotein>
 138. Kayikcioglu M. LDL Apheresis and Lp (a) apheresis: a clinician's perspective. *Curr Atheroscler Rep* 2021; 23: 15.
 139. The ASCEND Study Collaborative Group. Effects of aspirin for primary prevention in persons with diabetes mellitus. *N Engl J Med* 2018; 379: 1529-39.
 140. Gaziano JM, Brotons C, Coppolecchia R, et al. Use of aspirin to reduce risk of initial vascular events in patients at moderate risk of cardiovascular disease (ARRIVE):

- a randomised, double-blind, placebo-controlled trial. *Lancet* 2018; 392: 1036-46.
141. McNeil JJ, Wolfe R, Woods RL, et al. Effect of aspirin on cardiovascular events and bleeding in the healthy elderly. *N Engl J Med* 2018; 379: 1509-18.
 142. Zheng SL, Roddick AJ. Association of aspirin use for primary prevention with cardiovascular events and bleeding events: a systematic review and meta-analysis. *JAMA* 2019; 321: 277-87.
 143. Lacaze P, Bakshi A, Riaz M, et al. Aspirin for primary prevention of cardiovascular events in relation to lipoprotein(a) genotypes. *J Am Coll Cardiol* 2022; 80: 1287-98.
 144. Bhatia HS, Trainor P, Carlisle S, et al. Aspirin and cardiovascular risk in individuals with elevated lipoprotein(a): the multi ethnic study of atherosclerosis. *J Am Heart Assoc* 2024; 13: e033562.
 145. Chasman DI, Shiffman D, Zee RYL, et al. Polymorphism in the apolipoprotein(a) gene, plasma lipoprotein(a), cardiovascular disease, and low-dose aspirin therapy. *Atherosclerosis* 2009; 203: 371-6.
 146. Razavi AC, Richardson LC, Coronado F, et al. Aspirin use for primary prevention among US adults with and without elevated lipoprotein(a). *Am J Prev Cardiol* 2024; 18: 100674.
 147. Patel SM, Bonaca MP, Morrow DA, et al. Lipoprotein(a) and benefit of antiplatelet therapy: insights from the PEGASUS-TIMI 54 Trial. *JACC Adv* 2023; 2: 100675.
 148. Della Bona R, Giubilato S, Palmieri M, et al. Aspirin in primary prevention: looking for those who enjoy it. *J Clin Med* 2024; 13: 4148.
 149. Banach M, Surma S, Bielecka-Dąbrowa A, et al. Rosuvastatin-based combination treatment with acetylsalicylic acid or ezetimibe in the management of patients at high and very high cardiovascular risk. Expert opinion paper of the Polish Lipid Association 2025. *Arch Med Sci* 2025; 21: 1-15.
 150. Sosnowska B, Surma S, Banach M. Targeted treatment against lipoprotein (a): the coming breakthrough in lipid lowering therapy. *Pharmaceuticals* 2022; 15: 1573.
 151. O'Donoghue ML, Rosenson RS, López JAG, et al. The off-treatment effects of olpasiran on lipoprotein(a) lowering. *J Am Coll Cardiol* 2024; 84: 790-7.
 152. Nissen SE, Wang Q, Nicholls SJ, et al. Zerlasiran – a small-interfering RNA targeting lipoprotein(a): a phase 2 randomized clinical trial. *JAMA* 2024; 332: 1992.
 153. Nissen SE, Linnebjerg H, Shen X, et al. Lepodisiran, an extended-duration short interfering RNA targeting lipoprotein(a): a randomized dose-ascending clinical trial. *JAMA* 2023; 330: 2075-83.
 154. Kulkarni JA, Witzigmann D, Thomson SB, et al. The current landscape of nucleic acid therapeutics. *Nat Nanotechnol* 2021; 16: 630-43.
 155. Novartis Pharmaceuticals. A Randomized Double-blind, Placebo-controlled, Multicenter Trial Assessing the Impact of Lipoprotein (a) Lowering With Pelacarsen (TQJ230) on Major Cardiovascular Events in Patients With Established Cardiovascular Disease [Internet]. *clinicaltrials.gov*; 2025 [accessed 12 January 2025]. Report No.: NCT04023552. Available at: <https://clinicaltrials.gov/study/NCT04023552>
 156. Novartis Pharmaceuticals. A Randomized Double-blind, Placebo-controlled, Multicenter Trial Assessing the Impact of Lipoprotein(a) Lowering With Pelacarsen (TQJ230) on the Progression of Calcific Aortic Valve Stenosis [Lp(a)FRONTIERS CAVS] [Internet]. *clinicaltrials.gov*; 2024 [accessed 12 January 2025]. Report No.: NCT05646381. Available at: <https://clinicaltrials.gov/study/NCT05646381>
 157. Amgen. A Double-blind, Randomized, Placebo-controlled, Multicenter Study Assessing the Impact of Olpasiran on Major Cardiovascular Events in Participants With Atherosclerotic Cardiovascular Disease and Elevated Lipoprotein(a) [Internet]. *clinicaltrials.gov*; 2024 [accessed 12 January 2025]. Report No.: NCT05581303. Available at: <https://clinicaltrials.gov/study/NCT05581303>
 158. Eli Lilly and Company. A Phase 2, Randomized, Double-Blind, Placebo-Controlled Study to Investigate the Efficacy and Safety of LY3819469 in Adults With Elevated Lipoprotein(a) [Internet]. *clinicaltrials.gov*; 2025 [accessed 12 January 2025]. Report No.: NCT05565742. Available at: <https://clinicaltrials.gov/study/NCT05565742>
 159. Eli Lilly and Company. A Phase 3, Randomized, Double-Blind, Placebo-Controlled Study to Investigate the Effect of Lepodisiran on the Reduction of Major Adverse Cardiovascular Events in Adults With Elevated Lipoprotein(a) Who Have Established Atherosclerotic Cardiovascular Disease or Are at Risk for a First Cardiovascular Event - ACCLAIM-Lp(a) [Internet]. *clinicaltrials.gov*; 2024 [accessed 12 January 2025]. Report No.: NCT06292013. Available at: <https://clinicaltrials.gov/study/NCT06292013>
 160. Nicholls SJ, Nissen SE, Fleming C, et al. Muvalaplin, an oral small molecule inhibitor of lipoprotein(a) formation: a randomized clinical trial. *JAMA* 2023; 330: 1042.
 161. Morrow PK, D'Souza S, Wood T, et al. Abstract 17013: CTX320: An Investigational in vivo CRISPR-Based Therapy Efficiently and Durably Reduces Lipoprotein (a) Levels in Non-Human Primates After a Single Dose. *Circulation* 2023; 148 (Suppl_1). https://www.ahajournals.org/doi/10.1161/circ.148.suppl_1.17013
 162. Nicholls SJ. Therapeutic potential of lipoprotein(a) inhibitors. *Drugs* 2024; 84: 637-43.
 163. Nicholls SJ, Ni W, Rhodes GM, et al. Oral muvalaplin for lowering of lipoprotein(a): a randomized clinical trial. *JAMA* 2025; 333: 222-31.
 164. CTX320: An Investigational In Vivo CRISPR-Based Therapy Efficiently And Durably Reduces Lipoprotein(a) Levels In Non-Human Primates After A Single Dose. [Internet]. Available at: <https://ir.crisprtx.com/static-files/db310adc-944e-4d9a-9d68-5ab358716c2e>
 165. Nicholls SJ, Ditmarsch M, Kastelein JJ, et al. Lipid lowering effects of the CETP inhibitor obicetrapib in combination with high-intensity statins: a randomized phase 2 trial. *Nat Med* 2022; 28: 1672-8.
 166. Tsimikas S, Yang X, Hsieh A, Ditmarsch M, Davidson M, Kastelein J. Obicetrapib demonstrates significant reductions of Lp(a) on top of high-intensity statins. *Atherosclerosis* 2024; 395: 118412.
 167. Nicholls S, Nelson A, Ditmarsch M, et al. 4171281 Safety and efficacy of obicetrapib in patients with heterozygous familial hypercholesterolemia. *Circulation* 2024; 150: <https://doi.org/10.1161/CIR.0000000000001299>
 168. NewAmsterdam Pharma. A Phase 2 Study to Evaluate the Effect of Obicetrapib Alone and in Combination With Evolocumab (Repatha®) on Lipoprotein (a) in Patients With Mild Dyslipidemia: The VINCENT Study [Internet]. *clinicaltrials.gov*; 2024 [accessed January 2025]. Report No.: NCT06496243. Available at: <https://clinicaltrials.gov/study/NCT06496243>
 169. Gudbjartsson DF, Thorgeirsson G, Sulem P, et al. Lipoprotein(a) concentration and risks of cardiovascu-

- lar disease and diabetes. *J Am Coll Cardiol* 2019; 74: 2982-94.
170. Kamstrup PR, Nordestgaard BG. Lipoprotein(a) concentrations, isoform size, and risk of type 2 diabetes: a Mendelian randomisation study. *Lancet Diabetes Endocrinol* 2013; 1: 220-7.
 171. Paige E, Masconi KL, Tsimikas S, et al. Lipoprotein(a) and incident type-2 diabetes: results from the prospective Bruneck study and a meta-analysis of published literature. *Cardiovasc Diabetol* 2017; 16: 38.
 172. Tsimikas S. In search of a physiological function of lipoprotein(a): causality of elevated Lp(a) levels and reduced incidence of type 2 diabetes. *J Lipid Res* 2018; 59: 741-4.
 173. Farukhi ZM, Mora S. Lifelong low Lp(a) levels: genetics give a green light? *Eur Heart J* 2021; 42: 1157-9.
 174. Ye Z, Haycock PC, Gurdasani D, et al. The association between circulating lipoprotein(a) and type 2 diabetes: is it causal? *Diabetes* 2014; 63: 332-42.
 175. Gołowski M, Lejawa M, Banach M, Jóźwiak J, Gierlotka M, Osadnik T. Association between Lp(a) and T2D: a Mendelian randomization study. *Arch Med Sci* 2024; 20: 1002-5.
 176. Yeung MW, Said MA, Van De Vegte YJ, Verweij N, Dullaart RPF, Van Der Harst P. Associations of very low lipoprotein(a) levels with risks of new-onset diabetes and non-alcoholic liver disease. *Atheroscler Plus* 2024; 57: 19-25.
 177. Lejawa M, Gołowski M, Fronczek M, et al. Causal associations between insulin and Lp(a) levels in Caucasian population: a Mendelian randomization study. *Cardiovasc Diabetol* 2024; 23: 316.
 178. Sabouret P, Dib J, Ecarnot F, Banach M, Lellouche N. Eligibility for icosapent ethyl in a French population of coronary outpatients with type 2 diabetes. *Arch Med Sci* 2023; 19: 1901-3.
 179. Koniecznyńska M, Natarska J, Ząbczyk M, Undas A. Lipoprotein(a) and thromboembolism: current state of knowledge and unsolved issues. *Arch Med Sci* 2024; 20: 1770-83.