

Genetic causal associations between serum metabolites and infertility: a Mendelian randomization study

Jiahao Shi¹, Jiamei Zhu², Tian Tao^{3*}

¹Department of Urology, The Affiliated Wuxi People's Hospital of Nanjing Medical University, Wuxi, China

²Department of Obstetrics and Gynecology, Jingjiang People's Hospital Affiliated to Yangzhou University, Taizhou, China

³Department of Obstetrics and Gynecology, The First Affiliated Hospital of Soochow University, Suzhou, China

Submitted: 28 October 2024; **Accepted:** 22 April 2025

Online publication: 8 June 2025

Arch Med Sci 2026; 22 (3): 1794–1803

DOI: <https://doi.org/10.5114/aoms/204248>

Copyright © 2025 Termedia & Banach

***Corresponding author:**

Tian Tao

Department of Obstetrics and Gynecology

The First Affiliated

Hospital of

Soochow University

Suzhou, China

E-mail: taotian_1123@163.com

com

Abstract

Introduction: This study aimed to elucidate the causal relationships between serum metabolites and infertility in both men and women, and to identify key metabolic biomarkers.

Material and methods: This study employed a two-sample Mendelian randomization design, with circulating plasma metabolite genome-wide association study data as an exposure factor and FinnGen Consortium R10 genome-wide association study data for infertility in men and women as an outcome. The causal relation between plasma metabolites and infertility in men and women was assessed using five methods: inverse variance weighted, Egger regression, weighted median, maximum likelihood estimation, and simple mode.

Results: This analysis identified 17 and 10 metabolites positively and negatively associated with infertility in women, respectively. Similarly, 22 and 30 metabolites were positively and negatively associated with infertility in men, respectively. Galactonate and glycerate levels were identified as risk factors for infertility in both men and women. In addition, sphingomyelin exerts protective effects against infertility in both men and women. Metabolic pathway analysis revealed enrichment of critical metabolic pathways related to infertility.

Conclusions: This study identified several circulating metabolic biomarkers associated with infertility. These biomarkers can be used for the screening and prevention of infertility. In addition, they could be employed as candidate molecules for future mechanistic exploration and drug-targeting studies.

Key words: infertility, circulating plasma metabolites, Mendelian randomization.

Introduction

Infertility, a prevalent reproductive system disorder with rising global incidence, impacts individual well-being, family dynamics, and societal demographics. It is defined as the failure to achieve clinical pregnancy after 12 months of regular unprotected intercourse [1]. It affects 14.3% and 25% of reproductive-age couples in developed and developing countries, respectively [2]. Notably, male factors contribute to nearly

50% of cases, highlighting the need for early prevention, diagnosis, and targeted management strategies.

Integrative omics (transcriptomics, proteomics, metabolomics) has emerged as a pivotal tool in life sciences. Evidence indicates that metabolic disorders (e.g. hypertension, diabetes, obesity) are strongly linked to infertility in both sexes [3, 4]. Metabolomics, extending beyond genomics and proteomics [5, 6], identifies metabolic biomarkers and elucidates pathogenic pathways by comparing circulating metabolite profiles between infertile patients and healthy controls, offering novel therapeutic targets for infertility prevention and treatment.

Mendelian randomization (MR) is an epidemiological method that uses single-nucleotide polymorphisms (SNPs) as instrumental variables (IVs) to infer causal relationships between exposures and diseases, circumventing limitations of observational studies [7]. Leveraging genome-wide association study (GWAS) data, we applied a two-sample MR approach to investigate causal effects of serum metabolites on infertility and identify underlying metabolic pathways.

Material and methods

Study design

We used a two-sample MR approach based on GWAS summary statistics to investigate causal associations between circulating metabolites in humans and the risk of infertility in both men and women. This is a genetic epidemiological approach for investigating causality and addressing confounders by using IVs under three assumptions: (i) IVs are not associated with confounders; (ii) IVs are associated with exposure factors; and (iii) IVs are not associated with outcome variables, with IVs influencing outcomes only through exposure. Genetic information on circulating metabolites and infertility in humans was obtained from separate GWAS databases. The flowchart of this MR study is shown in Figure 1.

GWAS data

Chen *et al.* obtained a genome-wide association database of 870 blood metabolites [8]. This is the most comprehensive analysis of blood metabolites in humans and is included in the GWAS




Figure 1. Flowchart of the study design in our Mendelian randomization (MR) analysis

catalog. After applying the appropriate exclusion criteria, the Canadian Longitudinal Study of Aging finalized the identification of 8,299 participants between the ages of 45 and 85 years and performed genome-wide typing, identifying 248 linked loci. They were also tested for 1,091 circulating plasma metabolites and 309 metabolite ratios. Of the 1,091 metabolites, 850 had a known identity (including lipids, amino acids, xenobiotics, nucleotides, cofactors and vitamins, carbohydrates, peptides, and energy-related molecules), whereas the remaining 241 were defined as unknown or partially characterized molecules; metabolite ratios and unknown molecules were excluded from this study. Information on these metabolites is provided in Supplementary Table S1.

FinnGen is a large public-private cooperation project covering the entire territory of Finland to determine the etiology of various diseases and assess the role of genetics in health and disease. We downloaded GWAS data related to infertility in both men and women from the R10 version of the FinnGen consortium, which yielded 14,759 infertility cases in women, 111,583 controls, 1,429 infertility cases in men, and 130,139 controls (<https://r10.finnngen.fi/>).

IV selection

In this study, we selected SNPs as genetic IVs. We rigorously screened the IVs associated with serum metabolites using the following steps to ensure the robustness of the MR analyses (Figure 1). First, SNPs with minor allele frequency (MAF) > 0.01 in exposure and outcome were selected, followed by screening for SNPs satisfying the locus-wide significance threshold ($p < 1 \times 10^{-5}$) associated with exposure. Subsequently, SNPs were removed by performing the LD clumping procedure with $r^2 < 0.001$ and a window size = 10,000 kb to eliminate variants in linkage disequilibrium. Genetic variants with $F > 10$ were selected as strong IVs to avoid bias from weak IVs [9].

MR analysis

Five methods were used to evaluate the causal relation between exposure (plasma metabolites) and outcome (infertility), with the random-effects inverse variance weighted (IVW) model as the primary analytical method, along with the MR-Egger, weighted median, maximum likelihood, and simple mode methods [10]. If only one IV was available for the plasma metabolites, the Wald ratio was used. If two or more IVs were present, the IVW method was used. Results with significant p -values for the IVW method and β values for the remaining four methods in the same direction as the IVW method were considered meaningful. In

the heterogeneity test, Cochran's Q statistic had a p -value > 0.05, indicating no heterogeneity between SNPs at the statistical level. Second, horizontal pleiotropy was assessed using MR-Egger regression. A p -value < 0.05 for the intercept term indicated the presence of statistically significant horizontal pleiotropy. We detected reverse causality using the Steiger test of directionality, with TRUE indicating the absence of statistically significant reverse causality. The results show that all MR methods yielded results in the same direction and that the MR-Egger method did not detect statistically significant pleiotropy, supporting our main finding. Statistical analyses were conducted using R software (version 4.2.3), and MR analysis was performed using the TwoSampleMR package.

Metabolic pathway analysis

The RaMP database of Web-based Metaconflict 6.0 was used to analyze metabolic pathways leading to infertility (<https://www.metabcanalest.ca/>), and pathways with FDR < 0.05 were considered significantly enriched.

Results

MR results for infertility in men and women

After excluding unknown metabolites and metabolite ratios, 870 plasma metabolite-associated SNPs with known structures and functions were used. The number of IVs selected for plasma metabolites ranged from 3 to 32, and we derived causal associations between 870 metabolites and infertility in both men and women (Supplementary Tables SII and SIII). The IVW method showed a positive correlation between increased levels of 17 metabolites and the risk of infertility in women, including eight from the lipid metabolism pathway, three from the amino acid metabolism pathway, two from the cofactor and vitamin pathways, two from the carbohydrate pathway, one from the xenobiotic pathway, and one from a partially characterized molecular pathway. The top five of these metabolites were as follows: 7-methylxanthine (odds ratio [OR] = 1.187, 95% confidence interval [CI] = 1.076–1.311, $p = 0.0007$); octadecadienedioate (C18:2-DC) (OR = 1.086, 95% CI = 1.041–1.134, $p = 0.0002$); galactonate (OR = 1.192, 95% CI = 1.069–1.329, $p = 0.0016$); metabolonic lactone sulfate (OR = 1.067, 95% CI = 1.026–1.109, $p = 0.0011$); sphingomyelin (d18:0/18:0, d19:0/17:0) (OR = 1.112, 95% CI = 1.037–1.193, $p = 0.0030$). Furthermore, we observed 10 metabolites that were negatively associated with infertility in women. The top five of these were oleate/vaccenate (18:1) (OR = 0.865, 95% CI = 0.784–0.954, $p = 0.0037$); glyoursodeoxycholate (OR = 0.892, 95% CI = 0.827–




Figure 2. Forest plot of Mendelian randomization (MR) analysis between plasma metabolites and infertility in women
IVW – inverse variance weighted, SNP – single nucleotide polymorphism, OR – odds ratio, CI – confidence interval.

0.963, $p = 0.0032$); sphingomyelin (d18:2/23:1) (OR = 0.894, 95% CI = 0.819–0.975, $p = 0.0116$); glycocholate (OR = 0.923, 95% CI = 0.862–0.987, $p = 0.0196$); glycohyocholate (OR = 0.869, 95% CI = 0.772–0.979, $p = 0.0208$) (Figure 2 and Supplementary Table SIV).

A total of 22 metabolites were identified as positively associated with infertility in men. The top five metabolites were as follows: 1-oleoyl-GPI (18:1) (OR = 1.429, 95% CI = 1.151–1.773, $p = 0.0012$); 11 β -hydroxyandrosterone glucuronide (OR = 1.382, 95% CI = 1.134–1.683, $p = 0.0013$); tauro- β -muricholate (OR = 1.178, 95% CI = 1.051–1.319, $p = 0.0048$); 4-vinylguaiaicol sulfate (OR = 1.408, 95% CI = 1.110–1.788, $p = 0.0049$); trans-4-hydroxyproline (OR = 1.372, 95% CI = 1.095–1.718, $p = 0.0059$). Concurrently, 30 metabolites were negatively associated with infertility in men. Of these, the top five were 5 α -androstan-3 β , 17 β -diol monosulfate (2) (OR = 0.852, 95% CI = 0.767–0.947, $p = 0.0029$); 1-(1-enyl-palmitoyl)-2-linoleoyl-GPE (p-16:0/18:2) (OR = 0.742, 95% CI = 0.608–0.904, $p = 0.0032$); sphingomyelin (d18:1/24:1, d18:2/24:0) (OR = 0.659, 95% CI = 0.498–0.873, $p = 0.0036$); carnitine C14 (OR = 0.704, 95% CI = 0.543–0.914, $p = 0.0083$); androsterone sulfate (OR = 0.903, 95% CI

= 0.835–0.976, $p = 0.0100$) (Figure 3 and Supplementary Table SV).

Infertility in men and women shares several causal metabolites. Pyridoxal was positively associated with infertility in both women (OR = 1.096, 95% CI = 1.027–1.170, $p = 0.0058$) and men (OR = 1.257, 95% CI = 1.042–1.518, $p = 0.0171$). Galactonate and glycerate levels were risk factors for infertility in both men and women. Furthermore, sphingomyelin exerted a protective effect against infertility and was inversely correlated with infertility in women (d18:2/23:1). By contrast, in infertility in men, six sphingomyelin metabolites were identified: sphingomyelin (d18:0/18:0, d19:0/17:0); sphingomyelin (d18:0/20:0, d16:0/22:0); sphingomyelin (d18:1/17:0, d17:1/18:0, d19:1/16:0); sphingomyelin (d18:1/18:1, d18:2/18:0); sphingomyelin (d18:1/20:2, d18:2/20:1, d16:1/22:2); sphingomyelin (d18:1/24:1, d18:2/24:0).

Sensitivity analysis

The directionality of the MR results was consistent across all methods (Supplementary Figures S1 and S2). MR-Egger results showed that no horizontal pleiotropy was detected for 27 metabolites associated with infertility in women and




Figure 3. Forest plot of Mendelian randomization (MR) analysis between plasma metabolites and infertility in men

IVW – inverse variance weighted, SNP – single nucleotide polymorphism, OR – odds ratio, CI – confidence interval.

52 metabolites related to infertility in men (Supplementary Table S1). Scatter plots depict the effect of each SNP locus on infertility, with the slopes representing the magnitude of impact. Leave-one-out analysis showed no significant outliers (Supplementary Figures S3 and S4).

Metabolic pathway analysis

A study of metabolic pathways revealed that the five most prevalent metabolic pathways in infertility in women were glycerolipid metabolism, glycine and serine metabolism, valine/leucine/iso-

leucine degradation, bile acid biosynthesis, and thiamine metabolism (Figure 4 A). The five metabolic pathways strongly associated with infertility in men were de novo triacylglycerol biosynthesis, malate-aspartate shuttle, glycerol phosphate shuttle, cardiolipin biosynthesis, and bile acid biosynthesis. Additionally, five metabolic pathways were common to infertility in both men and women (Figure 4 B): bile acid biosynthesis, vitamin B6 metabolism, glycerolipid metabolism, transfer of acetyl groups into mitochondria, and valine/leucine/isoleucine degradation.




Figure 4. Enriched significant metabolic pathways of infertility. **A** – Significant metabolic pathways involved in infertility in women. **B** – Significant enrichment of metabolic pathways in men with infertility

Discussion

In this two-sample MR analysis utilizing large-scale GWAS data, we systematically investigated the potential causal associations between 871 plasma metabolites and infertility in both men and women. Multiple risk and protective factors associated with infertility, as well as key metabolic pathways, were identified. The results provide new evidence and insights into infertility pathogenesis and potential targets for prevention and treatment. Metabolomics, a widely used method for studying reproduction-related diseases, effectively identifies biomarkers in blood, semen, and follicular fluid. Among these, blood samples are particularly valuable because of their accessibility and rich metabolic profile. This study not only validated characteristic metabolites but also further elucidated key metabolic pathways leading to infertility, laying an important foundation for a deeper understanding of infertility's molecular mechanisms and the development of precise therapeutic strategies.

A decline in semen quality is a major cause of infertility in men. In some countries, semen quality problems have become a primary issue for young men with fertility problems [11]. One study found differences in the serum metabolic profiles of men with different sperm concentrations and identified the markers as peptides related to the protein complement C3f [12]. Zhang *et al.* identified 24 potential biological markers of serum metabolites in patients with azoospermia and in healthy controls. Taurine exhibits high levels in healthy individuals and is enriched in one of the metabolic pathways, gluconeogenesis, which is consistent with the results of the present study [13]. Androgen secretion-stimulating factors play a pivotal role in sperm production, potentially enhancing sperm viability and count [14]. Alipour *et al.* found that pyruvate and taurine enhanced sperm viability through a correlation between seminal plasma metabolomic profile and ejaculation-abstinence period length in normozoospermic men [15]. Tauro- β -muricholate is a compound that combines taurine and bile acids, exhibiting dual functionality as both bile acids and taurine. Bile acids are associated with idiopathic infertility. Furthermore, FXR binds to endogenous bile acids and impedes spermatozoa fertilization [16]. This study indicates that serum metabolites of tauro- β -muricholate may be a risk factor for infertility in men. This may be related to the biological function of bile acids, and further investigation is required to elucidate the specific biological mechanisms. In the present study, carnitine C18:2 was identified as a potential biomarker of infertility in men, whereas carnitine C14 was found to have a protective effect against this condition. The carnitine family

encompasses a range of active forms with distinct and significantly varying functions among the various metabolite members. Urine metabolomic analyses have shown that acylcarnitines, carnitine C8, and carnitine C10:2 are strongly associated with infertility in men [17]. However, other studies have indicated that carnitine, a naturally occurring antioxidant in mammals, is a potential therapeutic agent for improving infertility in men. *In vitro* experiments have demonstrated that L-carnitine significantly improves sperm quality in patients with severe spermatopenia [18]. These findings suggest that the roles of carnitine and its metabolites in infertility in men are complex and varied and that their specific functional mechanisms require further in-depth study.

A comprehensive literature review indicated that the principal metabolic pathways implicated in infertility in men are the tricarboxylic acid cycle, oxidative phosphorylation, glycolysis, and lipid metabolism pathways [19]. Our findings indicate that the metabolic pathways associated with infertility in men, including fatty acid biosynthesis, gluconeogenesis, and the Warburg effect, are linked to the tricarboxylic acid cycle, glycolysis, and lipid metabolism. Spermocyte motility and capacitation are dependent on glycolysis. It has been proposed that the pathogenesis of infertility in men may be related to a decrease in ATP due to a blockage of the glycolytic pathway in spermatozoa [20]. Lipid metabolism plays an important role in fertility in men. The synthesis of sperm cell membranes is dependent on lipids. In men with infertility having abnormal lipid metabolism, reduced levels of phospholipids and elevated levels of free fatty acids have been observed. Mutation in the sphingomyelin synthase gene cause spermatogenesis defects in mice [21]. In summary, the pathogenesis of infertility in men is closely related to various metabolic pathways, including the tricarboxylic acid cycle, glycolysis, lipid metabolism, and fatty acid biosynthesis. Collectively, these pathways regulate spermatogenesis, motility, and energetics by affecting the energy supply, cell membrane structure, and spermatozoa function.

A growing body of evidence has indicated that infertility is a metabolic disorder, with polycystic ovary syndrome (PCOS) being the most prevalent form of infertility in women. PCOS is characterized by anovulation and endocrine and metabolic disorders [22]. A prospective randomized controlled trial of patients with PCOS identified six plasma metabolites – glutamic acid, aspartic acid, 1-methylnicotinamide, acetylcarnitine, glycerophosphocholine, and oleamide – which may serve as potential biomarkers for PCOS [23]. As identified in this study, the metabolites associated with infertility in women were predominantly lipid metabolites. Lipids play pivotal roles in human reproduction.

Blood lipids can influence women's reproductive health by regulating the production of steroid hormones and may even have adverse effects on pregnant women and fetuses [24]. The composition of lipids in the follicular fluid influences the functionality of granulosa cells, which in turn affects oocyte development and maturation [25]. An MR study by Jansen *et al.* showed that elevated triglyceride, total cholesterol, and low-density lipoprotein cholesterol levels were associated with an increased risk of infertility in women [26]. The results of our study indicated that glycerate may be a risk factor for infertility in women. Glycerates are conventionally classified into triglycerides, 1,2- or 1,3-glycerol diesters and 1- or 2-glycerol monoesters according to the number and position of the acyl groups. Of these, triglycerides represent the most significant and prevalent constituents; thus, we hypothesized that the primary cause of infertility in women because of glycerate is the action of triglycerides. Furthermore, our findings indicated a positive causal relation between bilirubin levels and infertility in women. One study examined serum total bilirubin, conjugated bilirubin, and free bilirubin levels in infertile and normal fertile women. The results indicated that all three parameters were elevated in the infertile group compared with those in the control group. The implication of these results is that elevated bilirubin levels may contribute to poor pregnancy outcomes [27]. Hood *et al.* identified nine metabolites, including bilirubin, through the metabolomic analysis of follicular fluid from 125 infertile women undergoing IVF [28]. This is consistent with our findings.

Studies of women's metabolic pathways have found that infertility in women is associated with bile acid biosynthesis. A previous study revealed that serum metabolic pathways in infertile women include fatty acid metabolism, whereas metabolic pathways in follicular fluid include bile acid biosynthesis [29], which is comparable to our findings. Smith *et al.* identified bile acids and all enzymes involved in the bile acid synthesis pathway in human follicular fluid [30]. These findings suggest that the bile acid synthesis pathway is essential for follicular growth, development, and maturation. However, further investigations are required to elucidate the underlying mechanisms of infertility. A metabolomic analysis of follicular fluid in women with evidence of infertility because of decreased ovarian reserve function revealed that the metabolic pathways involved included pantothenate and CoA biosynthesis and glutathione metabolism [29], consistent with our findings. The biosynthesis of pantothenate and CoA plays a role in the metabolism of sugars, proteins, and fats, and also exhibits antioxidant effects. These processes influence oocyte development and maturation.

This study had several strengths. It represents the most comprehensive investigation to date of the potential causal relation between plasma metabolites and infertility. Second, this study employed five distinct statistical methods for MR to enhance the robustness and credibility of the findings.

The metabolic markers and pathways identified in this study have the following clinical translational potential. The metabolites can be used as non-invasive biomarkers for the screening and stratified diagnosis of infertility. The associated metabolic pathways (such as lipid and glucose metabolism) provide new perspectives for developing targeted therapeutic strategies. Furthermore, the monitoring of metabolic processes in high-risk populations (such as patients with a family history of infertility or metabolic syndrome) in combination with lifestyle interventions has been shown to reduce the risk of infertility. However, it should be noted that the study is subject to several limitations. First, the lack of detailed participant clinical characteristics in the database limits the in-depth interpretation of the causal relation between metabolites and infertility, as different etiologies can lead to significant differences in metabolite profiles. Failure to fully consider these clinical features may have affected the generalizability and clinical application of the findings. Future studies should incorporate more detailed clinical information combined with multicenter data to further validate and optimize the reliability of metabolites as infertility biomarkers. Second, the GWAS database employed in this study predominantly comprises data from European populations. Consequently, the generalizability of these results to other racial and geographical groups remains unclear. Third, although the present study revealed significant associations between some serum metabolites and infertility using MR, whether these metabolites directly contribute to infertility or whether their changes are a result of the disease remains uncertain. Some metabolites may be involved in pathological processes by affecting sperm or egg energy metabolism, oxidative stress, or cell membrane function, whereas others may simply reflect changes in the metabolic environment. Future studies combining functional experiments (including gene editing or metabolic interventions) and longitudinal cohort studies are needed to clarify the causal mechanisms.

In conclusion, the fundamental objective of metabolomics is the identification of specific biomarkers. In the present study, MR analysis was used to identify plasma metabolite risk factors and protective factors associated with infertility. The analysis involved 17 risk factors and 10 protective factors for infertility in women and 22 risk factors and 30 protective factors for infertility in

men. Certain metabolites play a common role in infertility in both women and men. For instance, pyridoxal was positively associated with infertility in both women and men, whereas galacturonic acid and glycolic acid were identified as common risk factors. Through metabolic pathway analysis, we identified the pivotal metabolic pathways associated with infertility. These findings have yielded significant candidate molecules and established a theoretical foundation for future in-depth research on the molecular mechanisms underlying infertility and the development of targeted intervention strategies.

Acknowledgments

We gratefully acknowledge the invaluable contributions of researchers from prestigious institutions, including those who provided the blood metabolite GWAS summary data, and the FinnGen consortium. Furthermore, we express our gratitude to the diligent researchers and dedicated participants of other genome-wide association study datasets used in this research. Their steadfast commitment has been crucial for advancing scientific knowledge. We also thank BioRender.com for assistance with the illustrations.

Jiahao Shi and Jiamei Zhu contributed equality to this work.

Funding

No external funding.

Ethical approval

Not applicable.

Conflict of interest

The authors declare no conflict of interest.

References

- Vander Borcht M, Wyns C. Fertility and infertility: definition and epidemiology. *Clin Biochem* 2018; 62: 2-10.
- Mascarenhas MN, Flaxman SR, Boerma T, Vanderpoel S, Stevens GA. National, regional, and global trends in infertility prevalence since 1990: a systematic analysis of 277 health surveys. *PLOS Med* 2012; 9: e1001356.
- Service CA, Puri D, Al Azzawi S, Hsieh TC, Patel DP. The impact of obesity and metabolic health on male fertility: a systematic review. *Fertil Steril* 2023; 120: 1098-111.
- Dai M, Hong L, Yin T, Liu S. Disturbed follicular microenvironment in polycystic ovary syndrome: relationship to oocyte quality and infertility. *Endocrinology* 2024; 165: bqae023.
- Kovac JR, Pastuszak AW, Lamb DJ. The use of genomics, proteomics, and metabolomics in identifying biomarkers of male infertility. *Fertil Steril* 2013; 99: 998-1007.
- Zhang A, Sun H, Wang P, Han Y, Wang X. Recent and potential developments of biofluid analyses in metabolomics. *J Proteomics* 2012; 75: 1079-88.
- Sekula P, Del Greco MF, Pattaro C, Köttgen A. Mendelian randomization as an approach to assess causality using observational data. *J Am Soc Nephrol* 2016; 27: 3253-65.
- Chen Y, Lu T, Pettersson-Kymmer U, et al. Genomic atlas of the plasma metabolome prioritizes metabolites implicated in human diseases. *Nat Genet* 2023; 55: 44-53.
- Davies NM, Holmes MV, Davey Smith G. Reading Mendelian randomisation studies: a guide, glossary, and checklist for clinicians. *BMJ* 2018; 362: k601.
- Burgess S, Davey Smith G, Davies NM, et al. Guidelines for performing Mendelian randomization investigations: update for summer 2023. *Wellcome Open Res* 2019; 4: 186.
- Jørgensen N, Joensen UN, Jensen TK, et al. Human semen quality in the new millennium: a prospective cross-sectional population-based study of 4867 men. *BMJ Open* 2012; 2: e000990.
- Courant F, Antignac JP, Monteau F, Le Bizec B. Metabolomics as a potential new approach for investigating human reproductive disorders. *J Proteome Res* 2013; 12: 2914-20.
- Zhang Z, Zhang Y, Liu C, et al. Serum metabolomic profiling identifies characterization of non-obstructive azoospermic men. *Int J Mol Sci* 2017; 18: 238.
- Zhang J, Huang Z, Chen M, et al. Urinary metabolome identifies signatures of oligozoospermic infertile men. *Fertil Steril* 2014; 102: 44-53.
- Alipour H, Duus RK, Wimmer R, et al. Seminal plasma metabolomics profiles following long (4-7 days) and short (2 h) sexual abstinence periods. *Eur J Obstet Gynecol Reprod Biol* 2021; 264: 178-83.
- Yang J, Zong X, Wu G, Lin S, al Feng Y, Hu J. Taurine increases testicular function in aged rats by inhibiting oxidative stress and apoptosis. *Amino Acids* 2015; 47: 1549-58.
- Malivindi R, Santoro M, De Rose D, et al. Activated-farnesoid X receptor (FXR) expressed in human sperm alters its fertilising ability. *Reproduction* 2018; 156: 249-59.
- Chang D, Li F, Kang Y, et al. The effects of L-carnitine and fructose in improved Ham's F10 on sperm culture in idiopathic severe asthenospermia within 24h. *PLoS One* 2025; 20: e0306235.
- Peña FJ, Ortiz-Rodríguez JM, Gaitskell-Phillips GL, Gil MC, Ortega-Ferrusola C, Martín-Cano FE. An integrated overview on the regulation of sperm metabolism (glycolysis-Krebs cycle-oxidative phosphorylation). *Anim Reprod Sci* 2022; 246: 106805.
- Eirefelt S, Stahlhut M, Svitacheva N, et al. Characterization of a novel non-steroidal glucocorticoid receptor agonist optimized for topical treatment. *Sci Rep* 2022; 12: 1501.
- Wittmann A, Grimm MOW, Scherthan H, et al. Sphingomyelin synthase 1 is essential for male fertility in mice. *PLoS One* 2016; 11: e0164298.
- Azziz R, Carmina E, Chen Z, et al. Polycystic ovary syndrome. *Nat Rev Dis Primers* 2016; 2: 16057.
- Ding X, Deng Y, Wang Y, et al. Serum metabolomic profiling reveals potential biomarkers in assessing the management of women with polycystic ovary syndrome: a randomized controlled trial. *Chin Med J (Engl)* 2022; 135: 79-85.
- Morel Y, Roucher F, Ploton I, Goursaud C, Tardy V, Mallet D. Evolution of steroids during pregnancy: maternal, placental and fetal synthesis. *Ann Endocrinol (Paris)* 2016; 77: 82-9.

25. Liu Y, Zhai J, Chen J, Wang X, Wen T. PGC-1 α protects against oxidized low-density lipoprotein and luteinizing hormone-induced granulosa cells injury through ROS-p38 pathway. *Hum Cell* 2019; 32: 285-96.
26. Jansen H, Lieb W, Schunkert H. Mendelian randomization for the identification of causal pathways in atherosclerotic vascular disease. *Cardiovasc Drugs Ther* 2016; 30: 41-9.
27. Mangione R, Pallisco R, Bilotta G, et al. Bilirubin concentration in follicular fluid is increased in infertile females, correlates with decreased antioxidant levels and increased nitric oxide metabolites, and negatively affects outcome measures of in vitro fertilization. *Int J Mol Sci* 2023; 24: 10707.
28. Hood RB, Liang D, Tan Y, et al. Serum and follicular fluid metabolome and markers of ovarian stimulation. *Hum Reprod* 2023; 38: 2196-207.
29. Li J, Zhang Z, Wei Y, Zhu P, al Yin T, Wan Q. Metabonomic analysis of follicular fluid in patients with diminished ovarian reserve. *Front Endocrinol (Lausanne)* 2023; 14: 1132621.
30. Smith LP, Nierstenhoefer M, Yoo SW, Penzias AS, Tobiasch E, Usheva A. The bile acid synthesis pathway is present and functional in the human ovary. *PLoS One* 2009; 4: e7333.