Associations of preoperative Oswestry Disability Index and EuroQoI-5D with long-term all-cause mortality in patients undergoing percutaneous vertebroplasty

Keywords

vertebroplasty, EQ-5D, ODI, mortality, fracture

Abstract

Introduction

Oswestry Disability Index (ODI) and EuroQoI-5D (EQ-5D) have been widely used to assess general health quality and function in clinical studies of patients with vertebral fractures. We aimed to investigate the associations of preoperative ODI and EQ-5D with long-term mortality in patients undergoing percutaneous vertebroplasty.

Material and methods

We retrospectively identified adult patients who had a single-level vertebral compression fracture and received percutaneous vertebroplasty between 2013 and 2020. Patients with traumatic fractures, burst fractures, and pathologic fractures, as well as those who had missing information on preoperative assessment of ODI and EQ-5D, were excluded. Survival status of the study patients was confirmed at the end of March 2021. The associations of preoperative ODI and EQ-5D with all-cause mortality were examined using Cox-proportional hazard models.

Results

A total of 167 patients were analyzed (mean age 75.8 \pm 9.3 years, 25.7% male). There were 28 patients who died during a median follow-up duration of 2.1 years (63.6 per 1000 patient-years). Preoperative ODI was significantly associated with all-cause mortality after vertebroplasty (HR 1.049, 95% CI 1.008 to 1.092, p=0.018). In contrast, preoperative EQ-5D was independently associated with a lower risk of all-cause mortality after the surgery (HR 0.202, 95% CI 0.043 to 0.936, p=0.041).

Conclusions

Preoperative assessment of ODI (HR 1.049, 95% CI 1.008 to 1.092) and EQ-5D (HR 0.202, 95% CI 0.043 to 0.936) may help determine postoperative long-term mortality risk in this aging surgical population.

Introduction

Vertebral fractures are the most common fracture associated with low bone mass and skeletal fragility [1-2]. Variance in clinical presentations of vertebral fractures may contribute partly to the wide variations of their prevalence and incidence in previous studies [3-4], with the highest rates in North America and Asia [5]. They commonly occurred in the elderly [6], and could cause substantial pain, disability, adverse health outcomes, and even mortality [7]. The impacts of sex and gender on orthopedic health are also important issues deserve further investigations, such as the higher rates of done deformities in females and osteoporosis in postmenopausal women [8]. Since populations are rapidly aging worldwide [9], vertebral fractures have given rise to a heavy healthcare burden and constitute a serious public health issue [1].

For patients suffering from a vertebral fracture, percutaneous vertebroplasty has been developed to relieve pain and improve their physical function and quality of life [10]. Although vertebroplasty has become a common surgical procedure, its effects on symptoms relief, quality of life, and long-term patient outcomes remain unclear [11-13]. Several patient-reported outcome measures are recommended for assessment of general health quality and function in patients who undergo spine surgery [14]. Among these, the Oswestry Disability Index (ODI) [15] and EuroQol-5D (EQ-5D) [16] have been widely used in clinical studies of patients with vertebral fractures [17-21]. However, whether patient-reported outcome measures were associated with long-term mortality in patients who received vertebroplasty is not yet clear. In this study, we aimed to investigate the associations of preoperative ODI and EQ-5D with long-term mortality in patients with vertebral compression fractures who underwent percutaneous vertebroplasty.

Materials and Methods

This retrospective study was conducted in a tertiary medical center. The study protocol was approved by the Institutional Review Board of Taichung Veterans General Hospital, Taichung, Taiwan (approval number CE22167A). Patient informed consent was waived due to the retrospective study design. We identified adult patients who had a single-level vertebral compression fracture and received percutaneous vertebroplasty between 2013 and 2020. In previous studies [22-23] that examined quality of life for osteoporosis-related conditions (including hip fracture, vertebral fracture, and distal forearm fracture), the follow-up duration was generally 18 months or less. In another study [24] that reported postoperative outcomes including all-cause mortality in patients undergoing thoracolumbar fusion for spinal deformity, 71% of patients were followed up for at least 10 months. Compared with these studies, our study period is sufficient to observe the effects of percutaneous vertebroplasty on patients' outcomes.

In our patients, surgical intervention was conducted after preoperative assessment including ODI [15] and EQ-5D [16]. Patients with traumatic fractures, burst fractures, and pathologic fractures according to discharge diagnosis, as well as those who had missing information on preoperative assessment of ODI and EQ-5D, were excluded. Pathologic fracture was defined as fractures that occur without an adequate traumatic event due to focal benign or malignant skeletal lesions [25]. For optimal preoperative care in an orthopedic department, nursing leadership and team work are important during the COVID-19 pandemic and post-COVID-19 era [26]. Assessment of ODI [15] and EQ-5D [16] was conducted by a trained-nurse with more than 8 years of experience as part of the preoperative evaluation [14] in our hospital. The assessment of ODI [15] contains ten questions (pain intensity, personal care, lifting, walking, sitting, standing, sleeping, sex life if applicable, social life, and traveling) on limitations of activities of daily living. Each item is rated on a 0-5 point scale, and a higher ODI score indicates greater severity of disability. The EQ-5D [16] consists of an assessment of five dimensions (mobility, self-care, usual activities, pain/discomfort, and anxiety/depression), and a higher value indicates a better quality of life. The reliability

of ODI and EQ-5D in patients undergoing spinal surgery have been assessed in previous studies [27-28]. The validity of ODI and EQ-5D to assess the effectiveness of lumbar fusion for degenerative lumbar spondylolisthesis was evaluated in a prospective cohort study [29]. The area under the ROC curve against patient-reported outcomes was 0.94 for ODI and 0.97 for EQ-5D. The specificity and sensitivity for ODI were ~85% and ~95%, respectively. The respective values for EQ-5D were ~100% and ~90% [29]. Baseline demographics, including age (range: 48 to 94 years), sex, and body mass index, and medical history, were collected from electronic medical records. Survival status of the study patients at the end of March 2021 was obtained from the Ministry of Health and Welfare, R.O.C. De-identified data were then used for analyses.

A statistician (CH Lin) conducted all statistical analyses independently using deidentified data. Categorical and continuous variables are presented as number (percentage) and mean ± SD, respectively. The outcome of interest was all-cause mortality. The independent variables examined were preoperative ODI and EQ-5D. The associations of preoperative ODI and EQ-5D with all-cause mortality were examined using the Cox-proportional hazards model which is a frequently used approach to investigate relationships between the time to event outcome and a set of explanatory variables. We considered the following variables as potential confounders, including age, sex, body mass index, smoking, diabetes, hypertension, chronic kidney disease, osteoporosis, and medical treatment for osteoporosis. These variables were adjusted in the Cox proportional hazards models to examine the independent associations of ODI and EQ-5D with all-cause mortality. A Cox proportional hazards model with restricted cubic splines [30] was used to display the relation between independent variables (preoperative ODI and EQ-5D) and all-cause mortality. All of the statistical analyses were performed using the Statistical Package for the Social Sciences (IBM SPSS version 22.0; International Business Machines Corp, NY, USA). A two-sided p-value of less than 0.05 was considered statistically significant.

Results

We identified 209 adult patients who had a single-level vertebral compression fracture and received percutaneous vertebroplasty. After excluding 18 patients with traumatic fractures (n=10), burst fractures (n=2), or pathologic fractures (n=6), and 24 patients who had missing information on preoperative ODI or EQ-5D, a total of 167 patients were analyzed (Table I). The mean age of the study sample was 75.8 ± 9.3 years (range from 48 to 94 years), and 25.7% of them were male. The prevalence rates of diabetes, hypertension, and chronic kidney disease were 19.8%, 53.3%, and 34.1%, respectively. Most of the patients had osteoporosis (74.3%) and received medical treatment (63.5%). The levels of vertebroplasty were T- (42.5%) or L-spine (57.5%). The distribution of operation level of T-spine vertebroplasty were T6 (n=2), T7 (n=3), T8 (n=3), T9 (n=3), T10 (n=6), T11 (n=15), and T12 (n=39). For L-spine, the distribution were L1 (n=41), L2 (n=29), L3 (n=13), L4 (n=9), and L5 (n=4). The mean pre-operative ODI and EQ-5D were 71.3 ± 12.4 and 0.25 ± 0.22 respectively. There were 28 patients who died during a median follow-up duration of 2.1 years (63.6 per 1000 patient-years). Among the 28 patients died during the follow-up period, 12 of them (42.9%) had T-spine fracture. None of the 28 patients died during the index hospitalization, while four of them died within 150 days after the surgery (range: 75-119 days). Hence, we suggested that the cause of death was not likely related to vertebroplasty. There was no significant difference regarding the distribution of the levels of spine fracture compared to those who were alive (T-spine facture 42.4% [59/139], p=0.968) by the end of study period.

The associations of preoperative ODI and EQ-5D with all-cause mortality were examined using Cox-proportional hazard models with multivariate adjustment (age, sex, body mass index, smoking, and concomitant chronic diseases). As shown in Table II, preoperative ODI was significantly associated with all-cause mortality after vertebroplasty (HR 1.060, 95% CI 1.023 to 1.098, p=0.001). The association remained significant (HR 1.049, 95% CI 1.008 to 1.092, p=0.018) after multivariate adjustment. Most previous studies examined epidemiology of vertebral fractures in aged people (for example, \geq 65 years) [1]. We examined findings of the multivariate adjusted models in our patients aged \geq 65 years (n=146), and similar findings were noted (HR 1.054, 95% CI 1.010 to 1.101, p=0.016). In contrast, preoperative EQ-5D was associated with a lower risk of all-cause mortality after the surgery (HR 0.102, 95% CI 0.025 to 0.420, p=0.002). The association was independent of age, sex, body mass index, and other comorbidities (HR 0.202, 95% CI 0.043 to 0.936, p=0.041). The findings were consistent in patients with age \geq 65 years (HR 0.168, 95% CI 0.035 to 0.808, p=0.026).

Cubic spline of preoperative ODI and EQ-5D versus risk of all-cause mortality by a Cox-proportional hazards model was conducted [30]. The Scaled Schoenfeld residuals were used to test the assumption of Cox-proportional hazards model, and no violation of the assumption was confirmed. Figure 1 shows the cubic spline of preoperative ODI versus risk of all-cause mortality in the study sample. An increase in all-cause mortality risk was observed in patients who had a preoperative ODI higher than 70. Figure 2 shows the cubic spline of preoperative EQ-5D versus risk of all-cause mortality. We observed an increase in all-cause mortality risk in those who had a preoperative EQ-5D lower than 0.2.

Discussion

In this study, we demonstrated that preoperative patient-reported outcome measures (ODI and EQ-5D) were associated with long-term all-cause mortality in patients who underwent a vertebroplasty for a single-level vertebral compression fracture. Although patient-reported outcome measures have been proposed for assessment of general health quality and function in those who received spine surgery [14], their associations with long-term outcomes are not yet clear. To the best of our knowledge, our study is the first to investigate the associations between preoperative patient-reported outcome measures (ODI and EQ-5D) and long-term mortality in patients who underwent a vertebroplasty for a single-level vertebral compression fracture.

Patients suffering from a vertebral compression fracture are typically aged ≥ 65 years [11-13], and age is a predictor of vertebral fractures [1]. Moreover, vertebral fractures have been associated with progression of kyphosis, which in turn is associated with reduced pulmonary function [31] and physical function [32], and possibly falls [33]. All of these effects may lead to impairment of quality of life and a higher risk of

mortality [1, 34-35]. A vertebroplasty may relieve symptoms and improve functional outcomes in these patients [13]; nevertheless, its effect on long-term mortality is not yet clear [11-12, 36]. Percutaneous vertebroplasty relieved pain and improved quality of life in patients (mean age 73.3 ± 7.9 years) with osteoporotic compression fractures in a prospective study [10]. These findings were similar to the results of an open-label prospective randomized trial [13], although there have been inconsistent studies [11-12]. These studies varied in selected populations, assessment of outcomes, and duration of follow-up. In the two randomized trials [11-12] which reported no beneficial effect of vertebroplasty compared with the control group, the authors included patients with a duration of pain of up to 12 months, and assessed their primary outcome (pain scores) at 1 and 3 months, respectively. In the other randomized study [13], the authors included patients with uncontrolled pain for less than 6 weeks, and the primary outcome (pain score) was assessed at 1 month and 1 year. Some other issues that may contribute to the variability in findings across different studies have been discussed [37].

Patients with compression fractures are at a higher risk of mortality. In a cohort study [38] of patients older than 60 years, the risk of long-term mortality doubled in those who had osteoporotic vertebral fractures. The increase in mortality risk after vertebral fractures might be partly attributed to functional disability, physical inactivity,

and impairment of quality of life [31-33], all of which have been associated with mortality risk in elderly populations [39-43]. Nevertheless, the effects of surgical intervention on these measures and mortality outcome in patients with vertebral fractures are not yet clear. Findings from previous reports [36, 44-45] on the risk of mortality after surgical interventions for vertebral fractures are inconsistent. In a large retrospective cohort study of more than 850000 patients with vertebral compression fractures and up to 4 years of follow-up [44], operated cohort (kyphoplasty or vertebroplasty) had a significantly lower risk of mortality (adjusted relative risk 0.63, 95% CI 0.62 to 0.64, p<0.001) compared with nonoperated cohort. In contrast, surgical treatment (kyphoplasty or vertebroplasty) for osteoporotic vertebral fractures had no significant difference in 1-year mortality risk (hazard ratio 0.92, 95% CI 0.81 to 1.04, p=0.18) compared with propensity score matched control group [36]. In another retrospective cohort study [45], vertebroplasy for vertebral compression fractures had a non-significantly higher risk of mortality after multivariate adjustment (adjusted hazard ratio 1.17, 95% CI 0.96 to 1.42), compared with control group after a mean follow-up period of nearly 4 years. All these reports are retrospective cohort studies, and there may be confounding factors that cannot not be addressed due to the retrospective study design. Moreover, patient-reported outcome measures (such as ODI and EQ-5D) were

not included in these analyses. In our study, we investigated the association between preoperative assessment of ODI and EQ-5D with postoperative long-term mortality in patients undergoing vertebroplasty. Preoperative assessment of patient-reported outcome measures [14] may represent baseline condition of disability and quality of life and help predict long-term outcomes in this aging population. Based on our findings, a higher preoperative ODI and a lower preoperative EQ-5D were independently associated with a higher risk of mortality in patients undergoing vertebroplasty for a single-level vertebral compression fracture during a median follow-up period of 2.1 years. Our results are in line with a recent retrospective study [46] which revealed that preoperative EQ-5D was negatively associated with postoperative mortality during a mean follow-up period of 7.5 years in patients undergoing total hip arthroplasty and knee arthroplasty. Taken together, we suggest that preoperative assessment of ODI and EQ-5D, both of which are recommended patient-reported outcome measures in spine surgery [14], may help determine postoperative long-term mortality risk in patients undergoing vertebroplasty for vertebral compression fractures.

The strength of this study is that the ODI and EQ-5D were assessed by a trained nurse at baseline. The reliability and validity of ODI and EQ-5D in patients undergoing spinal surgery have been reported [27-29]. Moreover, the status of survival was examined after a median follow-up duration of more than 2 years. In contrast, the

follow-up period of most previous studies was usually less than 18 months [22-24].

However, there were several limitations in this study. First, the retrospective study design might have led to patient selection bias. Moreover, the size of the study sample was relatively small. Second, although we had adjusted for some concomitant chronic diseases (such as diabetes and hypertension) in the analyses to determine the associations of baseline ODI and EQ-5D with all-cause mortality, we did not assess disease control status (such as blood glucose and blood pressure levels), which might have confounded our results. Third, the use of complex and lengthy questionnaires (ODI and EQ-5D) may have posed challenges in terms of patient comprehension and completion, possibly leading to incomplete or inaccurate data collection. Despite these limitations, we report for the first time that preoperative assessment of ODI and EQ-5D was associated with long-term all-cause mortality in patients undergoing percutaneous vertebroplasty. Our findings need to be confirmed in future studies with a larger sample size.

In summary, a higher preoperative ODI and a lower preoperative EQ-5D were independently associated with a higher risk of long-term all-cause mortality in patients undergoing vertebroplasty for a single-level vertebral compression fracture. Preoperative assessment of ODI and EQ-5D may help determine postoperative longterm mortality risk in this aging population. We suggest that careful consideration is required before performing vertebroplasty in elderly patients with a high ODI or a low EQ-5D.

Acknowledgements: None.

Declaration of Competing Interest: The authors declare that they have no competing interests.

Funding: This work was supported by research grants from Taichung Veterans General Hospital, Taichung, Taiwan [grant numbers TCVGH-1113502C, 2022; TCVGH-1123502C, 2023]. The funder was not involved in the study design, data collection, analysis, interpretation of the results, or preparation of the article.

References

[1] Schousboe JT. Epidemiology of Vertebral Fractures. J Clin Densitom. 2016; 19(1):8-22.

[2] Compston JE, McClung MR, Leslie WD. Osteoporosis. Lancet. 2019; 393: 364-376.

[3] Delmas PD, van de Langerijt L, Watts NB, et al; IMPACT Study Group.

Underdiagnosis of vertebral fractures is a worldwide problem: the IMPACT study. J

Bone Miner Res. 2005; 20(4): 557-563.

[4] Majumdar SR, Kim N, Colman I, et al. Incidental vertebral fractures discovered with chest radiography in the emergency department: prevalence, recognition, and osteoporosis management in a cohort of elderly patients. Arch Intern Med. 2005; 165(8): 905-909.

[5] Ballane G, Cauley JA, Luckey MM, El-Hajj Fuleihan G. Worldwide prevalence and incidence of osteoporotic vertebral fractures. Osteoporos Int. 2017; 28(5): 1531-1542.
[6] Cosman F, Krege JH, Looker AC, et al. Spine fracture prevalence in a nationally representative sample of US women and men aged ≥40 years: results from the National Health and Nutrition Examination Survey (NHANES) 2013-2014. Osteoporos Int. 2017; 28(6): 1857-1866.

[7] Ensrud KE, Thompson DE, Cauley JA, et al. Prevalent vertebral deformities predict

mortality and hospitalization in older women with low bone mass. Fracture Intervention Trial Research Group. J Am Geriatr Soc. 2000; 48(3): 241-249.

[8] Biz C, Khamisy-Farah R, Puce L, et al. Investigating and Practicing Orthopedics at the Intersection of Sex and Gender: Understanding the Physiological Basis, Pathology, and Treatment Response of Orthopedic Conditions by Adopting a Gender Lens: A Narrative Overview. Biomedicines. 2024; 12(5): 974.

[9] Chatterji S, Byles J, Cutler D, Seeman T, Verdes E. Health, functioning, and disability in older adults--present status and future implications. Lancet. 2015; 385(9967): 563-575.

[10] Alvarez L, Alcaraz M, Pérez-Higueras A, et al. Percutaneous vertebroplasty:functional improvement in patients with osteoporotic compression fractures. Spine(Phila Pa 1976). 2006; 31(10): 1113-1118.

[11] Kallmes DF, Comstock BA, Heagerty PJ, et al. A randomized trial of vertebroplasty for osteoporotic spinal fractures. N Engl J Med. 2009; 361(6): 569-579.

[12] Buchbinder R, Osborne RH, Ebeling PR, et al. A randomized trial of vertebroplasty for painful osteoporotic vertebral fractures. N Engl J Med. 2009; 361(6): 557-568.

[13] Klazen CA, Lohle PN, de Vries J, et al. Vertebroplasty versus conservative

treatment in acute osteoporotic vertebral compression fractures (Vertos II): an open-

label randomised trial. Lancet. 2010; 376(9746): 1085-1092.

[14] McCormick JD, Werner BC, Shimer AL. Patient-reported outcome measures in spine surgery. J Am Acad Orthop Surg. 2013; 21(2): 99-107.

[15] Fairbank JC, Pynsent PB. The Oswestry Disability Index. Spine (Phila Pa 1976).2000; 25(22): 2940-2952; discussion 2952.

[16] Tsuchiya A, Ikeda S, Ikegami N, et al. Estimating an EQ-5D population value set: the case of Japan. Health Econ. 2002; 11(4): 341-353.

[17] Lee HM, Park SY, Lee SH, Suh SW, Hong JY. Comparative analysis of clinical outcomes in patients with osteoporotic vertebral compression fractures (OVCFs): conservative treatment versus balloon kyphoplasty. Spine J. 2012; 12(11): 998-1005.
[18] Cheng X, Long HQ, Xu JH, Huang YL, Li FB. Comparison of unilateral versus bilateral percutaneous kyphoplasty for the treatment of patients with osteoporosis vertebral compression fracture (OVCF): a systematic review and meta-analysis. Eur Spine J. 2016; 25(11): 3439-3449.

[19] Verlaan JJ, Somers I, Dhert WJ, Oner FC. Clinical and radiological results 6 years after treatment of traumatic thoracolumbar burst fractures with pedicle screw instrumentation and balloon assisted endplate reduction. Spine J. 2015; 15(6): 1172-1178. [20] Diel P, Reuss W, Aghayev E, Moulin P, Röder C; SWISSspine Registry Group.SWISSspine-a nationwide health technology assessment registry for balloonkyphoplasty: methodology and first results. Spine J. 2010; 10(11): 961-971.

[21] Hübschle L, Borgström F, Olafsson G, et al; SWISSspine Registry Group. Real-life results of balloon kyphoplasty for vertebral compression fractures from the SWISSspine registry. Spine J. 2014; 14(9): 2063-2077.

[22] Si L, Winzenberg TM, de Graaff B, Palmer AJ. A systematic review and metaanalysis of utility-based quality of life for osteoporosis-related conditions. Osteoporos Int. 2014; 25(8): 1987-1997.

[23] Svedbom A, Borgstöm F, Hernlund E, et al. Quality of life for up to 18 months after low-energy hip, vertebral, and distal forearm fractures-results from the ICUROS.Osteoporos Int. 2018; 29(3): 557-566.

[24] Akbik OS, Al-Adli N, Pernik MN, et al. A Comparative Analysis of Frailty,Disability, and Sarcopenia With Patient Characteristics and Outcomes in Adult SpinalDeformity Surgery. Global Spine J. 2023; 13(8): 2345-2356.

[25] Wuennemann F, Kintzelé L, Weber MA, Kauczor HU, Rehnitz C. RadiologischeDiagnostik pathologischer Frakturen [Radiologic diagnosis of pathologic fractures].Radiologe. 2020; 60(6): 498-505.

[26] Biz C, Buffon L, Scapinello D, et al. Nursing Leadership in a Post-Pandemic
Elective Orthopaedic Theatre Department: A Detailed Thematic Analysis of an OpenEnded Qualitative Survey. Nurs Rep. 2024; 14(3): 1541-1552.

[27] Solberg TK, Olsen JA, Ingebrigtsen T, Hofoss D, Nygaard OP. Health-related quality of life assessment by the EuroQol-5D can provide cost-utility data in the field of low-back surgery. Eur Spine J. 2005; 14(10): 1000-1007.

[28] Stokes OM, Cole AA, Breakwell LM, Lloyd AJ, Leonard CM, Grevitt M. Do we have the right PROMs for measuring outcomes in lumbar spinal surgery? Eur Spine J. 2017; 26(3): 816-824.

[29] Godil SS, Parker SL, Zuckerman SL, Mendenhall SK, Glassman SD, McGirt MJ. Accurately measuring the quality and effectiveness of lumbar surgery in registry efforts: determining the most valid and responsive instruments. Spine J. 2014; 14(12): 2885-2891.

[30] Austin PC. Graphical methods to illustrate the nature of the relation between a continuous variable and the outcome when using restricted cubic splines with a Cox proportional hazards model. Stat Methods Med Res. 2024: 9622802241287707.

[31] Di Bari M, Chiarlone M, Matteuzzi D, et al. Thoracic kyphosis and ventilatory dysfunction in unselected older persons: an epidemiological study in Dicomano, Italy. J Am Geriatr Soc. 2004; 52(6): 909-915.

[32] Katzman WB, Huang MH, Lane NE, Ensrud KE, Kado DM. Kyphosis and decline in physical function over 15 years in older community-dwelling women: the Study of Osteoporotic Fractures. J Gerontol A Biol Sci Med Sci. 2013; 68(8): 976-983.

[33] Kado DM, Prenovost K, Crandall C. Narrative review: hyperkyphosis in older persons. Ann Intern Med. 2007; 147(5): 330-338.

[34] Kendler DL, Bauer DC, Davison KS, et al. Vertebral Fractures: Clinical Importance and Management. Am J Med. 2016; 129(2): 221.e1-10.

[35] Morin S, Lix LM, Azimaee M, Metge C, Caetano P, Leslie WD. Mortality rates after incident non-traumatic fractures in older men and women. Osteoporos Int. 2011; 22(9): 2439-2448.

[36] McCullough BJ, Comstock BA, Deyo RA, Kreuter W, Jarvik JG. Major medical outcomes with spinal augmentation vs conservative therapy. JAMA Intern Med. 2013; 173(16): 1514-1521.

[37] Clark W, Lyon S, Burnes J. Trials of vertebroplasty for vertebral fractures. N Engl J Med. 2009; 361(21): 2097-2098; author reply 2099-2100.

[38] Bliuc D, Nguyen ND, Milch VE, Nguyen TV, Eisman JA, Center JR. Mortality risk associated with low-trauma osteoporotic fracture and subsequent fracture in men and

women. JAMA. 2009; 301(5): 513-521.

[39] Son KY, Shin DW, Lee JE, Kim SH, Yun JM, Cho B. Association of timed up and go test outcomes with future incidence of cardiovascular disease and mortality in adults aged 66 years: Korean national representative longitudinal study over 5.7 years. BMC Geriatr. 2020; 20(1): 111.

[40] Cheong CY, Yap P, Gwee X, et al. Physical and functional measures predicting long-term mortality in community-dwelling older adults: a comparative evaluation in the Singapore Longitudinal Ageing Study. Aging (Albany NY). 2021; 13(23): 25038-25054.

[41] Ascencio EJ, Cieza-Gómez GD, Carrillo-Larco RM, Ortiz PJ. Timed up and go test predicts mortality in older adults in Peru: a population-based cohort study. BMC Geriatr. 2022; 22(1): 61.

[42] Phyo AZZ, Ryan J, Gonzalez-Chica DA, et al; ASPREE Investigator Group.
Health-related quality of life and all-cause mortality among older healthy individuals in
Australia and the United States: a prospective cohort study. Qual Life Res. 2021; 30(4):
1037-1048.

[43] Gobbens RJJ, van der Ploeg T. The prediction of mortality by quality of life assessed with the WHOQOL-BREF: a longitudinal analysis at the domain and item levels using a seven-year follow-up period. Qual Life Res. 2021; 30(7): 1951-1962.

[44] Edidin AA, Ong KL, Lau E, Kurtz SM. Mortality risk for operated and nonoperated vertebral fracture patients in the medicare population. J Bone Miner Res.2011; 26(7): 1617-1626.

[45] McDonald RJ, Achenbach SJ, Atkinson EJ, et al. Mortality in the vertebroplasty population. AJNR Am J Neuroradiol. 2011; 32(10): 1818-1823.

[46] Clement ND, Patton RFL, MacDonald DJ, Duckworth AD. Preoperative health-related quality of life is independently associated with postoperative mortality riskfollowing total hip or knee arthroplasty: seven to eight years' follow-up. Bone Jt Open.2022; 3(12): 933-940.

Figure legends:

Figure 1. Cubic spline of preoperative Oswestry Disability Index versus risk of all-cause mortality.

Figure 2. Cubic spline of preoperative EuroQol-5D versus risk of all-cause mortality.



Number of patients	167
Age, years	75.8 ± 9.3
Male sex, n (%)	43 (25.7)
Body mass index, kg/m ²	24.0 ± 4.1
Smoking, n (%)	12 (7.2)
Diabetes, n (%)	33 (19.8)
Hypertension, n (%)	89 (53.3)
Chronic kidney disease, n (%)	57 (34.1)
Osteoporosis, n (%)	124 (74.3)
Medication for osteoporosis, n (%) ^a	106 (63.5)
Level of vertebroplasty, n (%)	
T-spine	71 (42.5)
L-spine	96 (57.5)
Oswestry Disability Index	71.3 ± 12.4
EuroQol-5D	0.25 ± 0.22

Values are mean \pm SD or n (%). ^aBisphosphonate, receptor activator of nuclear factor kappa-B inhibitor, or parathyroid hormone.

	Hazard Ratio (95% CI)	Р
Oswestry Disability Index		
Model 1	1.060 (1.023, 1.098)	0.001
Model 2	1.061 (1.022, 1.102)	0.002
Model 3	1.049 (1.008, 1.092)	0.018
EuroQol-5D		
Model 1	0.102 (0.025, 0.420)	0.002
Model 2	0.106 (0.025, 0.442)	0.002
Model 3	0.202 (0.043, 0.936)	0.041

Table II – Associations of preoperative ODI and EQ-5D with all-cause mortality

EQ-5D, EuroQol-5D. ODI, Oswestry Disability Index. Model 1, unadjusted. Model 2, adjusted for age and sex. Model 3, adjusted for variables in Model 2 plus body mass index, smoking, diabetes, hypertension, chronic kidney disease, osteoporosis, and medical treatment for osteoporosis.



Figure 1. Cubic spline of preoperative Oswestry Disability Index versus risk of all-cause mortality.



Figure 2. Cubic spline of preoperative EuroQoI-5D versus risk of all-cause mortality.