

# Relationship between body weight and sexual functioning evaluation index

Maciej Białorudzki<sup>1,2\*</sup>, Joanna Mazur<sup>2</sup>, Zbigniew Izdebski<sup>2,3</sup>

<sup>1</sup>Faculty of Education, University of Warsaw, Poland

<sup>2</sup>Department of Humanization of Health Care and Sexology, Collegium Medicum, University of Zielona Gora, Poland

<sup>3</sup>Department of Biomedical Aspects of Development and Sexology, Faculty of Education, University of Warsaw, Poland

Submitted: 16 April 2025; Accepted: 10 June 2025

Online publication: 19 August 2025

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms/206973>

Copyright © 2025 Termedia & Banach

\*Corresponding author:

Maciej Białorudzki

Department of Humanization

of Health Care

and Sexology

Collegium Medicum

University of Zielona Gora

Poland

E-mail: [mbialorudzki@uz.zgora.pl](mailto:mbialorudzki@uz.zgora.pl)

## Abstract

**Introduction:** This study investigated the association of body weight, self-rated health, and physical attractiveness with sexual life evaluation in Polish healthcare workers.

**Material and methods:** A cross-sectional survey was conducted between February and April 2022 among 1,478 healthcare workers from 99 Polish hospitals and specialized clinics. A total of 27.5% of the respondents were physicians. Data were collected using an online and paper-based questionnaire assessing body mass index (BMI), self-rated health, physical attractiveness, stress, sleep, and sexual life evaluation. Statistical analysis included hierarchical linear regression and structural equation modeling (SEM) to identify direct and indirect relationships.

**Results:** The data indicate that 54.7% of subjects had a BMI indicating excessive weight. Of these, 17.8% had a BMI that indicated obesity. BMI, health assessment, and physical attractiveness significantly influenced sexual life evaluation. Regression models demonstrated that impaired body image and health perception were the strongest predictors of lower sexual life evaluation ( $R^2 = 0.365$ ,  $p < 0.001$ ). SEM analysis revealed BMI's indirect impact on sexual functioning, primarily mediated through attractiveness and health rating ( $-0.345$ ,  $-0.238$ ;  $p < 0.001$ ).

**Conclusions:** Our findings highlight the complex links between body weight, self-perception, and sexual functioning. Self-rated health and physical attractiveness emerged as critical mediators, underscoring the need for targeted interventions addressing body image and health perceptions to improve sexual well-being in populations at risk of excess body weight.

**Key words:** sexual life, assessment of sexual function, body mass index, self-rated health, self-rated physical attractiveness, healthcare workers.

## Introduction

Excess body weight poses significant public health challenges worldwide, with rising prevalence and serious health consequences [1, 2]. Problems associated with excess body weight impose a substantial burden on healthcare systems and lead to high social and economic costs [3]. According to the World Health Organization (WHO), being overweight is defined as having excessive fat deposits, while obesity is classified as a chronic, complex disease characterized by excessive fat deposits that

can impair health [4]. The negative effects of excess body weight on health-related quality of life are well documented [5, 6]. Along with its high prevalence, excess body weight has been linked to poorer physical, mental, and sexual health [7].

The approach to sexuality focuses mainly on medical aspects, often overlooking other equally important dimensions, such as subjective evaluations of sexual life, overall satisfaction levels, and perceptions of sexual performance [8, 9]. Sexual functioning can be described as the engagement of motivators such as sexual attraction and desire, while also considering the significance of health, psychosocial factors, and body image self-assessment [10, 11], all of which influence the evaluation of sexual life. It is closely linked to overall health and subjective well-being [12]. Furthermore, it represents a key component of psychosocial functioning and strongly correlates with overall quality of life [13]. Problems in this area may lead to mental health issues, reduced self-esteem, tension in interpersonal relationships, and negative effects on professional performance – including among the professional group of healthcare workers [14, 15]. Due to the physical, mental, and social demands of their profession, healthcare workers may be at higher risk of experiencing diminished sexual functioning [16–19].

Excess body weight is associated with sexual functioning, making it an important area of interest in public health [20]. Individuals with excess body weight are more likely to experience reduced libido, lower sexual satisfaction, erectile dysfunction, and negative body image, which are linked to lower satisfaction and decreased sexual activity [21, 22]. The potential mechanisms underlying the negative impact of excess body weight on sexual functioning are multifactorial and complex. These include the secretory activity of adipose tissue, the relationship between excess body weight and comorbidities affecting sexual health, hormonal imbalances, and psychological factors such as body image concerns, low self-esteem, and fear of rejection, all of which can lead to sexual dysfunction [23–26].

Given the increasing prevalence of overweight and obesity, a more comprehensive approach is necessary to analyze their relationship with sexual functioning. This approach should consider both direct and indirect mechanisms, including self-rated health and perceived physical attractiveness. Previous studies have primarily focused on sexual dysfunctions in women and erectile disorders in men. However, since sexuality involves multiple dimensions, including interactions between psychosomatic mechanisms inherent to individuals and the socio-cultural contexts in which they operate, it is essential to move beyond the analysis

of purely physical sexual dysfunctions. Instead, a broader perspective on the components of sexual health should be adopted.

The aim of the present study was to examine the extent to which body weight impacts sexual life evaluation among healthcare workers in Poland, considering such mediating factors as self-rated health, perceived physical attractiveness, stress, and sleep disturbances.

## Material and methods

### Design of the study

A nationwide cross-sectional quantitative study was conducted in Poland between 21 February and 28 April 2022, as part of a project examining the health and health behavior of health care workers. The sampling frame consisted of a database of 2,894 hospitals and outpatient clinics across all of Poland's 16 provinces (voivodships). The facilities were selected at random to ensure a representative distribution across demographic and healthcare structures in Poland. The analysis included staff from 99 randomly selected healthcare facilities nationwide, where the management consented to participate in the study. The study was conducted using a computer-assisted web interview (CAWI) supported by a paper-and-pencil interview (PAPI).

### Participants

A total of 1,478 respondents from three main employment categories in healthcare facilities were included in the analysis (physicians  $N = 407$ , nurses  $N = 928$ , paramedics  $N = 143$ ). The group sizes were diversified according to the project design assumptions, which specified the minimum target sample size, required representation of occupational groups, and geographic variation reflecting the employment structure in Polish healthcare [27]. Cases with missing data for any of the analyzed variables were excluded from the study. Consequently, respondents who did not respond to these variables were excluded from the analyses.

### Questionnaire

Respondents provided anonymous answers to closed-ended questions, primarily on nominal or ordinal scales, as well as visual analog scales. Participation in the survey was voluntary, with respondents providing informed consent and having the option to withdraw at any time.

### Research tools

In the present study, a subset of questions from the questionnaire was selected for analysis. The main outcome variable was a composite index

of sexual life evaluation, based on two questions developed by the research team. These questions were part of a broader block addressing various aspects of life, including *satisfaction with one's sexual life, sexual performance, and sexual capabilities*.

Responses were recorded using a five-point Likert scale, ranging from "very good" to "very poor," with an option to decline answering or skip the question for participants who had so far not become sexually active in their lives. The composite index ranged from 2 to 10 points, with higher scores indicating a better evaluation of sexual life. According to principal component analysis (PCA), this total index is homogeneous, with both factor loadings at 0.93, and demonstrates good internal consistency (Cronbach's  $\alpha = 0.93$ ).

The independent variables were defined such that higher levels indicated greater dysfunction.

### Body weight

The study included a question about current body weight in kilograms and height in centimeters, allowing for the calculation of BMI using the formula: weight divided by height squared ( $\text{kg}/\text{m}^2$ ). The following BMI categories were applied: 18.5–24.99 for normal weight, 25–29.99 for overweight, and  $>30$  for obesity.

### Self-assessment of health problems

Subjective health problems were assessed using a question from the same set of items that evaluated different aspects of life, including the dependent variable. The response options were standardized and grouped into three categories by combining the most extreme answers (very good + good rating and bad + very bad rating).

### Body image self-assessment

Another question from the same block focused on participants' subjective evaluations of their own physical attractiveness, which served as an indicator of body image self-assessment. Responses were recoded into three categories by combining the most extreme answers (very good + good rating and bad + very bad rating).

### Stress

Stress levels were measured using the 4-item Perceived Stress Scale (PSS-4), also known as Cohen's scale [28]. The scale includes five response categories, ranging from "never" to "very often", based on experiences during the past month. The total score ranges from 0 to 16 points, with higher scores reflecting higher stress levels (two items were reverse-coded in this respect). According to PCA analysis, the scale is not homogeneous, with

factor loadings of 0.91 and 0.92, respectively, and a single-factor reliability score of 0.57.

### Sleep

Sleep disturbances over the past month were evaluated using the 4-item Jenkins Sleep Scale (JSS-4). Responses ranged from "0" (not at all) to "5" (22–31 days). Total scores for sleep disturbances range from 0 to 20, with higher scores indicating more frequent sleep disturbances [29]. According to PCA analysis, the scale is homogeneous, with factor loadings from 0.84 to 0.92, and demonstrates good internal consistency (Cronbach's  $\alpha = 0.90$ ).

In addition, demographic and social characteristics were included in the analyses: sex, age, relationship status, and occupation.

### Statistical analysis

The factor structure of the applied scales was evaluated using PCA, and their reliability was assessed with Cronbach's  $\alpha$ .

In univariate analysis, correlations between continuous, quasi-continuous, or ordinal variables were examined using Spearman's coefficient. For comparisons of means, the nonparametric Mann-Whitney *U* test (for two independent samples) or the Kruskal-Wallis *H* test (for three independent samples) was applied. In the latter case, post hoc multiple comparisons with Bonferroni correction were also performed.

For multivariate analysis, a hierarchical linear regression model was estimated for the dependent variable – the sexual life evaluation index. Results from the four steps of the analysis were presented as unstandardized betas with 95% confidence intervals (CI), and model fit was assessed using the coefficient of determination ( $R^2$ ).

To identify complex pathways of relationships between variables, structural equation modeling (SEM) without latent variables was used. The models analyzed the direction of hypothetical relationships, the significance of path coefficients, root mean square error of approximation (RMSEA) with 95% CI, comparative fit index (CFI), normed fit index (NFI), and relative fit index (RFI). Modification indices (MI) were also considered, adding potential connections as needed, but without correlating error terms. The model for the entire sample was modified until all non-significant paths were eliminated and MI values fell below 4. Separate models for men, women, and healthcare worker groups were estimated using the same framework without further modifications, to highlight any non-significant relationships between variables. The analysis also examined matrices of total, direct, and indirect effects.

SPSS 29.0 and AMOS 29.0 (IBM Corp. Released 2024. IBM SPSS Statistics for Windows, Version 29.0. IBM Corp., Armonk, NY, USA) were used for data analysis. A significance level of  $p < 0.05$  was adopted for all tests.

## Results

### Demographics and sexual life evaluation index

The basic demographics of the analyzed sample are presented in Table I. The sample had the largest proportion of nurses (62.8%), followed by physicians (27.5%), and the smallest group consisted of paramedics (9.7%). The high proportion of women (77.6%) reflects the employment structure in Polish healthcare facilities. The average age of respondents was 46.74 years, with the highest average age observed among nurses (48.12 years) and the lowest among paramedics (37.68 years). The majority of respondents were over 50 years old (46.1%) (respondents over 65 years constitut-

**Table I.** Sexual life evaluation index by socio-demographic characteristics and BMI

Variable	N (%)	Sexual life evaluation index Mean (SD)
Total	1478	7.51 (1.86)
Sex		
Male	331 (22.4)	7.69 (1.73)
Female	1147 (77.6)	7.46 (1.89)
MW – p		0.092
Age		
20–29	156 (10.6)	7.99 (1.79)
30–49	641 (43.4)	7.61 (1.91)
> 50	681 (46.1)	7.31 (1.79)
KW – p		< 0.001
Relationship status		
Single	1305 (88.3)	7.61 (1.83)
In relationship	173 (11.7)	6.80 (1.91)
MW – p		< 0.001
Occupation		
Physician	407 (27.5)	7.38 (1.86)
Nurse	928 (62.8)	7.49 (1.87)
Paramedic	143 (9.7)	8.04 (1.68)
KW – p		< 0.001
BMI		
Normal	669 (45.3%)	7.72 (SD = 1.84)
Overweight	546 (36.9%)	7.47 (SD = 1.81)
Obese	263 (17.8%)	7.07 (SD = 1.93)
KW – p		< 0.001

MW – Mann-Whitney; KW – Kruskal-Wallis.

ed 2.1% of the total sample) and were currently in a relationship (88.3%). Respondents with excess body weight accounted for 54.7% of the sample.

Table I also shows the average evaluation of sexual life across different respondent groups. The overall mean index for the entire sample was  $7.51 \pm 1.86$ . Statistically significant differences ( $p < 0.001$ ) were found in relation to all factors except sex, where the result was borderline. Higher sexual life evaluations were reported by respondents aged 20–29, those in relationships, paramedics, and those with normal BMI.


Given the main focus of the present study, it is important to highlight the differences between BMI groups, particularly the consistent decline in sexual life self-assessment with increasing levels of excess body weight ( $p < 0.001$ ). Post hoc analysis revealed significant differences between all three group pairs – normal weight vs. overweight ( $p = 0.012$ ), normal weight vs. obesity ( $p < 0.001$ ), and overweight vs. obesity ( $p = 0.016$ ).

Figure 1 plots the sexual life evaluation index by BMI, divided by sex. Women in all BMI categories rated their sexual lives lower than men, with sex differences being significant in the overweight group ( $p = 0.036$ ). Despite a clear downward trend observed in both sexes, statistically significant differences between BMI groups were found only among women ( $p < 0.001$ ).

### Sexual life evaluation based on socio-demographic characteristics, BMI, and life assessment

Table II presents potential relationships between the analyzed continuous, quasi-continuous, and ordinal variables.

Sexual life evaluation, as a positively oriented variable, showed a significant negative correlation with age, health assessment, physical attractiveness assessment, stress, sleep, and BMI ( $p < 0.001$ ). The strongest correlations with sexual life evaluation were observed for health



**Figure 1.** BMI and sexual life evaluation index by sex

**Table II.** Spearman's correlations (rho) between analyzed variables

	M ± SD	1	2	3	4	5	6
1 Sexual life evaluation	7.51 ±1.86				1		
2 Age	Rho	46.74 ±11.30	-0.152			1	
	P-value		< 0.001				
3 Health problems self-assessment	Rho	2.25 ±0.85	-0.496	0.042		1	
	P-value		< 0.001	0.110			
4 Body image self-assessment	Rho	2.40 ±0.98	-0.513	-0.032	0.543		1
	P-value		< 0.001	0.223	< 0.001		
5 Stress	Rho	5.75 ±2.94	-0.207	-0.107	0.243	0.154	
	P-value		< 0.001	< 0.001	< 0.001	< 0.001	
6. Sleep	Rho	5.37 ±2.94	-0.290	0.031	0.383	0.304	0.236
	P-value		< 0.001	0.237	< 0.001	< 0.001	< 0.001
7. BMI	Rho	26.00 ±4.34	-0.140	0.234	0.161	0.239	-0.004
	P-value		< 0.001	< 0.001	< 0.001	< 0.001	0.023

**Table III.** Hierarchical linear regression\* for variability in sexual life valuation index

Variable	B	95%CI(B)		P-value
		Low	High	
<b>Model 1</b>				
Constant	10.827	9.967	11.688	< 0.001
BMI	-0.062	-0.084	-0.040	< 0.001
Age	-0.015	-0.024	-0.007	< 0.001
R <sup>2</sup>			0.065	
<b>Model 2</b>				
Constant	11.932	11.164	12.701	< 0.001
BMI	-0.022	-0.042	-0.002	0.034
Age	-0.017	-0.024	-0.009	< 0.001
Health problems assessment	-1.003	-1.100	-0.906	< 0.001
R <sup>2</sup>			0.270	
<b>Model 3</b>				
Constant	12.199	11.429	12.968	< 0.001
BMI	-0.024	-0.044	-0.004	0.017
Age	-0.018	-0.026	-0.011	< 0.001
Health problems assessment	-0.862	-0.967	-0.757	< 0.001
Stress	-0.071	-0.100	-0.042	< 0.001
Sleep	-0.033	-0.050	-0.016	< 0.001
R <sup>2</sup>			0.291	
<b>Model 4</b>				
Constant	12.335	11.606	13.063	< 0.001
BMI	0.006	-0.013	0.025	0.533
Age	-0.025	-0.032	-0.018	< 0.001
Health problems assessment	-0.529	-0.640	-0.417	< 0.001
Stress	-0.063	-0.091	-0.036	< 0.001
Sleep	-0.022	-0.038	-0.005	0.010
Impaired body image self-assessment	-0.637	-0.733	-0.541	< 0.001
R <sup>2</sup>			0.365	

\*Adjusted for sex, relationship status, and occupation; BMI, age, stress, and sleep were treated as continuous variables, while health and body image assessments were treated as ordinal variables based on the original 5 categories.

problems and two additional factors – impaired body image self-assessment and sleep disturbances.

In contrast, BMI showed a significant positive correlation with age, health problems, and impaired body image self-assessment ( $p < 0.001$ ).

### Variability in sexual life evaluation

Table III presents four regression models. Each model was adjusted for sex, relationship status, and occupation, which were recoded into two dummy variables (data not shown here).

In Model 1, BMI and age were identified as significant predictors of sexual life evaluation ( $R^2 = 0.065$ ;  $p < 0.001$ ).

In Model 2, an additional variable – health problems self-assessment – was introduced, leading to a substantial increase in  $R^2$  (0.270). BMI and age remained significant predictors.

Model 3 incorporated two more variables – stress and sleep – which resulted in a slight further increase in  $R^2$  (0.291), without altering the significance of other parameters.

Finally, Model 4 added impaired body image self-assessment, which significantly improved the model fit, as measured by  $R^2$  (0.365). The inclusion of this new factor – perceived physical attractiveness – eliminated BMI as a predictor of variability in sexual life evaluation. Additionally, the beta coefficient for age increased, whereas it decreased for other independent variables.

In the next step, a path model was estimated to identify the more complex mechanism of relationships between the analyzed variables, distin-

guishing between direct and indirect effects. This analytical approach corresponds to the estimation of a set of regression equations in which BMI can be the dependent or independent variable, also indirectly influencing the variability of the sexual life assessment index. For the defined path model estimated for the entire sample, the results showed  $c_{min}/df = 1.140$ , and the fit indices were as follows:  $RMSEA = 0.017$  (95% CI: 0.000–0.042). The NFI, RFI, and CFI values were 0.996, 0.983, and 0.999, respectively.

As presented in Table IV and illustrated in the graphical representation of the model (Figure 2), the structure consists of six interconnected linear equations. Age, as a non-modifiable factor, acts solely as an independent variable, while sexual life evaluation is treated as the only dependent variable, without influencing the variability of other variables. In the resulting model, sexual life evaluation has five significant predictors. BMI is determined only by age and is indirectly associated with sexual life evaluation, primarily through its link to physical attractiveness assessment. Sleep and stress levels are also associated with sexual life evaluation.


Of particular note is the negative assessment of physical attractiveness, which was found to have four significant predictors and emerged as the strongest factor influencing variability in sexual life evaluation.

The matrices of indirect and direct effects (Appendix 1, Supplementary Tables SI–SIII) indicate that while physical attractiveness has the strongest direct effect on variability in sexual life eval-

**Table IV.** Standardized regression weights for SEM models in the general population

Path		Estimate	S.E.	P-value
BMI	HEALTH	0.185	0.005	< 0.001
BMI	ATRRA	0.193	0.005	< 0.001
HEALTH	JSS-4	0.323	0.146	< 0.001
HEALTH	SEX	-0.238	0.057	< 0.001
HEALTH	ATRRA	0.462	0.027	< 0.001
ATRRA	SEX	-0.345	0.048	< 0.001
ATRRA	PSS4	0.065	0.092	0.033
AGE	BMI	0.212	0.010	< 0.001
AGE	ATRRA	-0.107	0.002	< 0.001
AGE	PSS4	-0.106	0.007	< 0.001
AGE	SEX	-0.15	0.003	< 0.001
PSS4	SEX	-0.106	0.014	< 0.001
PSS4	JSS-4	0.183	0.043	< 0.001
PSS4	HEALTH	0.24	0.008	< 0.001
JSS-4	SEX	-0.063	0.008	0.007
JSS-4	ATRRA	0.091	0.004	< 0.001

HEALTH – health problems self-assessment, ATRRA – body image self-assessment, SEX – sexual life evaluation index, PSS-4 – stress, JSS-4 – sleep.



**Figure 2.** Model SEM in the general population

ation ( $-0.345$ ), health problems produce the strongest indirect effect ( $-0.197$ ). However, the second most important factor generating an indirect effect is BMI ( $0.150$ ), which is the only variable with a zero direct effect (Table IV). A comparison of gender specific SEM models is provided in Appendix 2.

## Discussion

This study evaluated the relationship between body weight and sexual life evaluation in healthcare workers in Poland, using both linear modeling and direct and indirect associations. While numerous studies suggest a link between increased body weight and impaired sexual functioning, only a few have conducted extensive analyses of both direct and indirect relationships [20, 30–32]. In the path model, five factors were directly correlated with variability in sexual life evaluation: impaired body image self-assessment, health assessment, elevated stress levels, sleep disturbances, and age. Our findings indicate that body weight is associated with sexual functioning through an indirect relationship with self-rated health and physical attractiveness.

Health issues among healthcare workers, including those related to excess body weight, are particularly important for public health, as they can impact their ability to provide patient care [33]. In our study, the average BMI was 26, which aligns with findings from other studies, where the average BMI among healthcare workers was 25.4 [34]. Additionally, our results show that BMI was significantly positively correlated with both age and self-rated health ( $p < 0.001$ ). Furthermore, weight gain among healthcare workers has shown an upward trend, exacerbated by the COVID-19 pandemic and its negative effects on both physical and mental health [35].

Global prevalence data indicate that reduced sexual life evaluation, including lower sexual functioning, represents a significant public health issue [36]. In our bivariate analysis, higher BMI was associated with poorer sexual life evaluation ( $p < 0.001$ ), which is consistent with other studies [20, 24]. However, self-rated health and physical attractiveness proved to be stronger differentiators of sexual life evaluation than BMI ( $p < 0.001$ ). In our study, hierarchical regression analysis showed that adding health assessment and physical attractiveness to the model eliminated BMI as a predictor of variability in sexual life evaluation. This suggests that these variables may play a mediating or moderating role in this correlation. These findings align with other studies evaluating the relationship between body weight and sexual functioning, where health factors and physical attractiveness were identified as key variables influencing this association [20, 23–25]. The results of the hierarchical model prompted further exploration of these connections.

The remainder of this discussion section primarily focuses on self-rated health and physical attractiveness, as these factors were most strongly associated with BMI and sexual life evaluation.

In our study, 29.1% of respondents reported a lower health assessment, including 9.3% who indicated serious health problems. Given its ability to predict mortality and morbidity, self-rated health is a critical indicator of population health [33, 37]. Our path model demonstrated that health assessment had a strong direct association with variability in sexual life evaluation ( $-0.238$ ), consistent with previous studies [20, 24].

Another key variable in our study is physical attractiveness (body-image self-assessment). Self-assessment reflects self-worth, encompassing beliefs about oneself and emotional reactions to those beliefs [38]. As such, self-assessment can

act as a predictor of life satisfaction, representing self-value in specific contexts and environments. Western studies on physical attractiveness have largely focused on body appearance, particularly weight and body shape [39], which are core components of body image [40]. In our study, 13.1% of respondents reported impaired physical attractiveness assessment, while 27% had moderately impaired assessments. Additionally, higher BMI was associated with lower body image ratings ( $p < 0.001$ ), a finding in line with other studies showing BMI's influence on perceptions of physical attractiveness [41]. Weaver and Byers [42] found that women who view themselves as more attractive report higher sexual satisfaction, likely due to greater confidence and comfort in intimate relationships. These findings are consistent with a meta-analysis by Cash and Pruzinsky, which identified positive body image as a key predictor of high sexual satisfaction [43]. In our study, the path model demonstrated that physical attractiveness assessment had the strongest direct relationship with variability in sexual life evaluation ( $-0.345$ ). These results are supported by prior studies, which showed that body image and self-perceived attractiveness strongly correlate with overall life satisfaction, including sexual satisfaction, by enhancing self-esteem and emotional well-being [44, 45]. These findings reinforce that a positive body image, including views of genital appearance and self-perception, is associated with higher sexual satisfaction and better sexual life evaluation. This highlights the need for interventions aimed at improving body image and addressing cultural pressures related to appearance [46].

In our study, other factors potentially influencing sexual life evaluation were also identified. Age was negatively associated with sexual life evaluation through lower self-rated attractiveness. Older age is also linked to poorer health outcomes, higher risk of sexual dysfunctions, and a greater number of individuals without romantic partners, which may further impact sexual functioning [47]. Sleep disturbances also play a key role in human health, including mental, physical, and sexual health. Greater sleep disturbances were associated with poorer sexual life evaluation, while better sleep was linked to higher scores [48, 49]. Elevated stress levels also emerged as a significant factor negatively influencing sexual life evaluation, consistent with previous research [50]. Finally, sex differences were observed, with women reporting poorer sexual life evaluations as BMI increased ( $p < 0.001$ ).

This study was cross-sectional in design; therefore, causal relationships could not be established. However, the use of structural equation modeling (SEM) enabled the identification of complex path-

ways between key variables and allowed for the determination of influence trajectories. The sexual life evaluation index was based on two questions that are not part of a validated scale. Nevertheless, these questions have been successfully used since 1997 in cyclical studies on sexuality and health. The analyses also did not consider the presence of comorbidities, which may affect sexual functioning. However, self-rated health, one of the key variables, is closely related to the occurrence of comorbidities [37].

Further research should include longitudinal studies to better illustrate cause-and-effect relationships. Future studies assessing the relationship between body weight and sexual functioning should consider changes in body weight as a key indicator and examine the interactions of these factors in more depth. Additionally, future research should include a detailed assessment of sexual functioning using validated scales to capture a more comprehensive picture of the relationships involved.

In conclusion, excess body weight is associated with numerous chronic diseases, lower self-esteem, and reduced sexual life evaluation. Our study provides a comprehensive analysis of the correlation BMI and sexual life evaluation, exploring both direct and indirect associations, while accounting for self-rated health and physical attractiveness. The findings highlight a strong link between body weight and sexual life evaluation, primarily mediated by self-rated health and, in particular, physical attractiveness, which emerged as the strongest predictor in our study. In the future, educational programs addressing sexual health should be developed and implemented among individuals with excess body weight as part of treatment strategies.

## Funding

Funding source: Medical Research Agency.

## Ethical approval

Ethics Committee of the Faculty of Education at the University of Warsaw (decision no. 2021/8).

## Conflict of interest

The authors declare no conflict of interest.

## References

1. Dai H, Alsalhe TA, Chalghaf N, Riccò M, Bragazzi NL, Wu J. The global burden of disease attributable to high body mass index in 195 countries and territories, 1990–2017: an analysis of the Global Burden of Disease Study. *PLoS Med* 2020; 17: e1003198.
2. Jaacks LM, Vandevijvere S, Pan A, et. al. The obesity transition: stages of the global epidemic. *Lancet Diabetes Endocrinol* 2019; 7: 231-40.

3. Welsh A, Hammad M, Piña IL, Kulinski J. Obesity and cardiovascular health. *Eur J Prev Cardiol* 2024; 31: 1026-35.
4. World Health Organization. Obesity and overweight; 2024. <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight> [Accessed February 15, 2025].
5. Fontaine KR, Barofsky I. Obesity and health-related quality of life. *Obes Rev* 2001; 2: 173-82.
6. Dixon JB. The effect of obesity on health outcomes. *Mol Cell Endocrinol* 2010; 316: 104-8.
7. Esfahani SB, Pal S. Obesity, mental health, and sexual dysfunction: a critical review. *Health Psychol Open* 2018; 5. <https://doi.org/10.1177/20551029187868>.
8. Epstein S, Mamo L. The proliferation of sexual health: Diverse social problems and the legitimization of sexuality. *Soc Sci Med* 2017; 188: 176-90.
9. Masoudi M, Maasoumi R, Bragazzi NL. Effects of the COVID-19 pandemic on sexual functioning and activity: a systematic review and meta-analysis. *BMC Public Health* 2022; 22: 189.
10. DeLamater J, Karraker A. Sexual functioning in older adults. *Curr Psychiatry Rep* 2009; 11: 6-11.
11. Wallwiener S, Strohmaier J, Wallwiener LM, et al. Sexual function is correlated with body image and partnership quality in female university students. *J Sex Med* 2016; 13: 1530-8.
12. Stephenson KR, Meston CM. The conditional importance of sex: exploring the association between sexual well-being and life satisfaction. *J Sex Marital Ther* 2015; 41: 25-38.
13. Knowles SR, Gass C, Macrae F. Illness perceptions in IBD influence psychological status, sexual health and satisfaction, body image and relational functioning: a preliminary exploration using structural equation modeling. *J Crohns Colitis* 2013; 7: e344-50.
14. Izdebski Z, Mazur J. *Zdrowie, życie seksualne i funkcjonowanie w związku z okresem pandemii COVID-19 [Health, sex life and relationship functioning during the COVID-19 pandemic]*. PWN; Poland 2024.
15. Izdebski Z, Mazur J, Furman K, Kozakiewicz A, Białorudzki M. Humanization of the treatment process and clinical communication between patients and medical staff during the COVID-19 pandemic. University of Warsaw; Poland 2023.
16. Gray P, Senabe S, Naicker N, Kgalamono S, Yassi A, Spiegel JM. Workplace-based organizational interventions promoting mental health and happiness among healthcare workers: a realist review. *Int J Environ Res Public Health* 2019; 16: 4396.
17. Eroglu U, Balci M, Coser S, et al. Impact of the COVID-19 Pandemic on the Psychosexual Functions of Healthcare Workers. *J Sexual Med* 2022; 19: 182-7.
18. Pan A, Wu Y, Chen X, et al. A qualitative study of psychological stress among China's frontline nurses fighting COVID-19. *Arch Med Sci* 2022; 18: 1407-12.
19. Marczewski KP, Piegza M, Gospodarczyk NJ, et al. Evaluation of selected factors influencing sleep disorders in paramedics during the COVID-19 pandemic. *Arch Med Sci* 2024; 20: 86-93.
20. Sarwer DB, Hanson AJ, Voeller J, Steffen K. Obesity and sexual functioning. *Curr Obes Rep* 2018; 7: 301-7.
21. McNabney SM, Gletsu-Miller N, Rowland DL. Sexual function and satisfaction in the context of obesity. *Curr Diabetes Rep* 2023; 23: 315-27.
22. Kolotkin RL, Zunker C, Østbye T. Sexual functioning and obesity: a review. *Obesity* 2012; 20: 2325-33.
23. Harvey I, Boudreau A, Stephens JM. Adipose tissue in health and disease. *Open Biol* 2020; 10: 200291.
24. Rowland DL, McNabney SM, Mann AR. Sexual function, obesity, and weight loss in men and women. *Sex Med Rev* 2017; 5: 323-38.
25. Sarwer DB, Steffen KJ. Quality of life, body image and sexual functioning in bariatric surgery patients. *Eur Eat Disord Rev* 2015; 23: 504-8.
26. Finkelstein EA, Khavjou OA, Thompson H, et al. Obesity and severe obesity forecasts through 2030. *Am J Prev Med* 2012; 42: 563-70.
27. Matyszko K, Pędziński B, Maślach D, Krzyżak M, Marciniowicz L. Medical staff in Poland in 2012-2022 - challenges related to the distribution of human resources. *Ann Agric Environ Med* 2024; 31: 382-7.
28. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983; 24: 385-96.
29. Monterrosa-Castro Á, Portela-Buelvas K, Salguedo-Madrid M, Mo-Carrascal J, Duran-Méndez Leidy C. Instruments to study sleep disorders in climacteric women. *Sleep Sci* 2016; 9: 169-78.
30. Kolotkin RL, Binks M, Crosby RD, Østbye T, Gress RE, Adams TD. Obesity and sexual quality of life. *Obesity* 2006; 14: 472-9.
31. Moore RH, Sarwer DB, Lavenberg JA, et al. Relationship between sexual function and quality of life in obese persons seeking weight reduction. *Obesity* 2013; 21: 1966-74.
32. Syed AH, Chandnani A, Khan A, et al. Association of weight loss with improved sexual function in females. *Cureus* 2021; 13: e16849.
33. Hartanto A, Yong JC, Toh WX. Bidirectional associations between obesity and cognitive function in midlife adults: a longitudinal study. *Nutrients* 2019; 11: 2343.
34. Lee DH, Keum N, Hu FB, et al. Predicted lean body mass, fat mass, and all cause and cause specific mortality in men: prospective US cohort study. *BMJ* 2018; 362: k2575.
35. Chen ZM, Lin RT. Rotating shift and BMI increase among healthcare workers in a military hospital: pre- and post-pandemic analysis in Taiwan. *Ann Occup Environ Med* 2024; 36: e15.
36. Dunn KM, Croft PR, Hackett GI. Satisfaction in the sex life of a general population sample. *J Sex Marital Ther* 2000; 26: 141-51.
37. Lorem G, Cook S, Leon DA, Emaus N, Schirmer H. Self-reported health as a predictor of mortality: a cohort study of its relation to other health measurements and observation time. *Sci Rep* 2020; 10: 4886.
38. McClure AC, Tanski SE, Kingsbury J, Gerrard M, Sargent JD. Characteristics associated with low self-esteem among US adolescents. *Acad Pediatr* 2010; 10: 238-44.e2.
39. Tiggemann M. Body image across the adult life span: stability and change. *Body Image* 2004; 1: 29-41.
40. Biro FM, Striegel-Moore RH, Franko DL, Padgett J, Bean JA. Self-esteem in adolescent females. *J Adolesc Health* 2006; 39: 501-7.
41. Watkins JA, Christie C, Chally P. Relationship between body image and body mass index in college men. *J Am Coll Health* 2008; 57: 95-100.
42. Weaver AD, Byers ES. The relationships among body image, body mass index, exercise, and sexual functioning in heterosexual women. *Psychology Women Quart* 2006; 30: 333-9.
43. Cash TF, Pruzinsky T. *Body Image: A Handbook Of Theory, Research, And Clinical Practice*. Guilford Press 2002.
44. Satinsky S, Reece M, Dennis B, Sanders S, Bardzell S. An assessment of body appreciation and its relationship to sexual function in women. *Body Image* 2012; 9: 137-44.

45. Meland E, Breidablik HJ, Thuen F, Samdal GB. How body concerns, body mass, self-rated health and self-esteem are mutually impacted in early adolescence: a longitudinal cohort study. *BMC Public Health* 2021; 21: 496.
46. Alavi-Arjas F, Goodman MP, Simbar M, Alavi Majd H, Nahidi F. The strength of correlation between female genital self-image and sexual function: a systematic review and meta-analysis. *J Sex Med* 2023; 20: 1376-83.
47. Izdebski Z. *Zdrowie i życie seksualne Polek i Polaków w wieku 50-74 lat w 2017 roku. Perspektywa starzejącego się społeczeństwa [Health and sex life of Polish men and women aged 50-74 in 2017 A perspective of an ageing society]*. Warsaw University Publishers, Poland 2023.
48. Biaforudzki M, Undra M, Izdebski Z. Impact of body weight on the relationships between sleep quality in healthcare workers and their somatic health, sexual life, occupational burnout and stress. *Int J Occup Med Environ Health* 2024; 37: 545-56.
49. Zhu L, Gao Q, Guo X, Xu Z, Zhang J. Causal relationship between sleep traits and erectile dysfunction: evidence from Mendelian randomization analysis. *Arch Med Sci* 2025; 21: 597-604.
50. Biaforudzki M, Izdebski Z. Changes in the body mass of adult residents of rural and urban areas in the initial months of the COVID-19 pandemic vs. their mental, physical and sexual health. *Ann Agric Environ Med* 2021; 28: 667-75.