

Shared Decision Making in primary care: barriers and facilitators from the physician's and patient's perspective—a scoping review.

Keywords

primary care, barriers, communication, facilitators, doctor–patient relationship, shared decision making

Abstract

Introduction

Health-related communication and Shared Decision Making (SDM) are crucial elements of patient-centered care. Multiple studies have showed that the decision-making process is often far from ideal, where an ideal process involves the patient being active, understanding their treatment options, and expressing preferences clearly, while the physician has sufficient time and willingness to share knowledge and involve the patient in the decision-making process. The aim of this review was to identify the key barriers and facilitators influencing the adoption of Shared Decision Making (SDM) approach from both the physician's and patient's perspectives in primary healthcare in relation to the 5-step model of SDM.

Material and methods

A scoping review of resources published within the last five years was conducted in June 2024 on the basis of PRISMA-ScR. Specific barriers hindering the adoption of SDM were identified, which can arise at any of the five steps.

Results

The most common barriers include: time constraints and physician's workload, lack of SDM culture and tradition, paternalistic model of healthcare, patients' passive attitude and belief they are unable to make decisions. SDM facilitators include: trust, use of educational materials by physicians, belief that SDM will lead to better treatment outcomes, and training in communication for physicians.

Conclusions

Despite the challenges to public health, action should be taken to overcome these barriers, given the well-established benefits of SDM, although this will require time and necessary adaptations.

1 Title: Shared Decision Making in primary care: barriers and facilitators from the physician's
2 and patient's perspective—a scoping review.

3 Abstract

4 Background: Health-related communication and Shared Decision Making (SDM) are
5 crucial elements of patient-centered care. The aim of this review was to identify the key
6 barriers and facilitators influencing the adoption of Shared Decision Making (SDM)
7 approach from both the physician's and patient's perspectives in primary healthcare in
8 relation to the 5-step model of SDM.

9 Methods: A scoping review of resources published within the last five years was
10 conducted in June 2024 on the basis of PRISMA-ScR. Specific barriers hindering the
11 adoption of SDM were identified, which can arise at any of the five steps.

12 Results: The most common barriers include: time constraints and physician's workload,
13 lack of SDM culture and tradition, paternalistic model of healthcare, patients' passive
14 attitude and belief they are unable to make decisions. SDM facilitators include: trust, use
15 of educational materials by physicians, belief that SDM will lead to better treatment
16 outcomes, and training in communication for physicians.

17 Conclusions: Despite the challenges to public health, action should be taken to overcome
18 these barriers, given the well-established benefits of SDM, although this will require time
19 and necessary adaptations.

20
21 Key words: shared decision making, barriers, facilitators, primary care, communication,
22 doctor–patient relationship

23 Introduction

24 Effective communication plays a key role in the healthcare decision-making process [1].
25 Health-related communication and Shared Decision Making (SDM) are crucial elements of
26 patient-centered care [2,3]. The evolution of the SDM concept originated from the work on
27 informed consent that began in the 1980s [4,5]. Advances in information and data
28 management (collection, analysis, sharing), along with greater access to information, have
29 led to expectations that patients will take a more active role in managing their health. SDM
30 is based on three pillars: two-way exchange of information between the patient and the
31 physician, identifying and acknowledging the patient's preferences, and engaging both the
32 patient and the physician in the final decision-making process [6]. In 2015, the Agency for
33 Healthcare Research and Quality (AHRQ) released a revised, more practical 5-step model
34 of SDM, replacing the previous 9-step version, called SHARE (Seek, Help, Assess,
35 Reach, Evaluate) [7]. The objective of the first step is to understand the situation and help
36 the patient become aware of the aspects that demand action ("Seek your patient's
37 participation"). In the second step, the physician should provide the patient with evidence-
38 based information about the available treatment options ("Help your patient explore and
39 compare treatment options"). The next step should take into account the patient's values
40 and preferences in the context of the suggested treatment options ("Assess your patient's
41 values and preferences"). Finally, both parties should make a decision together ("Reach a
42 decision with your patient") and evaluate it ("Evaluate your patient's decision") to ensure it
43 balances benefits, risks, and the patient's preferences, maximizing the likelihood of
44 achieving the desired outcome. [8] In primary care, multiple health-related decisions are
45 made during each visit [9]. Studies have showed that engaging patients in the decision-
46 making process can reduce regret and anxiety related to the chosen option, improve
47 treatment outcomes, satisfaction, and quality of life, as well as enhance the quality and
48 safety of healthcare [10,11]. Engaging patients in decision-making conversations about

their chronic condition is crucial when it comes to lifestyle, as it encourages them to adopt necessary changes [12]. This is particularly important for decisions involving long-term actions, such as physiotherapy for chronic pain or changing the diet due to food allergy [13-15]. In addition, SDM significantly improves health literacy and reduces decision-related conflicts [16]. In contrast, poor SDM practices have been linked to worse patient-reported quality outcomes, lower overall quality of care, and increased dependence on healthcare services [11].

Multiple studies have showed that the decision-making process is often far from ideal, where an ideal process involves the patient being active, understanding their treatment options, and expressing preferences clearly, while the physician has sufficient time and willingness to share knowledge and involve the patient in the decision-making process [16]. The debate on SDM has been going on for many years, yet many physicians still believe that making decisions together with the patient is difficult, mainly due to social, cultural (lack of SDM tradition), and organizational reasons (healthcare systems do not recognize SDM as a standard of care) [18-20]. One major barrier to effective SDM implementation in medical practice is the lack of a universal definition and consistent understanding of SDM. There are differing perspectives on the definition of shared decision-making and how it should be implemented in practice. [21] The way SDM is conceptualized influences how its feasibility is assessed. Examples of SDM conceptualizations include: negative right, informed decision-making, and tailored partnership. Conceptualizing SDM as a "negative right" implies that patients have the right to refuse treatment, but do not necessarily take an active role in selecting therapeutic options. Decisions are primarily made by physicians based on professional standards, with patient involvement emerging mainly when they disagree with the proposed course of action. The informed decision-making model assumes that the physician provides the patient with medical knowledge and available options, while the patient independently

makes the treatment decision. In this view, the physician acts as an advisor rather than a decision-maker, although in practice physicians often offer recommendations, as patients tend to rely on their expertise and authority. Conceptualizing SDM as a tailored partnership emphasizes a flexible approach that takes into account the individual characteristics of the patient, such as age, health condition, and personal preferences. Decisions are made jointly, but the process may vary—potentially even including a jointly agreed-upon, more paternalistic approach when appropriate. [21] The article is based on Makula's definition of SDM, where it is defined as an approach in which clinicians and patients collaborate by sharing the best available evidence in making health care decisions, while supporting patients in weighing their options to arrive at informed preferences. [22] Communication and interaction between healthcare providers and patients is key for health-literate decision-making [23].

Existing overviews focus on specific elements, such as barriers, facilitators, or either the patient's or physician's perspective. This review integrates the barriers, facilitators and both clinician and patient perspectives, framed within the 5-step SHARE model of SDM, to provide a comprehensive understanding of its challenges and opportunities in primary care.

Methodology

The scoping review was conducted on the basis of the PRISMA- ScR instruction (Figure 1). The framework was applied as follows:

Identification: A comprehensive search strategy was developed based on predefined keywords: barriers, facilitators, shared decision making, general practice, primary care, patients perspective, GP's perspective. The scoping review was conducted in June 2024 and included the most recent studies conducted over the past five years. The following databases were searched: Pub Med, Cochrane, Willey, and Goggle Scholar.

Screening: After removal of duplicates, all titles and abstracts were independently screened by two reviewers to assess relevance against the eligibility criteria. Studies were included if they: (1) were an original collection of data, systematic review, scoping review or meta-analysis; (2) involved adult population only (aged >19); (3) looked at SDM in primary care; (4) captured SDM barriers and facilitators; (5) the results spanned the last five years. They also included guidelines, instructions, and SDM recommendations published on government websites. Studies not meeting the inclusion criteria were excluded at this stage. The exclusion criterion was studies involving hospital settings, pediatric populations, or patients with mental illness or cancer (Table S1).

Eligibility: Titles, abstracts, and potentially relevant articles were reviewed independently in full text by the authors (KDŽ and MKP) based on the eligibility criteria. Doubts were resolved through discussion (Figure 1). Due to the scoping nature of this review and in accordance with PRISMA-ScR guidelines, no formal critical appraisal of the quality of the study was conducted. The aim of this review was to map the existing literature and identify key barriers and facilitators to shared decision-making (SDM) in primary care, rather than to assess the effectiveness or rigour of interventions. Given the methodological heterogeneity of the included studies (e.g. qualitative interviews, surveys, reviews, policy documents), the use of a standardised quality assessment tool was considered inappropriate. However, to ensure the relevance of the included sources, all studies were independently screened by two reviewers using pre-defined inclusion and exclusion criteria, and any disagreements were resolved through discussion.

Inclusion: Studies meeting all criteria were included in the final synthesis.

Data extraction and quality assessment: An inductive thematic analysis was used to group findings into common themes. Results were presented narratively and mapped across the SDM framework from both the physician and patient perspectives.

1. Data screening - The researchers independently reviewed the full texts of all included studies to gain an in-depth understanding of their content, with a particular focus on reported barriers and facilitators of SDM from both the physician and patient perspectives.

2. Initial Coding - Two researchers (e.g., KDŽ and MKP) independently coded relevant segments of text related to SDM barriers and facilitators. Coding was conducted inductively, with codes assigned to data extracts that described specific factors impeding or supporting SDM.

3. Search for Themes - The identified codes were then grouped into broader categories or themes that reflected common patterns (e.g., time constraints, health literacy, communication style, decision-making roles). Barriers and facilitators were organized within the framework of the 5-step SHARE model of SDM (Seek, Help, Assess, Reach, Evaluate).

4. Reviewing Themes - The initial thematic structure was discussed and refined by the research team to ensure consistency and coherence. Redundant or overlapping themes were merged, and ambiguous codes were clarified.

5. Defining and Naming Themes - Final themes were clearly defined, labeled, and classified into either “barriers” or “facilitators.” Each theme was mapped to the corresponding SDM step and perspective (physician or patient), providing a structured thematic matrix.

6. Data Extraction and Synthesis - Themes were entered into a standardized extraction table, summary findings from the original studies. This matrix was used to compare thematic patterns across the five SDM steps and between physician and patient perspectives.

7. Validation and Consensus- The final thematic framework was reviewed by two independent experts in SDM. Disagreements were resolved through discussion with the principal investigator to ensure the validity and reliability of the thematic interpretation.

Results

Table I shows that barriers were identified for each step of SDM, both from the patient's and the physician's perspective. A shared barrier in the first step is the lack of a culture or tradition of SDM [9, 25, 26]. In addition, physicians cite time constraints and view consultations involving SDM as burdensome [25-28]. Evidence suggests that this may ultimately result in a default decision-making approach with little or no personalization. [29] Time pressures affect the physician's intuitive prioritization of which decisions to discuss. In this context, the diversity of patients and their problems also poses a challenge. There is concern that SDM will extend the length of the visit and thus unpredictably affect the physician's schedule. [30,31] On the other hand, a passive attitude—stemming from preconceived notions about consultations and socio-demographic factors like age, education, or gender—is a major barrier to patient engagement in the SDM process. This passive attitude is more commonly observed among elderly patients, males, and less educated individuals [23-39]. SDM facilitators in the first step include trust and recognizing that there are two experts in the clinical encounter [35]. Patients with the following personal traits tend to be more engaged in SDM: kind, self-confident, empowered, proactive [35]. Also, physicians who believe in the beneficial influence of SDM on patient outcomes and have good communication skills find it easier to implement SDM [24].

When treatment options are discussed and compared, SDM may be impeded if the patient feels overloaded by the amount of information, does not understand the information due to overly technical language, is unable to compare different options because only one was

presented, or perceives the physician as unwilling to provide explanations [25,36,37].
Conversely, the physician may be unable to discuss treatment options with the patient due
to time constraints, an overloaded consultation schedule, or a focus on quantity over
quality in patient consultations [25, 33].

In the third step, factors impeding the assessment of the patient's values and preferences
include the patient's hesitation in expressing their preferences or the physician's
unwillingness to listen to and acknowledge the patient's concerns and opinions. This stage
can be supported if the patient is aware of their preferences while the physician is caring,
empathic, calm and open [23, 27,35].

The most barriers on the patient's and physician's sides were identified in the fourth step,
where the decision should be made. From the patient's perspective, the key barriers to
making a treatment decision include: insufficient health literacy (difficulty accessing
detailed information before the visit), belief that only doctors can make decisions,
expectation that visits should provide definitive solutions, uncertainty about the best
option, lack of empowerment, financial limitations—only fully or partially state-funded
options are affordable, and additional stress from decision-making (acting under time
pressure) [24,25,28,34,36]. On the physician's side, barriers include: authority-based
decision-making style, reluctance to engage patients in the decision-making process—
believing they don't want to participate and it is not need—and a lack of communication
and interpersonal skills [24,25]. Supporting factors at this stage include: self-education
before the visit, fostering empowerment in self-care and responsibility for own decisions
and well-being, health literacy—understanding options, uncertainty in medical evidence,
and disease history—viewing the physician as a decision-making partner, having a nurse
present during consultations, and involving family members in the treatment process
[39,40]. Also, training for physicians supports shared decision-making [29]. In the last
step, the physician should evaluate the decision. A limiting factor at this stage can be the

physician's fear of facing a higher liability risk [25]. Furthermore, the decision may be difficult to evaluate in the long-term if there is no continuity in services—consultations are held by different specialists who give contradictory recommendations, causing confusion. Sometimes, the attending physician changes [25,35]. The evaluation process is supported when the physician feels responsibility is shared, providing a sense of protection from negative outcomes of the decision, or by maintaining detailed case records, documenting each visit, including concerns, patient preferences, and communication details. [25]

Table I Barriers and facilitators from the patient's and physician's perspectives in relation to 5 steps of SDM

Discussion

The shift towards the SDM model of healthcare is driven by the increasing number of treatment options, the growing complexity of the decision-making process, and the rising awareness that the paternalistic model does not translate to better patient outcomes [41]. Therefore, both SDM barriers and facilitators must be identified to enable more effective SDM integration into everyday medical practice. This review offers a wider perspective—that of patients and physicians at each of the five steps of SDM model.

Step 1: Seek your patient's participation

The barriers identified at this stage include primarily the lack of SDM tradition and culture. For cultural reasons, paternalism dominates in healthcare. On one hand, the physician makes decisions *a priori*, considering the patient's best interests, while on the other hand, the patient expects to be offered a ready-made solution, trusting the physician's knowledge and experience [9]. Shared Decision Making depends on a number of co-existing factors, including health literacy, patient's emotional status, patient's and physician's personal traits, quality of the patient-physician relationship, and the nature of the decision [42]. Studies have shown that elderly patients or less educated individuals rarely participate in SDM. Potential

reasons include difficulties in understanding information and a noticeably low level of engagement. This may discourage physicians from involving these patients in SDM, as they believe the patients are less willing or capable of participating in the decision-making process [43]. Moreover, patients tend to underestimate the value of their contribution to the consultation. Eager to please the physician, patients often feel they should conform and remain passive in the patient-physician relationship, believing they have limited permission to actively participate in the consultation [19,44]. However, patients' attitudes and beliefs can be modified through guidance (unlike unmodifiable socio-demographic factors, such as age), helping them accept a more active role [45]. To increase the chances of implementing SDM, physicians should be made aware of the benefits of building high-quality relationships with their patients. Usually, this requires improved communication skills [46]. In this context, personnel training is seen as the most effective intervention, enhancing relationship-building skills and facilitating the implementation of the SDM model as a standard in clinical practice [37]. On the other hand, time constraints has been identified as a significant barrier to introducing SDM. Studies have shown that improved time management could solve the majority of communication-related barriers [47].

Step 2: Help your patient explore and compare treatment options

At step two, where treatment options are discussed, the main barriers are health literacy and comprehension. For patients to effectively use the information in the subsequent steps of SDM, they must be able to receive and understand it, while physicians need strong communication skills to convey the information clearly [48]. The physician should present alternative treatment options, outlining the risks, benefits, and potential uncertainties, based on up-to-date, evidence-based, and validated data. [48] To improve the effectiveness of communication, physicians may use different techniques and tools enhancing health literacy. Educational aids can help patients make informed decisions [49]. These include: educational materials (brochures, graphics) and option grids, simplified information, no medical jargon, reduced amount of text in educational materials, presenting the most important information

first or in isolation, presenting figures logically, use of visual language, examples, and comparisons [23, 50, 51].

Step 3 Assess your patient's values and preferences

Understanding patients' values and preferences is a fundamental aspect of patient-centered care. However, eliciting this knowledge can be challenging since patients represent a heterogenic group in this regard. In addition, their values and preferences depend on many factors and may change in time [51]. At this step, both parties should seek congruency in the consultation—the clinical objectives should be in line with the patient's goals and values [51]. A recommended approach may turn out to be ineffective if the patient is not motivated to adhere to the treatment scheme or feels that the benefits do not justify the effort [52]. "Doctors might be experts in clinical matters, but the patient is the expert in their lived experience" [53]. To be able to express personal values and preferences, patients should be equipped with cognitive and social skills. Evidence indicates that these health-related skills can be enhanced through education [48].

Step 4 Reach a decision with your patient

In the 1990s, the practice of engaging patients in the decision-making process was criticized by many. Opponents of the SDM model argued that patients were often unwilling to participate in decision-making, that disclosing medical uncertainties could have negative consequences, that providing complete information on all treatment options was impractical, and that it might lead to an increase in demand for unnecessary and costly procedures [54]. Studies confirm that the paternalistic model can lead to satisfactory outcomes for some patients, while leaving others dissatisfied and with suboptimal results. However, over the past 20 years, there has been an increasing trend in patients preferring to be actively involved in the decision-making process regarding their health [52]. It is a common myth that most patients prefer their doctor to make decisions for them. While this may be true in some situations, researchers emphasize that it is not a reason to abandon SDM, but rather an

opportunity to take initiative. Patient's unwillingness to make a decision can be a starting point to a deeper discussion about the available options, recommendations, and patient's expectations. This attitude may benefit the patient-physician relationship, strengthening mutual respect and trust [42,52].

Step 5 Evaluate your patient's decision

Ensuring that the patient understands the key aspects of the decision, including its implications for their health, is a crucial final step in the model [42]. However, there is limited evidence in the literature on practical guidelines for assessing whether the patient has correctly understood the information necessary to make an informed health decision [55]. Nonetheless, there are tools to assess a patient's ability to make health decisions, which can help physicians evaluate the patient's situation in a structured manner [56]. However, the long-term evaluation process can be impeded by a lack of continuity in services and frequent staff rotations, which may result in the potential loss of information. The new staff may have no knowledge of the decision-making process from previous visits, which can undermine the patient-physician relationship [57].

The circumstances that limit the adoption of SDM should not be ignored. They include situations where the physician must balance individual wishes and preferences with the safety of the individual and the well-being of the wider population. For example, a patient may pressure the physician to prescribe antibiotics when it is not medically justified, or an individual addicted to psychoactive substances may demand opiates outside the recommended treatment regimen. Another situation arises when specific medical procedures or tests are mandated by the applicable law, such as newborn screening, or when medical evidence is insufficient to compare treatment options. Additionally, there are cases where life-saving actions must be taken or when a patient is deemed incapable of making decisions [58,59].

Based on the findings of Scholl et al. (2018), effectively addressing barriers to shared decision making (SDM) requires targeted actions at multiple levels. At the policy level, integrating SDM into healthcare regulations and providing financial incentives can promote its adoption. This includes embedding SDM principles into clinical guidelines and quality assessment frameworks to ensure systemic commitment. At the institutional level, fostering an organizational culture that values SDM is critical. This involves providing healthcare professionals with appropriate training, allocating sufficient time and resources for patient engagement, and embedding SDM tools into routine workflows and electronic health records. At the individual level, enhancing communication skills among both clinicians and patients, promoting patient empowerment and health literacy, and cultivating mutual respect and trust are necessary to facilitate effective collaborative decision making. Addressing barriers through this multi-level approach can help to mitigate structural, cultural, and interpersonal obstacles, thereby supporting more consistent and meaningful implementation of SDM in clinical practice.

The aim of this review was to update the current state of knowledge on the barriers and facilitators of SDM from the physician's and patient's perspectives in primary care. To mitigate the risk of imprecision and bias we used a specific methodology of article screening and data extraction. Nevertheless, the study has several limitations. First, since no precise definition of SDM exists, the authors decided to include studies where SDM matched the definition proposed by Makoul and Clayman [22]. The decision was based on its integrative nature and wide acceptance in the literature, allowing for consistent inclusion criteria across heterogeneous studies. Fundamental studies have included many socio-demographic factors of both healthcare providers and patient populations that may affect SDM, but only within a limited scope. Our search was limited to publications in English, potentially resulting in the omission of relevant studies published in other languages. The restriction to English-language publications may limit the generalizability of the findings, as relevant evidence published in other languages and cultural contexts could have been excluded. This may

particularly affect the identification of region-specific barriers and facilitators to SDM implementation. The search criteria may not have been comprehensive enough to capture all relevant studies reflecting both the physician's and the patient's perspectives. Additionally, our approach to data synthesis may have lacked sufficient depth, potentially failing to adequately assign barriers to the five steps of SDM. A more granular, step-by-step synthesis of barriers and facilitators—tailored to specific populations or healthcare systems—was not feasible due to the variability in study design, reporting standards, and the limited level of detail in some of the included sources. The study did not take into account barriers related to factors in the healthcare system (care settings), such as limitations in referral options for tests and specialist consultations, long waiting times for services outside the primary care system and others. This is a broad area that can affect patient involvement in decision-making. A more in-depth exploration of this issue may be a direction for future research. Finally, the conclusions of this review could be influenced by publication bias, as it is common for studies reporting only significant outcomes to be published.

Conclusions

Shared Decision Making (SDM) remains a challenge in primary healthcare but offers well-documented benefits, including improved treatment outcomes, patient satisfaction, and strengthened physician–patient communication. To facilitate its broader implementation, targeted actions are needed to eliminate persistent barriers.

Effective interventions include structured communication training for healthcare professionals, the use of simplified decision aids and educational materials, and promoting continuity of care to strengthen long-term clinical relationships. Programs such as MAGIC in the UK have demonstrated that a supportive institutional culture is essential for successful integration of SDM as a routine component of clinical practice. [19]

Future research should investigate the effectiveness of specific interventions across different healthcare settings, assess methods for integrating SDM into electronic medical records, and

monitor changes in barriers and facilitators over time. Emphasis should be placed on the role of digital and health literacy in enabling active patient participation. Such evidence will be critical for translating SDM principles into everyday practice in primary care.

Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and materials: All data generated or analysed during this study are included in this published article and its supplementary information files.

Competing interests: The authors declare that they have no competing interests

Funding: National Institute of Public Health NIH - National Research Institute: ZK1/2024

Authors' contributions: KDŽ: conceptualization of research, MKP: defining the main goal, KDŽ and MKP independently reviewed titles, abstracts and potentially relevant articles, KDŽ wrote a draft of the article; MKP verified a draft of the article; MKP: obtained financial resources. All authors read and approved the final manuscript.

References

1. Ratna H. The importance of effective communication in healthcare practice. Harvard Public Health Review. 2019;23.DOI:10.54111/0001/W4

2. Siebinga VY, Driever EM, Stiggelbout AM, Brand PLP. Shared decision making, patient-centered communication and patient satisfaction - A cross-sectional analysis. *Patient Educ Couns.* 2022;105(7):2145-2150. doi: 10.1016/j.pec.2022.03.012.
3. Expósito-Jiménez A, Alcaide-Leyva JM, Jiménez-Mérida MdR, Martínez-Angulo P. Health communication and shared decision-making between nurses and older adults in community setting: An integrative review. *Journal of Clinical Nursing.* 2024;33: 2922–2935. <https://doi.org/10.1111/jocn.17152>
4. US Preventive Services Task Force. Collaboration and Shared Decision-Making Between Patients and Clinicians in Preventive Health Care Decisions and US Preventive Services Task Force Recommendations. *JAMA.* 2022;327(12),1171–1176. doi:10.1001/jama.2022.3267
5. President's Commission for the Study of Ethical Problems in Medicine and Biomedical and Behavioral Research. Making health care decisions. The ethical and legal implications of informed consent in the patient— practitioner relationship. Washington DC 1982: https://repository.library.georgetown.edu/bitstream/handle/10822/559354/making_health_care_decisions.pdf
6. Ankolekar A, Dekker A, Fijten R, Berlanga A. The Benefits and Challenges of Using Patient Decision Aids to Support Shared Decision Making in Health Care. *JCO Clin Cancer Inform.* 2018; 2:1-10. doi: 10.1200/CCI.18.00013.
7. Agency for Healthcare Research and Quality (AHRQ). The SHARE Approach. 2018; <https://www.ahrq.gov/health-literacy/curriculum-tools/shareddecisionmaking/index.html>
8. Hargraves IG, Fournier AK, Montori VM, Bierman AS. Generalized shared decision making approaches and patient problems. Adapting AHRQ's SHARE Approach for Purposeful SDM. *Patient Educ Couns.* 2020;103(10):2192-2199. doi: 10.1016/j.pec.2020.

9. Ng CJ, Lee PY. Practising shared decision making in primary care. *Malays Fam Physician*. 2021; 22;16(1), 2-7. doi: 10.51866/cm0001.
10. Buljac-Samardzic M, Clark MA, van Exel NJA, van Wijngaarden JDH. Patients as team members: factors affecting involvement in treatment decisions from the perspective of patients with a chronic condition. *Health Expect*. 2022; 25; 138-148. <https://doi.org/10.1111/hex.13358>
11. Faiman B, Tariman J. Shared Decision Making: Improving Patient Outcomes by Understanding the Benefits of and Barriers to Effective Communication. *Clinical journal of oncology nursing*. 2019; 23: 540-542. 10.1188/19.CJON.540-542.
12. van der Horst DEM, Hofstra N, van Uden-Kraan CF, Stiggelbout AM et al. Shared Decision Making in Health Care Visits for CKD: Patients' Decisional Role Preferences and Experiences. *AJKD*. 2023; 20: doi: 10.1053/j.ajkd.2023.04.012
13. Matthias MS, Talib TL, Huffman MA. Managing chronic pain in an opioid crisis: what is the role of shared decision-making? *Health Commun* 2020; 1-9
14. Anagnostou A, Hourihane JOB, Greenhawt M. The role of shared decision making in pediatric food allergy management. *J Allergy Clin Immunol Pract*. 2020; 8: 46-51. [10.1016/j.jaip.2019.09.004](https://doi.org/10.1016/j.jaip.2019.09.004)
15. Sobczak K, Leoniuk K. Doctors' attitudes in the situation of delivering bad news: patients' experience and expectations. *Archives of Medical Science*. 2023;19(4):921-929. doi:10.5114/aoms/112756.
16. Hoque F. Shared Decision Making: Pinnacle for Patient-Clinician Relationships. *Journal of BMANA*. 2023; 2(1): 1-4.
17. Wieringa TH, Rodriguez-Gutierrez R, Spencer-Bonilla G, de Wit M, Ponce OJ, Sanchez-Herrera MF, Espinoza NR, Zisman-Ilani Y, Kunneman M, Schoonmade LJ, Montori VM, Snoek FJ. Decision aids that facilitate elements of shared decision making in chronic illnesses: a systematic review. *Syst Rev*. 2019;8,121. <https://doi.org/10.1186/s13643-019-1034-4>

18. Berger Z. Navigating the unknown: shared decision-making in the face of uncertainty. J Gen Intern Med. 2015; 15; 359:675-8.
19. Joseph-Williams N, Lloyd A, Edwards A, et al. Implementing shared decision making in the NHS: lessons from the MAGIC programme. BMJ 2017; 357, j1744. 10.1136/bmj.j1744
20. Montori VM, Ruissen MM, Hargraves IG, Brito JP, Kunneman M. Shared decision-making as a method of care. BMJ Evid Based Med. 2023; 28(4): 213-217. doi: 10.1136/bmjebm-2022-112068.
21. Moleman M, Regeer BJ, Schuitmaker-Warnaar TJ. Shared decision-making and the nuances of clinical work: Concepts, barriers and opportunities for a dynamic model. J Eval Clin Pract. 2021;27: 926–934. <https://doi.org/10.1111/jep.13507>
22. Makoul G, Clayman ML. An integrative model of shared decision making in medical encounter. Patient Educ. Couns. 2006; 60(3):301-312.
23. Harzheim L, Lorke M, Schulz S, Jünger S. Health literacy and shared decision-making in predictive medicine — professionals' perceptions and communication strategies. J Public Health (Berl.). 2023: <https://doi.org/10.1007/s10389-023-02110-0>
24. Schoefs E, Vermeire S, Janssens R, Huys I. HSD68 Barriers and Facilitators of Shared Decision Making in Clinical Practice: An Umbrella Review. Value in Health. 2022; 25(12): 287. <https://doi.org/10.1016/j.jval.2022.09.1411> NICE, 2021;
25. National Institute for Health and Care Excellence (NICE). (2021). Evidence review for effectiveness of approaches and activities to increase engagement in shared decision making and the barriers and facilitators to engagement: Shared decision making: Evidence Review. (NICE Guideline, No. 197.) <https://www.ncbi.nlm.nih.gov/books/NBK572433>
26. Eales OO, Smith S. Do socio-economically disadvantaged patients prefer shared decision-making? S Afr Fam Pract. 2021;17;63(1), e1-e6. doi: 10.4102/safp.v63i1.5293.

27. Caverly TJ, Hayward RA. Dealing with the Lack of Time for Detailed Shared Decision-making in Primary Care: Everyday Shared Decision-making. *J Gen Intern Med*. 2020 Oct;35(10):3045-3049.
28. Tan NQP, Maki KG, López-Olivo MA, Geng Y, Volk RJ. Cultural influences on shared decision-making among Asian Americans: A systematic review and meta-synthesis of qualitative studies. *Patient Educ Couns*. 2023;106:17-30. doi: 10.1016/j.pec.2022.10.350.
29. Creaney R. What Works to Support and Promote Shared Decision Making: A synthesis of recent evidence. Scottish Government 2019.
<https://www.gov.scot/binaries/content/documents/govscot/publications/research-and-analysis/2019/03/works-support-promote-shared-decision-making-synthesis-recent-evidence/documents/works-support-promote-shared-decision-making-synthesis-recent-evidence/wor> (accessed on 7 Jun 2024)
30. Grad R, Sandhu A, Majdpour D, Kitner S, Rodriguez C, Elwyn G. Perspectives of family physician educators on shared decision making in preventive health care: A Qualitative Descriptive Inquiry. *Patient Education and Counseling*. 2025; 134: 108681
31. Yahanda AT, Mozersky J. What's the Role of Time in Shared Decision Making? *AMA J Ethics*. 2020 May 1;22(5):E416-422. doi: 10.1001/amajethics.2020.416.
32. Hu Q, Feng Z, Zong Q, Wang J, Zheng Z, Feng D. Analysis of factors that promote the participation of patients with chronic diseases in shared decision making on medication: a cross-sectional survey in Hubei Province, China. *BMC Public Health* 2023; 23, 2440. <https://doi.org/10.1186/s12889-023-17099-0>
33. Bartels EA, Meijboom BR, Nahar-van Venrooij LMW, de Vries E. How service modularity can provide the flexibility to support person-centered care and shared decision-making. *BMC Health Serv Res*, 2021;18;21(1),1245. doi: 10.1186/s12913-021-07267-6. Erratum in: *BMC Health Serv Res*. 2022, 23;22(1),815.

34. Alsulamy N, Lee A, Thokala P, Alessa T. Views of stakeholders on factors influencing shared decision-making in the Eastern Mediterranean Region: a systematic review. *East Mediterr Health J.* 2021; 23;27(3),300-311. doi: 10.26719/emhj.20.139.
35. Tang C, Wang A, Yan J. Exploring motivations and resistances for implementing shared decision-making in clinical practice: a systematic review based on a structure–process–outcome model. *Health Expect.* 2022; 25: 1254-1268. doi:10.1111/hex.13541
36. Li X, Yang D, Meng M, Zhao J, Yin Y, Wang H, Zhang X, Liu Q, Li M, Liu J, Hao Y. Shared decision-making in healthcare in mainland China: a scoping review. *Front Public Health.* 2023; 7(11): 1162993. doi: 10.3389/fpubh.2023.1162993.
37. Légaré F, Härter M, Stiggelbout AM, Thomson R, Stacey D, North J.(2020). Choosing treatments and the role of shared decision-making. In: E. Nolte, S. Merkur, A. Anell, (Ed.). *Achieving Person-Centred Health Systems: Evidence, Strategies and Challenges* European Observatory on Health Systems and Policies. Cambridge University Press 2020; p. 283-316
38. Caverly TJ, Hayward RA. Dealing with the Lack of Time for Detailed Shared Decision-making in Primary Care: Everyday Shared Decision-making. *J Gen Intern Med.* 2020; 35(10),3045-3049. doi: 10.1007/s11606-020-06043-2.
39. Ousseine YM, Durand MA, Bouhnik AD, Smith A, Mancini J. Multiple health literacy dimensions are associated with physicians' efforts to achieve shared decision-making. *Patient Educ Couns.* 2019;102(11):1949-1956. doi: 10.1016/j.pec.2019.05.015.
40. Goto Y, Miura H, Son D, Scholl I, Kriston L, Härter M, Sato K, Kusaba T, Arai H. Association between physicians' and patients' perspectives of shared decision making in primary care settings in Japan: The impact of environmental factors. *PLoS One.* 2021;10;16(2), e0246518. doi: 10.1371/journal.pone.0246518
41. Hawley ST, Morris AM. Cultural challenges to engaging patients in shared decision making. *Patient Educ Couns.* 2017;100(1),18-24. doi: 10.1016/j.pec.2016.07.008.

42. Fisher KA, Tan ASL, Matlock DD, Saver B, Mazor KM, Pieterse AH. Keeping the patient in the center: Common challenges in the practice of shared decision making. *Patient Educ Couns*. 2018; 101(12): 2195-2201. doi: 10.1016/j.pec.2018.08.007.
43. Keij SM, de Boer JE, Stiggelbout AM, Bruine de Bruin W, Peters E, Moaddine S, Kunneman M, Pieterse AH. How are patient-related characteristics associated with shared decision-making about treatment? A scoping review of quantitative studies. *BMJ Open*. 2022; 24,12(5), e057293. doi: 10.1136/bmjopen-2021-057293.
44. Büdenbender B, Köther AK, Grüne B, Michel MS, Kriegmair MC, Alpers GW. When attitudes and beliefs get in the way of shared decision-making: A mediation analysis of participation preference. *Health Expect*. 2023; 26(2):740-751. doi: 10.1111/hex.13699.
45. Ajzen I, Schmidt P. Changing behavior using the theory of planned behavior. In: MS Hagger, LD Cameron, K Hamilton, N Hankonen, & T Lintunen, (Ed). *The Handbook of Behavior Change*. Cambridge University Press, 2020, p.17-31
46. Alsulamy N, Lee AP, Thokala P, Alessa T. What Influences the Implementation of Shared Decision Making: An Umbrella Review. *Patient Education and Counseling*. 2020; 103(12):2400-7. <https://doi.org/10.1016/j.pec.2020.08.009>.
47. Roodbeen R, Vreke A, Boland G, Rademakers J, van den Muijsenbergh M, Noordman J, van Dulmen S. Communication and shared decision-making with patients with limited health literacy; helpful strategies, barriers and suggestions for improvement reported by hospital-based palliative care providers. *PLoS One*. 2020;15(6), e0234926. doi: 10.1371/journal.pone.0234926.
48. Muscat DM, Shepherd HL, Nutbeam D, Trevena L, McCaffery KJ. Health Literacy and Shared Decision-making: Exploring the Relationship to Enable Meaningful Patient Engagement in Healthcare. *J Gen Intern Med*. 2021; 36(2): 521-524. doi: 10.1007/s11606-020-05912-0.
49. Stacey D, Légaré F, Lewis K, Barry MJ, Bennett CL, Eden KB, Holmes-Rovner M, Llewellyn-Thomas H, Lyddiatt A, Thomson R, Trevena L. Decision aids for people

facing health treatment or screening decisions. Cochrane Database Syst.Rev , 4,
CD001431.

50. Visscher BB, Steunenber B, Heijmans M, Hofstede JM, Deville W, van der Heide I, Rademakers J. Evidence on the effectiveness of health literacy interventions in the EU: a systematic review. BMC Public Health. 2018; 18(1).

51. Tringale M, Stephen G, Boylan AM, Heneghan C. Integrating patient values and preferences in healthcare: a systematic review of qualitative evidence. BMJ Open. 2022 18;12(11), e067268. doi: 10.1136/bmjopen-2022-067268.

52. Bialosky JE, Cleland JA, Mintken P, Beneciuk JM, Bishop MD. The healthcare buffet: preferences in the clinical decision-making process for patients with musculoskeletal pain. J Man Manip Ther. 2022; 30(2),68-77. doi: 10.1080/10669817.2021.1989754.

53. Thériault G, Grad R, Dickinson JA, Breault P, Singh H, Bell NR, Szafran O. To share or not to share: When is shared decision making the best option? Can Fam Physician. 2020; 66(5), 327-331.

54. Coulter A. Partnerships with patients: the pros and cons of shared clinical decision-making. J Health Serv Res Policy. 1997; 2(2): 112-21. doi: 10.1177/135581969700200209.

55. Cox CL. Patient understanding: how should it be defined and assessed in clinical practice? J Eval Clin Pract. 2023; 29: 1127-1134. [doi:10.1111/jep.13882](https://doi.org/10.1111/jep.13882)

56. Barstow C, Shahan B, Roberts M. Evaluating Medical Decision-Making Capacity in Practice. Am Fam Physician, 2018; 1,98(1):40-46.

57. Verwijmeren D, Grootens KP. Shifting Perspectives on the Challenges of Shared Decision Making in Mental Health Care. Community Ment Health J. 2024; 60:292–307. <https://doi.org/10.1007/s10597-023-01170-6>

58. Laight D. Risk communication: a pillar of shared decision making. Prescriber 2022; 33: 24-28. <https://doi.org/10.1002/psb.1993>

59. Elwyn G, Price A, Franco JVA, Gulbrandsen P. The limits of shared decision making. BMJ Evid Based Med. 2023; 28(4): 218-221. doi: 10.1136/bmjebm-2022-112089.

566 60. Scholl I, LaRussa A, Hahlweg P, Kobrin S, Elwyn G. Organizational- and system-level
567 characteristics that influence implementation of shared decision-making and
568 strategies to address them - a scoping review. Implement Sci. 2018 Mar 9;13(1):40.
569 doi: 10.1186/s13012-018-0731-z.

570

Preprint

SHARED DECISION MAKING IN PRIMARY CARE: BARRIERS AND FACILITATORS FROM THE PHYSICIAN'S AND PATIENT'S PERSPECTIVE — A SCOPING REVIEW

AIM

Integration of barriers, facilitators and perspectives of both clinicians and patients into the SHARE SDM 5-step model to provide a comprehensive understanding of its challenges and opportunities in primary care



METHODS

Scoping review (based on PRISMA-ScR)
Publications from last 5 years
Thematic data analysis



BARRIERS TO SDM

- Time constraints and physician's workload
- Lack of Shared Decision Making (SDM) culture and tradition
- Paternalistic model of healthcare
- Patients' passive attitude and belief they are unable to make decisions



FACILITATORS OF SDM

- Trust between physicians and patient
- Use of educational materials by physicians,
- Belief that SDM will lead to better treatment outcomes
- Physician communication training



CONCLUSIONS

Despite the challenges to public health, action should be taken to overcome barriers, given the well-established benefits of SDM, although this will require time and necessary adaptations.



Preprint

Table I Barriers and facilitators from the patient's and physician's perspectives in relation to 5 steps of SDM

BARRIERS		
	Patient	Physician
Step 1 Seek patient's participation	No SDM culture or tradition	No SDM culture or tradition
	Patient's passive attitude	Consultation involving SDM is viewed as burdensome
	Patient's anxiety related to consultations	
	General view of how a medical consultation should look like	Lack of time: Busy workload (too many patients per consultation)
	Socio-demographic factors (age, education level, gender): passive attitude is displayed by the elderly patients, males, and less educated individuals	
Step 2 Explore and compare treatment options	Being overwhelmed by the level of participation—"the less I know, the better"	Prioritizing the quantity of visits over their quality (patients' needs)
	Lack of explanation, incomplete information	Lack of initiative to provide information if not prompted
	Use of medical terminology, jargon—incomprehensible to the patient	
	Presenting only one treatment option where there are many to choose from	Presenting only one treatment option
	Physicians disregarding patients or becoming impatient when asked about the disease or treatment options	
Step 3 Assessment of patient's values and preferences	Hesitancy in expressing preferences	Disregarding patients' concerns and opinions
Step 4 Reaching a decision with the patient	Insufficient health literacy	Lack of communication or interpersonal skills
	Assuming that only doctors can make health decisions	Authority-based style of decision making
	Expecting that the visit should provide specific solutions	Assuming that patients prefer not to engage in the deliberation process and it is not needed
	Financial issues—only the fully or partially state-funded option is affordable	Unwillingness to engage patients in the decision-making process
	Uncertainty as to which option is best	
	Additional stress caused by the need to make a decision	
	Lack of empowerment	
	Difficulties in finding detailed health-related information, especially in the Internet	
	Not enough time to make a decision	

Step 5 Evaluate patient's decision		Physicians' fear of facing a higher liability risk
		No continuity in services—consultations are held by different specialists who give contradictory recommendations, hindering decision-making and causing confusion
		Staff rotation—attending physician changes
FACILITATORS		
	Patient	Physician
Step 1 Seek patient's participation	Trust—the foundation of a good relationship	Understanding that two experts participate in the clinical encounter
	Understanding that two experts participate in the clinical encounter	Physician's good communication and interpersonal skills
	Viewing healthcare as something that patients pay for, not something that is guaranteed, encourages patients to engage in SDM	Policy that supports clinicians—working time management and professional training (basic communication skills, patient engagement)
	Personal traits: kind, self-confident, empowered, proactive, ready to share decision-making	Believing that SDM leads to better treatment outcomes
Step 2 Explore and compare treatment options		Referring patients to credible evidence-based sources tailored to their specific needs
		Providing complete information in a comprehensible way
		Access to SDM facilitating tools, like patient care plans, educational materials
Step 3 Assessment of patient's values and preferences	Being aware of one's preferences	Care, empathy, calmness, openness expressed by the physician
Step 4 Reaching a decision with the patient	Self-education before the visit	Training for physicians
	Fostering empowerment in self-care	
	Health literacy—understanding options, uncertainty in medical evidence, and disease history	
	Being responsible for one's decisions and well-being	
	Viewing the physician as a decision-making partner	
	Involving family members in the treatment process	
	Having a nurse present during patient-physician consultations	
Step 5 Evaluate patient's decision		Physician maintains detailed case records, documents each visit, including concerns, patient preferences, and communication details

		Sharing responsibility—if responsibility for the decision is shared, physicians feel protected from potential negative outcomes
--	--	---

Preprint

