

Dissecting the potential impact of appendectomy on colorectal cancer through metabolites and inflammatory proteins: a Mendelian randomization study

Ming Qiu¹, Chongyuan Lan¹, Zhiyuan Xia¹, Hongtang Wu¹, Zizhuo Wang¹, Jiayi Wei¹, Minglin Lin^{1,2*}

¹The First Affiliated Hospital of Guangxi Medical University, China

²Guangxi Key Laboratory of Enhanced Recovery after Surgery for Gastrointestinal Cancer, China

Submitted: 17 September 2024; **Accepted:** 17 July 2025

Online publication: 6 September 2025

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms/208360>

Copyright © 2025 Termedia & Banach

***Corresponding author:**

Minglin Lin

The First Affiliated Hospital
of Guangxi Medical
University

Guangxi Key Laboratory
of Enhanced Recovery
after Surgery for
Gastrointestinal Cancer
China

E-mail: linminglin@sr.gxmu.edu.cn

Abstract

Introduction: Traditional longitudinal epidemiological investigations have demonstrated an association between appendectomy and the cumulative incidence of colorectal cancer (CRC). However, there is a paucity of evidence for a causal relationship between appendectomy and CRC from alternative methodologies.

Material and methods: We extracted summary statistics for CRC, appendectomy, 1,400 metabolites, and 91 inflammatory proteins from the largest European ancestry genome-wide association studies (GWAS) and the FinnGen study. Using two-sample bidirectional Mendelian randomization (MR), we explored the potential causal relationship between genetic liability to appendectomy and CRC. Moreover, we employed a two-step MR approach to identify metabolites and inflammatory proteins that may mediate the effects of genetic liability to appendectomy on CRC risk and analyzed their potential mediator variables. Furthermore, we used the significant single nucleotide polymorphisms (SNPs) associated with the risk of CRC due to appendectomy, searched for the nearest genes in the FinnGen, and explored the relationship between these genes and CRC prognosis by analyzing transcriptomic data from The Cancer Genome Atlas (TCGA) and Gene Expression Omnibus (GEO).

Results: Two-sample MR analysis revealed evidence supporting a potential causal effect of genetic liability to appendectomy on an increased risk of CRC. Two-step MR analysis identified carnitine and interleukin-13 (IL-13) as key mediators potentially involved in the positive causal relationship between genetic liability to appendectomy and CRC. Survival analysis showed that upregulation of HLX and OSR1 expression is positively correlated with poor prognosis in CRC patients.

Conclusions: This MR investigation provides evidence supporting a potential causal relationship between genetic liability to appendectomy and CRC, with carnitine and IL-13 potentially acting as partial mediators of this effect.

Key words: appendectomy, colorectal cancer, metabolites, inflammatory proteins, Mendelian randomization.

Introduction

Colorectal cancer (CRC) is one of the most common malignancies in the digestive system. According to global cancer statistics, CRC is the third most common cancer and the second leading cause of cancer-re-

lated deaths, with over 1.9 million new cases and approximately 935,000 deaths reported in 2020 [1]. Although the efficacy of CRC treatment has improved to some extent with surgery-based comprehensive treatment, the overall efficacy for CRC patients has not fundamentally improved. The treatment landscape for CRC has shifted significantly in recent years. Initially, the focus was on radical tumor resection combined with postoperative adjuvant radiotherapy and chemotherapy. This has gradually evolved into a comprehensive treatment model that includes surgical resection following neoadjuvant therapy or conversion therapy. Moreover, there has been a transition from traditional systemic chemotherapy to precision therapy, using biological targeted drugs or immunotherapy, based on molecular biomarkers. Furthermore, the emergence of multidisciplinary treatment models has significantly extended the survival of CRC patients and improved their quality of life [2].

Many factors influence the incidence of CRC and patient survival [3]. Research has shown that appendectomy can lead to gut microbiota dysbiosis and impaired gut barrier function, increasing the risk of CRC [4]. The appendix contains a high density of gut-associated lymphoid tissue, which plays an important role in gut immune regulation and microbial composition in synergy with the gut microbiome [5, 6]. The appendix is often considered a vestigial organ that can be removed without significant consequences. However, recent studies have found a rich presence of immune cells in the appendix, including various innate immune cells such as natural killer cells and intraepithelial CD8⁺ T cells [7]. Furthermore, although studies have shown that patients who have undergone appendectomy have a significantly reduced severity of colitis, transcriptomic analysis and animal experiments have suggested that the presence of the appendix may confer protective T cell immunity, potentially involving protection against tumor initiation rather than tumor progression [8]. Additionally, appendectomy has been closely associated with the occurrence of various diseases, including CRC, in the long term [9–12].

Mendelian randomization (MR) is a method that uses genetic variants, such as single nucleotide polymorphisms (SNPs), as instrumental variables (IVs) to assess potential causal relationships between exposures and outcomes [13]. Since associations between exposures and outcomes may be attributed to unmeasured confounders and reverse causality, using genetic determinants of exposure as IVs allows for the estimation of the causal component of these observed associations under specific assumptions. Additionally, IVs must satisfy three criteria: 1) they must be associated

with the exposure, 2) they must be independent of confounders of the exposure-outcome association, and 3) they must influence the outcome only through the exposure (i.e., satisfy the exclusion restriction assumption) [14]. The advantage of MR analysis lies in its ability to reduce bias from certain confounding factors, providing additional evidence for etiological research. This method has been widely used to explore potential causal relationships between biological or medical variables [15–17]. However, the use of MR to study the potential causal relationship between appendectomy and CRC has yet to be explored.

In this study, we used data from genome-wide association studies (GWAS) and public datasets to explore the causal relationship between appendectomy and CRC risk using two-sample bidirectional MR. We employed two-step MR to identify metabolites and inflammatory proteins that may mediate the impact of appendectomy on CRC risk. Additionally, using SNPs significantly associated with CRC risk following appendectomy, we identified the five nearest genes from the FinnGen study and employed transcriptomic analysis to further investigate the impact of these related genes on the survival prognosis of CRC patients.

Material and methods

Study design

To comprehensively explore the causal relationship between appendectomy and CRC risk and potential mediating effects, we conducted an in-depth MR analysis. This approach used summary statistics from published GWAS. The MR framework employs SNPs as instrumental variables to estimate the potential causal effect of an exposure on an outcome. The MR design adheres to three fundamental assumptions: (1) relevance assumption: SNPs are associated with the exposure; (2) independence assumption: SNPs are independent of confounding factors; and (3) exclusion restriction assumption: SNPs influence the outcome only through the exposure. The second and third assumptions are related to confounding and horizontal pleiotropy, respectively, and violations of these assumptions can be assessed using a series of statistical methods. Our study design is summarized in the Figure 1.

Data source

In this study, summary data on appendectomy were obtained from the FinnGen database (https://r10.finnngen.fi/pheno/K11_APPENDECTOMY), including 28,601 cases and 383,580 controls. The summary data for CRC were obtained from the study by Huyghe *et al.* [18], which included 19,948 CRC patients, 12,124 healthy individuals,

and 38,356,021 SNPs. These data are accessible at (<https://gwas.mrcieu.ac.uk/datasets/ebi-a-GCST012879/>). The summary data for 1,400 metabolites were sourced from the study by Chen *et al.* [19]. The data can be obtained from the GWAS Catalog database (GWAS Catalog) with accession numbers GCST90199621-GCST90201020. The summary data for 91 inflammatory proteins were derived from the study by Zhao *et al.* [20]. The GWAS Catalog accession numbers are GCST90274758-GCST90274848. Transcriptome sequencing data used to validate the nearest genes to SNPs and assess their associations with CRC patient prognosis were obtained from the TCGA database (<https://portal.gdc.cancer.gov/>) and the GEO database (<https://www.ncbi.nlm.nih.gov/geo/>) with data summarized by Marisa *et al.* [21].

Instrumental variable selection

In our MR analysis, we used SNPs as IVs to study the potential causal relationship between exposure and outcome at the genetic level. Significant SNPs for appendectomy, CRC, 1400 metabolites, and 91 inflammatory proteins were selected using a uniform criterion of $p < 5.0 \times 10^{-8}$. However, due to the limited availability of genome-wide significant SNPs for metabolites, we adopted a more lenient threshold for metabolites in the MR analysis, specifically $p < 1 \times 10^{-5}$ [22]. This adjustment allowed us to obtain a broader range of SNPs associated with metabolites and CRC.

Simultaneously, we used an r^2 threshold of < 0.001 and a clumping window of 10,000 kb to select independent SNPs and exclude those in linkage disequilibrium (LD) with lead SNPs [23]. This strategy allowed us to exclude SNPs that were correlated due to LD. Additionally, to minimize bias from weak instruments, we calculated the F -statistic for each SNP. SNPs with an F -statistic < 10 were considered weak instruments [24]. SNPs displaying palindromic alleles or ambiguous strand orientation were systematically excluded from the IVs used in the MR analysis [25].

MR analysis

We conducted a comprehensive two-sample bidirectional MR analysis using inverse variance weighted (IVW), MR-Egger, weighted median, weighted mode, and simple mode methods to explore the potential causal relationship between appendectomy and the outcome of CRC. IVW is one of the primary methods in MR studies, aggregating the Wald ratios of each SNP for a summary estimate [26]. The IVW method assumes that the genetic variants included in the analysis are valid instrumental variables. Due to its high statistical power and widespread use in MR analyses, the




Figure 1. Study design of bidirectional and mediated Mendelian randomization analyses
SNP – single nucleotide polymorphism.

IVW method was chosen as the main analytical approach, with the other four methods serving as supplementary analyses [27, 28]. When the direction of the causal relationship remains consistent across these five methods, it is considered to provide stronger evidence supporting a potential causal relationship. According to the results of the MR analysis, a p -value < 0.05 indicates statistically significant evidence for an association between the genetically predicted exposure and the outcome under the MR assumptions [29]. To assess the robustness and validity of the results, we conducted sensitivity analyses using Cochran's Q statistic to evaluate heterogeneity among the IVs, and Mendelian randomization pleiotropy residual sum and outlier (MR-PRESSO) to test for pleiotropy, with a p -value < 0.05 indicating significant heterogeneity and pleiotropy [30, 31]. The MR-PRESSO method can not only detect horizontal pleiotropic outliers but also identify and exclude outlier genetic variants and provide corrected causal effect estimates. Additionally, it evaluates whether there is a difference between the results before and after correction [32]. The "leave-one-out" approach is used to detect whether any instrumental variables disproportionately influence the causal effect estimate. This involves sequentially removing individual SNPs and calculating the meta-effect of the remaining SNPs. We assessed whether the results changed after removing each SNP to determine whether the outcome was influenced by any specific SNP [33]. Horizontal pleiotropy was assessed using the MR-Egger intercept, and if the p -value was > 0.05 , it was considered that there

was no statistically significant evidence of directional horizontal pleiotropy [34]. Furthermore, we used two-step MR analysis to identify potential mediating variables of metabolites and inflammatory proteins in the association between appendectomy and CRC. In the two-step MR analysis, we calculated the causal effect estimate of appendectomy on the mediator and the causal effect of the mediator on CRC. We then used the delta method to estimate the significance of the mediated effect and the proportion of the mediated effect in the total effect [35].

Transcriptomic analysis

Based on the SNPs identified in the MR analysis as being associated with the potential causal relationship between appendectomy and CRC, we further explored the association between the expression of the nearest genes, identified using the FinnGen database, and survival of CRC patients. Survival analysis was conducted using the “survival” package in R, based on transcriptomic data from TCGA and GEO (GSE39582). A p -value < 0.05 was considered statistically significant for differences in survival.

Results

MR analysis revealed a potential positive causal relationship between appendectomy and CRC

In the two-sample MR analysis, SNPs were used as IVs, with appendectomy as the exposure and CRC as the outcome. GWAS data for appendectomy were used as the summary association statistics for the exposure, and GWAS data for CRC were used as the summary association statistics for the outcome. Using the extracted 6 SNPs for appendectomy that met the selection criteria, we found evidence supporting a potential positive causal relationship between appendectomy and CRC risk (Table I). Since the IVW method has high statistical power, it was used as the main analysis method. The IVW analysis showed evidence sup-

porting a potential causal relationship between appendectomy and an increased risk of CRC (OR = 1.328, 95% CI: 1.073–1.643, $p = 0.009$). Although the results of MR-Egger (OR = 1.575, 95% CI: 0.782–3.171, $p = 0.272$), simple mode (OR = 1.365, 95% CI: 0.957–1.949, $p = 0.146$), and weighted mode (OR = 1.341, 95% CI: 0.972–1.849, $p = 0.133$) did not reach statistical significance, the effect estimates from the other MR methods were directionally consistent with those of the IVW analysis. As shown in the scatter plot and leave-one-out plot (Figure 2), no individual SNP appeared to disproportionately influence the overall causal effect estimate. Furthermore, heterogeneity assessment showed little evidence of heterogeneity in the association ($p_{\text{heterogeneity}} = 0.862$). Additionally, the MR-Egger intercept and MR-PRESSO analyses showed little evidence of horizontal pleiotropy ($p_{\text{pleiotropy}} = 0.642$, $p_{\text{presso}} = 0.909$). The reverse MR analysis (Supplementary Table S1) showed no evidence supporting a potential causal relationship between genetic susceptibility to CRC and appendectomy (OR = 0.957, 95% CI: 0.898–1.020, $p_{\text{IVW}} = 0.179$). This provides additional support for the robustness of the findings.

To conclude, our study results provide evidence supporting a potential positive causal relationship between appendectomy and the risk of CRC, suggesting that appendectomy may be associated with an increased risk of CRC.

Mediation analyses of potential metabolites and inflammatory proteins

To explore the potential mechanisms underlying the association between appendectomy and CRC, we conducted two-step MR analyses with 1,400 metabolites and 91 inflammatory proteins as potential mediators. First, in the MR analysis of 1,400 metabolites on CRC risk, based on $p_{\text{IVW}} < 0.05$, consistent OR values across five methods, and satisfying the requirements of the MR-PRESSO method [32] (excluding metabolites with fewer than 3 SNPs), we identified 82 metabolites associated with CRC risk (Supplementary Tables SII

Table I. Six genome-wide significantly related SNPs were used as IVs to study the potential causal relationship between appendectomy and colorectal cancer

SNP	Gene	EA	OA	EAF	Beta exposure	P exposure	Beta outcome	P outcome	F
rs11931959	PITX2	G	A	0.307581	0.112966	5.01E-34	0.0320035	0.0855185	148
rs2044674	PITX2	T	C	0.389854	0.055885	3.40E-10	0.0139629	0.455235	39
rs2484697	HLX	G	A	0.497524	0.0558568	1.43E-10	0.0187648	0.273563	41
rs2584229	NR2F2	C	T	0.17521	-0.0734796	2.40E-10	-0.0362832	0.112237	40
rs3905887	C4orf32	T	C	0.646271	0.0498745	4.91E-08	-0.00684343	0.704644	30
rs6707044	OSR1	G	A	0.217896	0.0725584	3.00E-12	0.0260994	0.247414	49

SNP – single nucleotide polymorphism, EA – effect allele, OA – other allele, EAF – effect allele frequency, F – F-statistic.




Figure 2. Mendelian randomization analyses showing evidence supporting a potential causal relationship between appendectomy and colorectal cancer. **A** – Scatter plots and causal effect estimates for five different methods. **B** – Leave-one-out plot of SNPs associated with appendectomy and colorectal cancer. **C** – IVW estimates of the association between appendectomy and colorectal cancer

SNP – single nucleotide polymorphism, IVW – inverse variance weight, OR – odds ratio, CI – confidence interval.

and SIII). Some evidence of heterogeneity in SNP effects was observed (Supplementary Table SIV), but the leave-one-out analysis did not indicate that the causal estimate was driven by specific SNPs (Supplementary Figure S1). Next, the MR-PRESSO method was used to test for horizontal pleiotropy among these 82 metabolites, and 14 metabolites were found to exhibit evidence of horizontal pleiotropy ($p_{\text{pleiotropy}} < 0.05$) (Supplementary Table SIII). After re-analyzing the metabolites showing evidence of horizontal pleiotropy using MR-PRESSO to exclude outlier SNPs and obtain corrected causal effect estimates, 5 metabolites had $p_{\text{IVW}} < 0.05$ and consistent OR values across all testing methods. In summary, 73 metabolites were identified as potentially causally associated with CRC risk (Figure 3). Subsequently, MR analysis was performed between appendectomy and the 73 metabolites, and the results showed that carnitine levels (GCST90199621) may partially mediate the association between appendectomy and CRC, with a proportion mediated of 4.47% (95% CI: –9.73–18.7%).

Similarly, using the same method, we analyzed the potential mediating roles of 91 inflammatory proteins in the association between appendectomy and CRC risk. We identified 18 inflammatory proteins associated with CRC risk (Supplementary Tables SV and SVI), with some SNPs showing heterogeneity (Supplementary Table SVII). However, the leave-one-out analysis indicated that no single SNP drove the causal estimate (Supplementary Figure S2). The MR-PRESSO test for horizontal pleiotropy identified 3 inflammatory proteins with evidence of horizontal pleiotropy ($p_{\text{pleiotropy}} < 0.001$) (Supplementary Table SVI). After re-analyzing these 3 proteins with MR-PRESSO to exclude outlier SNPs and obtain corrected causal effect estimates, 1 inflammatory protein had $p_{\text{IVW}} < 0.05$ with consistent OR values across all testing methods. Therefore, we identified 16 inflammatory proteins potentially causally associated with CRC risk (Figure 4). Performing MR analysis with these 16 inflammatory proteins as outcomes and appendectomy as the exposure, we found that interleukin-13 (IL-13, GCST90274799) may partially




Figure 3. Two-step Mendelian randomization analysis showing evidence supporting a potential causal relationship between metabolites and colorectal cancer, and between appendectomy and GCST90199621

SNP – single nucleotide polymorphism, IVW – inverse variance weight, OR – odds ratio, CI – confidence interval.




Figure 4. Two-step Mendelian randomization analysis showing evidence supporting a potential causal relationship between inflammatory proteins and colorectal cancer, and between appendectomy and GCST90274799

SNP – single nucleotide polymorphism, IVW – inverse variance weight, OR – odds ratio, CI – confidence interval.

mediate the association between appendectomy and CRC, with a proportion mediated of 12.8% (95% CI: –2.76–28.4%).

Taken together, our study results suggest that alterations in carnitine and IL-13 levels may partially mediate the association between appendectomy and CRC.

Appendectomy may influence CRC risk through genes located near the associated SNPs

To explore whether appendectomy may be associated with CRC risk at the genetic level, we identified the nearest genes to the 6 significant SNPs associated with the potential causal relationship between appendectomy and CRC risk using the FinnGen database. The genes identified were HLX, OSR1, PITX2, C4orf32, and NR2F2, indicating that these genes are located near SNPs associated with the potential relationship between appendectomy and CRC.

To further investigate the potential relevance of HLX, OSR1, PITX2, C4orf32, and NR2F2 in CRC, we conducted survival analyses using transcriptomic sequencing data from CRC patients in the TCGA and GEO databases. The results showed that high expression of HLX and OSR1 was positively correlated with poor prognosis in CRC patients in both TCGA and GSE39582 datasets (Supplementary Figure S3). In summary, these findings suggest that HLX and OSR1 may represent candidate genes warranting further investigation in the association between appendectomy and CRC, although the present analyses do not establish that appendectomy affects their

expression or influences CRC prognosis through these genes.

Discussion

In this MR study, we explored the potential causal relationship between appendectomy and the risk of developing CRC. By using genetic variants as IVs for appendectomy, our analysis provided evidence supporting a potential positive causal relationship between appendectomy and the risk of CRC. Additionally, we found that carnitine and IL-13 may act as potential mediators of this relationship.

Appendectomy, a common surgical procedure, is primarily used to treat acute appendicitis [36]. The appendix is rich in lymphoid tissue and is a major producer of IgA in the large intestine [37, 38]. Appendectomy may impair gut immunity, leading to decreased surveillance and reduced ability to identify and kill tumor cells. The appendix's rich biofilm and secluded anatomical location make it a reservoir for gut microbiota; its removal may alter the composition of the gut microbiota [39, 40], which may contribute to the development of CRC. Past studies have suggested a possible link between appendectomy and an increased risk of CRC [4], but the exact mechanisms of this relationship remain unclear. Our Mendelian randomization study provides further evidence supporting a potential positive causal relationship between appendectomy and an increased risk of CRC.

MR is an analytical method used to assess the potential causal relationship between exposures and clinically relevant outcomes [41]. As an emerging epidemiological method for causal re-

search, MR complements traditional epidemiological studies by helping to identify potential causal risk factors for diseases [42]. In our study, genetic variants were used as instrumental variables for appendectomy, which helped to reduce the impact of confounding factors and strengthened causal inferences. Through this approach, we found evidence supporting a significant positive potential causal relationship between appendectomy and the risk of CRC. This finding is consistent with some previous epidemiological studies that also reported an association between appendectomy and an increased risk of CRC [43–47].

Furthermore, our study identified carnitine and IL-13 as potential mediators in the association between appendectomy and CRC risk. These findings suggest that metabolic pathways and inflammatory responses may contribute to the association between appendectomy and CRC. Specifically, our two-step MR analysis suggested that carnitine may partially mediate this association. Additionally, IL-13 was identified as another potential mediator, suggesting that inflammatory pathways may also be involved.

Emerging evidence reveals a sophisticated immunomodulatory network involving vitamin D homeostasis, Th17 cell dynamics, the IL-31/IL-33 signaling axis, and gut microbial ecology in regulating immune-mediated inflammatory responses. Th17 cells – a proinflammatory CD4+ T-cell subset – are pivotal drivers of chronic inflammatory processes [48]. The alarmin cytokine IL-33 mediates dual activation of innate and adaptive immunity through ST2 receptor signaling, inducing Th2-polarized cytokine production (IL-5, IL-13) [49]. Murdaca *et al.* demonstrated that vitamin D insufficiency correlates with alterations in the IL-31/IL-33 signaling pathway, disrupting the Th1/Th17–Treg equilibrium and potentially facilitating bacterial translocation, thereby exacerbating intestinal immune dysfunction [50, 51]. Conversely, vitamin D sufficiency has been associated with a reduced risk of colorectal carcinogenesis in some clinical studies [52]. The gut microbiome's compositional diversity is fundamental to immune homeostasis, aligning with the hygiene hypothesis. Physiological microbial colonization orchestrates the differentiation of Th1/Th2/Th17/Treg lineages, while commensal-derived metabolites expand peripheral Treg populations to maintain mucosal immune tolerance [53]. Based on these findings, we hypothesize that appendectomy-related immunological alterations, including changes in microbial composition and immune signaling pathways, may contribute to the observed association between appendectomy and CRC risk. Surgical removal of the appendix may influence mucosal immune regulation, microbial ecology, and metabolic pathways, including carnitine metabolism,

although these mechanisms require further investigation.

Despite our study providing evidence supporting a potential causal relationship between appendectomy and an increased risk of CRC, there are some limitations to consider. First, MR studies rely on genetic variants as IVs, and their results may be affected by violations of MR assumptions, including genetic heterogeneity and pleiotropy. Second, our analysis is based on specific genetic variations and may not cover all relevant biological pathways. Therefore, future research should aim to further explore the mechanisms underlying this association and validate our findings, considering more genetic variations and biomarkers, as well as factors such as individuals' genetic backgrounds and lifestyles, to better reveal the complex interactions between appendectomy and CRC.

In conclusion, our study provides evidence supporting an association between appendectomy and CRC risk, and identifies metabolites and inflammatory proteins as potential mediators requiring further investigation. These findings may help improve prevention strategies and risk assessment methods for CRC, thereby reducing its incidence and related mortality.

Acknowledgments

We acknowledge the efforts of the GWAS and FinnGen study consortia for providing high quality resources for researchers.

Funding

This study was supported by the National Natural Science Foundation of China (82003072), and Innovation Project of Guangxi Graduate Education (No. YCSW2024258).

Ethical approval

Ethical approval and participant consent were obtained in the original studies. Patients or the public were not involved in the design, conduct, reporting, or dissemination plans of our research.

Conflict of interest

The authors declare no conflict of interest.

References

1. Sung H, Ferlay J, Siegel RL, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2021; 71: 209-49.
2. Zhang Q, Zhou Y, Song L, et al. China special issue on gastrointestinal tumors-Improved survival after multidisciplinary team decision for patients with advanced gastrointestinal cancer: a multicenter, noninterventional, controlled study. *Int J Cancer* 2023; 153: 1885-93.

3. Bożyk A, Krawczyk P, Reszka K, et al. Correlation between KRAS, NRAS and BRAF mutations and tumor localizations in patients with primary and metastatic colorectal cancer. *Arch Med Sci* 2022; 18: 1221-30.
4. Shi F, Liu G, Lin Y, et al. Altered gut microbiome composition by appendectomy contributes to colorectal cancer. *Oncogene* 2023; 42: 530-40.
5. Agrawal M, Allin KH, Mehandru S, Faith J, Jess T, Colombel JF. The appendix and ulcerative colitis – an unsolved connection. *Nature Rev Gastroenterol Hepatol* 2023; 20: 615-24.
6. Vitetta L, Chen J, Clarke S. The vermiform appendix: an immunological organ sustaining a microbiome inoculum. *Clin Sci* 2019; 133: 1-8.
7. Ishimoto Y, Tomiyama-Miyaji C, Watanabe H, et al. Age-dependent variation in the proportion and number of intestinal lymphocyte subsets, especially natural killer T cells, double-positive CD4+ CD8+ cells and B220+ T cells, in mice. *Immunology* 2004; 113: 371-7.
8. Collard MK, Tourneur-Marsille J, Uzzan M, et al. The appendix orchestrates T-cell mediated immunosurveillance in colitis-associated cancer. *Cell Mol Gastroenterol Hepatol* 2023; 15: 665-87.
9. Dai C, Jiang M, Huang YH. Appendectomy and future risk of microscopic colitis: correlation or causation? *Clin Gastroenterol Hepatol* 2023; 21: 1376-7.
10. Harnoy Y, Bouhnik Y, Gault N, et al. Effect of appendectomy on colonic inflammation and neoplasia in experimental ulcerative colitis. *J Br Surg* 2016; 103: 1530-8.
11. Lee KH, Lee S, Park JH, et al. Risk of hematologic malignant neoplasms from abdominopelvic computed tomographic radiation in patients who underwent appendectomy. *JAMA Surg* 2021; 156: 343-51.
12. Yang CY, Wu MC, Lin MC, Wei JCC. Risk of irritable bowel syndrome in patients who underwent appendectomy: a nationwide population-based cohort study. *EClinicalMedicine* 2020; 23. <https://doi.org/10.1016/j.eclinm.2020.100383>.
13. Emdin CA, Khera AV, Kathiresan S. Mendelian randomization. *JAMA* 2017; 318: 1925-6.
14. Glymour MM, Tchetgen Tchetgen EJ, Robins JM. Credible Mendelian randomization studies: approaches for evaluating the instrumental variable assumptions. *Am J Epidemiol* 2012; 175: 332-9.
15. Bouras E, Karhunen V, Gill D, et al. Circulating inflammatory cytokines and risk of five cancers: a Mendelian randomization analysis. *BMC Med* 2022; 20: 3.
16. Liu X, Tong X, Zou Y, et al. Mendelian randomization analyses support causal relationships between blood metabolites and the gut microbiome. *Nat Genet* 2022; 54: 52-61.
17. Sanna S, van Zuydam NR, Mahajan A, et al. Causal relationships among the gut microbiome, short-chain fatty acids and metabolic diseases. *Nat Genet* 2019; 51: 600-5.
18. Huyghe JR, Bien SA, Harrison TA, et al. Discovery of common and rare genetic risk variants for colorectal cancer. *Nat Genet* 2019; 51: 76-87.
19. Chen Y, Lu T, Pettersson-Kymmer U, et al. Genomic atlas of the plasma metabolome prioritizes metabolites implicated in human diseases. *Nat Genet* 2023; 55: 44-53.
20. Zhao JH, Stacey D, Eriksson N, et al. Genetics of circulating inflammatory proteins identifies drivers of immune-mediated disease risk and therapeutic targets. *Nat Immunol* 2023; 24: 1540-51.
21. Marisa L, de Reyniès A, Duval A, et al. Gene expression classification of colon cancer into molecular subtypes: characterization, validation, and prognostic value. *PLoS Med* 2013; 10: e1001453.
22. Yang J, Yan B, Zhao B, et al. Assessing the causal effects of human serum metabolites on 5 major psychiatric disorders. *Schizophr Bull* 2020; 46: 804-13.
23. Folkersen L, Gustafsson S, Wang Q, et al. Genomic and drug target evaluation of 90 cardiovascular proteins in 30,931 individuals. *Nat Metabol* 2020; 2: 1135-48.
24. Choi KW, Chen CY, Stein MB, et al. Assessment of bidirectional relationships between physical activity and depression among adults: a 2-sample Mendelian randomization study. *JAMA Psychiatry* 2019; 76: 399-408.
25. Chen L, Yang H, Li H, He C, Yang L, Lv G. Insights into modifiable risk factors of cholelithiasis: a Mendelian randomization study. *Hepatology* 2022; 75: 785-96.
26. Pierce BL, Burgess S. Efficient design for Mendelian randomization studies: subsample and 2-sample instrumental variable estimators. *Am J Epidemiol* 2013; 178: 1177-84.
27. Boehm FJ, Zhou X. Statistical methods for Mendelian randomization in genome-wide association studies: a review. *Comp Struct Biotechnol J* 2022; 20: 2338-51.
28. Liu Y, Si M, Qian Y, et al. Bidirectional Mendelian randomization analysis investigating the genetic association between primary breast cancer and colorectal cancer. *Front Immunol* 2024; 14: 1260941.
29. Hu X, Zhao J, Lin Z, et al. Mendelian randomization for causal inference accounting for pleiotropy and sample structure using genome-wide summary statistics. *Proc Natl Acad Sci* 2022; 119: e2106858119.
30. Bowden J, Holmes MV. Meta-analysis and Mendelian randomization: a review. *Res Synth Meth* 2019; 10: 486-96.
31. Morrison J, Knoblauch N, Marcus J H, Stephens M, He X. Mendelian randomization accounting for correlated and uncorrelated pleiotropic effects using genome-wide summary statistics. *Nat Genet* 2020; 52: 740-7.
32. Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet* 2018; 50: 693-8.
33. Feng Y, Liu X, Tan H. Causal association of peripheral immune cell counts and atrial fibrillation: a Mendelian randomization study. *Front Cardiovasc Med* 2023; 9: 1042938.
34. Bowden J, Del Greco MF, Minelli C, Davey Smith G, Sheehan NA, Thompson JR. Assessing the suitability of summary data for two-sample Mendelian randomization analyses using MR-Egger regression: the role of the I² statistic. *Int J Epidemiol* 2016; 45: 1961-74.
35. Carter AR, Sanderson E, Hammerton G, et al. Mendelian randomisation for mediation analysis: current methods and challenges for implementation. *Eur J Epidemiol* 2021; 36: 465-78.
36. Jalava K, Sallinen V, Lampela H, et al. Role of preoperative in-hospital delay on appendiceal perforation while awaiting appendectomy (PERFECT): a Nordic, pragmatic, open-label, multicentre, non-inferiority, randomised controlled trial. *Lancet* 2023; 402: 1552-61.
37. Masahata K, Umamoto E, Kayama H, et al. Generation of colonic IgA-secreting cells in the caecal patch. *Nat Commun* 2014; 5: 3704.
38. Mörbé UM, Jørgensen PB, Fenton TM, et al. Human gut-associated lymphoid tissues (GALT): diversity, structure, and function. *Mucosal Immunol* 2021; 14: 793-802.
39. Cai S, Fan Y, Zhang B, et al. Appendectomy is associated with alteration of human gut bacterial and fungal communities. *Front Microbiol* 2021; 12: 724980.

40. Kooij I, Sahami S, Meijer S, Buskens C, Te Velde A. The immunology of the vermiform appendix: a review of the literature. *Clin Exp Immunol* 2016; 186: 1-9.
41. Sekula P, Fabiola Del Greco M, Pattaro C, Köttgen A. Mendelian randomization as an approach to assess causality using observational data. *J Am Soc Nephrol* 2016; 27: 3253-65.
42. Luo J, Thomassen JQ, Bellenguez C, et al. Genetic associations between modifiable risk factors and Alzheimer disease. *JAMA Network Open* 2023; 6: e2313734.
43. Dahabra L, Tallat M, Abouyassine A, Deeb L. S134 Does appendectomy alter the risk of colorectal cancer? a large population-based study in the United States. *Am J Gastroenterol* 2021; 116: S58-9.
44. Lee S, Jang EJ, Jo J, Park SJ, Ryu HG. Long-term impacts of appendectomy associated with increased incidence of inflammatory bowel disease, infection, and colorectal cancer. *Int J Colorectal Dis* 2021; 36: 1643-52.
45. Shi F, Liu G, Lin Y, Guo C, Han J, She J. IDDF2021-ABS-0054 Appendectomy increases the risk of colorectal cancer through causing gut microbiota dysbiosis in a large population-based study. *Gut* 2021; 70: A17-9.
46. Stellingwerf ME, de Koning MA, Pinkney T, Bemelman WA, D'Haens GR, Buskens CJ. The risk of colectomy and colorectal cancer after appendectomy in patients with ulcerative colitis: a systematic review and meta-analysis. *J Crohn's Colitis* 2019; 13: 309-18.
47. Wu SC, Chen WTL, Muo CH, Ke TW, Fang CW, Sung FC. Association between appendectomy and subsequent colorectal cancer development: an Asian population study. *PLoS One* 2015; 10: e0118411.
48. Balan Y, Packirisamy RM, Mohanraj PS. High dietary salt intake activates inflammatory cascades via Th17 immune cells: impact on health and diseases. *Arch Med Sci* 2022; 18: 459-65.
49. Aggeletopoulou I, Tsounis E P, Triantos C. Molecular mechanisms underlying IL-33-mediated inflammation in inflammatory bowel disease. *Int J Mol Sci* 2023; 24: 623.
50. Murdaca G, Gangemi S, Greco M. The IL-33/IL-31 axis in allergic and immune-mediated diseases. *Int J Mol Sci* 2023; 24: 9227.
51. Murdaca G, Tagliafico L, Page E, Paladin F, Gangemi S. Gender differences in the interplay between vitamin D and microbiota in allergic and autoimmune diseases. *Biomedicines* 2024; 12: 1023.
52. Fu Q, Liao H, Li Z, Chen X, Zhang X, Di J. Preventive effects of 13 different drugs on colorectal cancer: a network meta-analysis. *Arch Med Sci* 2023; 19: 1428-45.
53. Murdaca G, Greco M, Borro M, Gangemi S. Hygiene hypothesis and autoimmune diseases: a narrative review of clinical evidences and mechanisms. *Autoimmun Rev* 2021; 20: 102845.