

Causal Associations between Sleep Apnea and Dementia: A Two-Sample Mendelian Randomization Study

Keywords

Mendelian Randomization, Dementia, Sleep Apnea, Causality, Genome-Wide Association Study

Abstract

Introduction

Observational studies suggest a link between sleep apnea and dementia, but causality and directionality are unclear. This study investigated bidirectional causal relationships between sleep apnea and various dementia types using two-sample Mendelian randomization (MR).

Material and methods

This study utilized summary-level data from genome-wide association studies (GWAS). Sleep apnea (including obstructive sleep apnea (OSA) and generalized sleep apnea) exposure data were from a European ancestry study. Dementia (general, Alzheimer's, vascular, etc.) outcome/exposure data were from the Finnish FinnGen consortium. The inverse variance weighted (IVW) method was used as the primary analysis, supplemented by multiple analyses (MR-Egger, weighted median, weighted mode). Robustness was assessed using several sensitivity analyses, including MR-PRESSO, Cochran's Q test, and leave-one-out analysis.

Results

Forward analysis, after MR-PRESSO outlier correction, showed OSA associated with reduced unspecified dementia risk (IVW, OR=0.830, 95%CI=0.700-0.970). Reverse analysis, post-outlier removal, linked general dementia with reduced OSA risk (IVW, OR=0.9399, 95%CI=0.9187-0.9615) and generalized sleep apnea risk (IVW, OR=0.9141, 95%CI=0.8863-0.9427). Alzheimer's dementia was also associated with reduced OSA and generalized sleep apnea risk. Sensitivity analyses did not reveal significant horizontal pleiotropy for the main findings, and heterogeneity was generally within acceptable limits.

Conclusions

This MR study reveals complex, bidirectional genetic associations between sleep apnea and dementia subtypes. These findings offer new genetic insights into the complex sleep apnea-dementia interplay, highlighting subtype-specific associations.

Causal Associations between Sleep Apnea and Dementia: A Two-Sample Mendelian Randomization Study

Running title: Sleep Apnea and Dementia: MR Study

Lirong Zhao^{1#}, Yamin Xu^{2#}, Lu Xia^{3*}, Yan Zhang^{4*}

¹Department of Neurology, Huadong Hospital Affiliated to Fudan University, Shanghai, 200040, China

²Day Care Chemotherapy, Huadong Hospital Affiliated to Fudan University, Shanghai, 200040, China

³Day Surgery Unit, Huadong Hospital Affiliated to Fudan University, 200040, Shanghai, China

⁴Department of Geriatrics, Huadong Hospital Affiliated to Fudan University, Shanghai, 200040, China

[#]These authors are co-first authors

***Corresponding author**

Lu Xia

Address: No. 221, Yan'an West Road, Jing'an Temple Subdistrict, Jing'an District, 200040, Shanghai, China.

Email: Huadongxialu@163.com

Tel: +86-18121220713

Yan Zhang

Email: Huadongzhangyan@163.com

Abstract

Introduction: Observational studies suggest a link between sleep apnea and dementia, but causality and directionality are unclear. This study investigated bidirectional causal relationships between sleep apnea and various dementia types using two-sample Mendelian randomization (MR).

Material and methods: This study utilized summary-level data from genome-wide association studies (GWAS). Sleep apnea (including obstructive sleep apnea (OSA) and generalized sleep apnea) exposure data were from a European ancestry study. Dementia (general, Alzheimer's, vascular, etc.) outcome/exposure data were from the Finnish FinnGen consortium. The inverse variance weighted (IVW) method was used as the primary analysis, supplemented by multiple analyses (MR-Egger, weighted median, weighted mode). Robustness was assessed using several sensitivity analyses, including MR-PRESSO, Cochran's Q test, and leave-one-out analysis.

Results: Forward analysis, after MR-PRESSO outlier correction, showed OSA associated with reduced unspecified dementia risk (IVW, OR=0.830, 95%CI=0.700-0.970). Reverse analysis, post-outlier removal, linked general dementia with reduced OSA risk (IVW, OR=0.9399, 95%CI=0.9187-0.9615) and generalized sleep apnea risk (IVW, OR=0.9141, 95%CI=0.8863-0.9427). Alzheimer's dementia was also associated with reduced OSA and generalized sleep apnea risk. Sensitivity analyses did not reveal significant horizontal pleiotropy for the main findings, and heterogeneity was generally within acceptable limits.

Conclusions: This MR study reveals complex, bidirectional genetic associations between sleep apnea and dementia subtypes. These findings offer new genetic insights into the complex sleep apnea-dementia interplay, highlighting subtype-specific

associations.

Keywords: Sleep Apnea; Dementia; Mendelian Randomization; Causality;
Genome-Wide Association Study

Preprint

Introduction

Dementia is a progressive syndrome characterized by cognitive decline beyond normal aging, affecting memory, thinking, and daily functioning [1, 2]. It poses a significant global health challenge, particularly in aging populations globally [3]. As the global population continues to age, the current global estimate of 50 million individuals with dementia is projected to triple by 2050, reaching 150 million, necessitating innovative approaches to address this burgeoning public health crisis [4]. Despite its impact, public awareness remains low, and research is limited [3]. Current pharmacological interventions provide only symptomatic relief, yet they are often accompanied by adverse effects that may pose significant challenges in clinical management, such as nausea, vomiting, insomnia and even seizures [5]. This highlights the urgent need for new diagnostic techniques and therapeutic strategies with fewer side effects.

Non-pharmacological therapies are increasingly important in delaying or preventing unhealthy aging [1].

Obstructive Sleep Apnea (OSA) is a common respiratory disorder characterized by recurrent upper airway collapse during sleep, leading to intermittent hypoxemia and sleep fragmentation [6]. Its prevalence ranges from 2-10% worldwide, with higher rates in middle-aged men (14-49%) [7, 8]. Risk factors include obesity, age, male gender, and smoking [8, 9]. OSA significantly impacts quality of life and is associated with increased risks of hypertension, cardiovascular diseases, stroke, diabetes, and heart failure [10-12]. Given its prevalence and health consequences, early recognition and treatment of OSA are key to prevent adverse outcomes.

Recent observational studies have increasingly highlighted a strong link between OSA and cognitive decline, with OSA being associated with an elevated risk of developing

dementia, including Alzheimer's disease (AD) and vascular dementia [13, 14]. Specifically, OSA has been associated with alterations in brain morphology, demonstrable deficits in memory, attention, and executive control, as well as higher risks of developing AD and even Parkinson's disease [14]. The prevalence of OSA is significantly higher in AD patients compared to cognitively healthy individuals [15]. Studies have investigated AD biomarkers in OSA patients, including cerebrospinal fluid, blood, neuroimaging, and nuclear medicine biomarkers. While continuous positive airway pressure (CPAP) treatment may alleviate some damage, it does not fully reverse it [16]. However, while observational studies have provided numerous clues regarding the association between sleep apnea and dementia, their findings are susceptible to various confounding factors and cannot definitively exclude the possibility of reverse causation. Therefore, elucidating whether a true causal relationship exists between sleep apnea and dementia, and determining the direction of this causality, is of critical clinical and scientific importance for developing effective dementia prevention strategies and improving cognitive outcomes in patients with sleep apnea.

Mendelian Randomization (MR) is an epidemiological method that utilizes genetic variants, randomly allocated during meiosis, as instrumental variables (IVs) to assess the causal effect of an exposure on an outcome [17]. To be more specific, MR employs genetic variants that influence the exposure (susceptibility to sleep apnea) as proxies to test whether the exposure has a causal impact on the outcome (dementia). Under the core assumptions that these genetic variants are strongly associated with the exposure, independent of confounding factors, and affect the outcome solely through the exposure, MR can mimic the design of a randomized controlled trial. As genetic variants are determined at conception and are generally not influenced by postnatal environmental

factors or disease status, MR analysis can substantially mitigate confounding bias and reverse causation, which often plague traditional observational studies, thereby providing more robust evidence for causal inference [18]. In recent years, MR studies have begun to explore causal relationships between various exposures and either dementia or sleep apnea; however, systematic investigations into the bidirectional causal links between sleep apnea (specifically differentiating OSA and generalized SA) and multiple dementia subtypes remain scarce.

Given this context, the present study aims to utilize publicly available summary-level data from large-scale genome-wide association studies (GWAS) within a two-sample MR framework. We will systematically investigate the potential bidirectional causal associations between sleep apnea and various types of dementia. We expect to unravel the complex causal network between these two common and impactful diseases, thereby providing novel genetic evidence to inform early dementia prevention and the long-term management of patients with sleep apnea, potentially identifying new avenues for intervention.

Materials and Methods

Study Design

This study employed a two-sample MR study design. This methodology relies on several assumptions, as presented in **Figure 1**. For causal estimates derived from MR studies to hold validity, three fundamental assumptions must be satisfied: (1) IVs are robustly linked with the exposure; (2) IVs exhibit no association with potential confounding factors; (3) IVs do not directly influence the outcome, except through the exposure [19]. As this study is based on publicly available GWAS summary data, no

further ethical approval was required. Our study is reported in accordance with the Strengthening the Reporting of Observational Studies in Epidemiology using MR (STROBE-MR) checklist [20].

Data Sources

GWAS summary statistics for sleep apnea (including OSA and generalized sleep apnea) were sourced from a large published GWAS [21], primarily based on individuals of European ancestry. This included 126,695 cases and 303,630 controls for OSA, and 152,031 cases and 278,027 controls for generalized sleep apnea. GWAS summary statistics for various types of dementia were sourced from the FinnGen consortium (Release R12), which predominantly includes individuals of Finnish ancestry. Specifically, the 'Dementia' phenotype (GWAS ID: F5_DEMENTIA), comprising 24,864 cases and 469,981 controls, was defined based on International Classification of Diseases 10th Revision (ICD-10) codes F00-F09 from hospital discharge records. This block (F00-F09) comprises a range of mental disorders with a demonstrable etiology in cerebral disease, brain injury, or other insult leading to cerebral dysfunction. Other specific dementia phenotypes (e.g., Unspecified dementia, Vascular dementia, with sample sizes detailed in **Supplementary Table 1**) were also derived from FinnGen R12, based on their respective ICD-10 classifications. Both forward MR (sleep apnea as exposure, dementia as outcome) and reverse MR (dementia as exposure, sleep apnea as outcome) analyses were conducted. Detailed descriptions of the specific GWAS summary samples used in this study are presented in **Supplementary Table 1**.

Instrumental Variable Selection

To ensure the accuracy and reliability of conclusions regarding causal relationships, a series of quality control measures were implemented to select the most appropriate

genetic instruments. Initially, SNPs that met the genome-wide significance threshold ($p < 5 \times 10^{-8}$) were used as IVs for sleep apnea exposures, and for Dementia, Unspecified Dementia, and Dementia in Alzheimer disease when these were considered exposures in the reverse MR [22]. For other dementia types as exposures, a threshold of $p < 5 \times 10^{-6}$ was applied. Only SNPs with a minor allele frequency (MAF) greater than 0.01 were retained. To minimize the impact of linkage disequilibrium (LD), SNPs were further filtered based on an LD threshold ($r^2 < 0.001$) and a genomic region size limit of 10,000kb [23]. The F-statistic, calculated as $R^2 \times (N-2)/(1-R^2)$ for each IV, was used to assess the instrumental strength, where R^2 is the proportion of phenotypic variance explained by each genetic variant in the exposure, and N is the sample size [24]. An F-statistic exceeding 10 was set as the threshold for robust IVs [25]. In instances where selected SNPs were absent in the outcome summary data, proxy SNPs with high LD association ($r^2 > 0.8$) were used as substitutes [26].

MR Analyses

We conducted two-sample MR analyses using the TwoSampleMR package in R (version 4.0.5). The methodologies employed included inverse variance weighted (IVW), MR-Egger regression, weighted median (WM), and weighted mode approaches. Our principal analysis was based on the IVW method [25]. MR-Egger assesses causality while allowing for the presence of directional pleiotropy [27]. Weighted median analysis was utilized to assess causality, assuming validity in at least half of the IVs [19]. Initially, causal estimates were computed using the fixed effects IVW method for each IV. Upon detection of significant heterogeneity ($P < 0.05$), the analysis was shifted to the random effects IVW method. To visualize the causal relationships and the influence

of each SNP on the outcome, forest, scatter, and funnel plots were generated. Multiple testing correction was applied using the false discovery rate (FDR) method, with associations deemed statistically significant at $PFDR < 0.05$.

Sensitivity and Pleiotropy Analysis

Pleiotropy detection was conducted via MR-Egger, where a significant intercept ($P < 0.05$) is indicative of pleiotropy [27]. Cochran's Q test was utilized to evaluate heterogeneity among IVs for each phenotypic exposure, with a P value above 0.05 signifying an absence of heterogeneity [28]. MR-PRESSO was employed to identify and eliminate outliers ($P < 0.05$) and to adjust for horizontal pleiotropy [29]. A leave-one-out analysis was performed to determine the influence of each SNP on the overall causality by excluding each SNP sequentially before conducting MR analyses on the remaining SNPs [30].

Result

Selection of IVs

In the forward analysis, when OSA and sleep apnea were exposures, we identified 78 and 112 Instrument Variables, respectively. The mean F-statistics for these Instrument Variables were 42.1 (OSA) and 40.41 (sleep apnea); minimum F-statistics were 29.75 (OSA) and 29.78 (sleep apnea); and maximum F-statistics were 380.34 (OSA) and 392.23 (sleep apnea), respectively. For 11 single nucleotide polymorphisms (SNP) not directly available in outcome datasets, proxy SNP were utilized where possible (e.g., rs370508348 by rs7594555). An additional 4 SNP were removed during data harmonization. Detailed data for IV selection in the forward analysis are provided in **Supplementary Table S2- S4**.

In the reverse analysis, the exposures included dementia due to parkinson's disease, dementia in alzheimer disease, dementia in other diseases classified elsewhere, dementia (general), frontotemporal dementia, unspecified dementia, and vascular dementia, with the number of instrument variables being 13, 17, 20, 31, 2, 8, and 18, respectively. For the 13 IVs for dementia due to parkinson's disease, the mean F-statistic was 24.14, the minimum F-statistic was 21.14, the maximum F-statistic was 35.66, and the R^2 value was 0.001442. For the remaining exposures, mean F-statistics ranged from 24.93 (dementia in other diseases classified elsewhere) to 220.72 (dementia in alzheimer disease). All minimum F-statistics exceeded 20 (thus satisfying the $F > 10$ criterion), with the highest maximum F-statistic being 2628.18 (dementia in alzheimer disease). when OSA or Sleep apnea served as outcomes, 9 SNP were unmatched in their summary datasets; proxy SNP were employed where possible (e.g., rs140887881 by rs145562973). Subsequently, 2 SNP were excluded for the OSA outcome and 5 SNP for the Sleep apnea outcome during harmonization. Detailed data for IV selection in the reverse analysis are provided in **Supplementary Table S2, S5 and S6**.

Mendelian Randomization Analysis

Forward analysis

In the initial forward analysis for the association between OSA (OSA) and unspecified dementia, the IVW method did not show a statistically significant association (OR=0.600, 95%CI=0.310-1.160, P=0.1275, **Table 1, Figure S1**). However, the weighted median method suggested a significant protective effect (OR=0.750, 95%CI=0.590-0.950, P=0.0178, **Table 1, Figure S1**). The MR-PRESSO analysis identified the presence of outliers for this association (**Table 2**). After manual removal outliers, the IVW analysis revealed a statistically significant association, indicating that

OSA was associated with a reduced risk of unspecified dementia (OR=0.830, 95%CI=0.700-0.970, P=0.021, FDR=0.021, **Table 1, Figure S2**). This finding was further supported by the weighted median method (OR=0.740, 95%CI=0.580-0.940, P=0.015, FDR=0.015, **Figure S2**) using the outlier-corrected dataset (**Table 1**). Detailed data for all forward MR analyses, including results before and after outlier removal for all methods, are provided in **Supplementary Table S7-S8**.

The MR-Egger method results showed no significant pleiotropy for the association between OSA and unspecified dementia after outlier removal (P for intercept = 0.767, Table 3). For the IVW analysis of OSA and unspecified dementia after outlier removal, the Q statistic did not suggest significant heterogeneity (P = 0.535, Table 3). Detailed sensitivity analysis data are provided in **Supplementary Table S9-S12**.

Reverse analysis

Reverse analysis initially revealed several significant associations. General dementia was associated with a reduced risk of OSA (IVW method: OR=0.9418, 95%CI=0.9107-0.9740, P=0.0005, FDR=0.0017, **Table 1, Figure S3**) and generalized sleep apnea (IVW method: OR=0.9412, 95%CI=0.9122-0.9711, P=0.0001, FDR=0.0007, **Table 1, Figure S4**). The MR-PRESSO analysis detected outliers for both associations (**Table 2**). After manual removal of these outliers, general dementia remained significantly associated with a reduced risk of OSA (IVW method: OR=0.9399, 95%CI=0.9187-0.9615, P<0.001, FDR<0.001, **Table 1, Figure S5**) and generalized sleep apnea (IVW method: OR=0.9141, 95%CI=0.8863-0.9427, P<0.001, FDR<0.001, **Table 1, Figure S6**).

Furthermore, unspecified dementia (IVW method: OR=0.9644, 95%CI=0.9334-0.9964, P=0.0293, **Table 1, Figure S7**), dementia due to Parkinson's disease (IVW method:

OR=0.9760, 95%CI=0.9546-0.9979, P=0.0315, **Table 1, Figure S8**). and dementia in Alzheimer's disease (IVW method: OR=0.9668, 95%CI=0.9514-0.9824, P<0.001, FDR=0.0003, **Table 1, Figure S9**) were associated with a reduced risk of OSA.

Dementia in Alzheimer's disease was also associated with a reduced risk of generalized sleep apnea (IVW method: OR=0.9680, 95%CI=0.9552-0.9809, P<0.001, FDR<0.001, **Table 1, Figure S10**).

Furthermore, vascular dementia was associated with a reduced risk of sleep apnea (IVW method: OR=0.9671, 95%CI= 0.9419 - 0.9931, P=0.0133 FDR=0.031, **Table 1**) and OSA (IVW method: OR=0.9668, 95%CI=0.9596 - 0.9845, P=0.0016, FDR=0.0044, **Table 1**) However, MR-PRESSO analysis identified outliers for this association, and after their manual removal, the association was no longer statistically significant. Detailed data for all reverse MR analyses, including results before and after outlier removal where applicable, and their corresponding FDR values, are provided in **Supplementary Table S13-S14**.

For the reverse causal associations that remained significant after outlier removal, such as general dementia with OSA (Cochran's Q P=0.154; MR-Egger intercept P=0.343) and generalized sleep apnea (Cochran's Q P=0.218; MR-Egger intercept P=0.701), sensitivity analyses indicated no significant heterogeneity and horizontal pleiotropy (**Table 3**). Other positive reverse associations (Unspecified dementia with OSA, Dementia in Alzheimer's disease with OSA, and Dementia due to Parkinson's disease with OSA) also showed no evidence of pleiotropy via the MR-Egger intercept, although some exhibited heterogeneity (**Table 3**). This observed heterogeneity is generally permissible given that the primary IVW method employed in this study utilized a random-effects model. Detailed sensitivity analysis data, including results for

associations before outlier removal, are provided in **Supplementary Table S15-S18**.

Discussion

This study employed a two-sample MR approach to systematically investigate potential causal relationships between sleep apnea and various dementia subtypes. Our principal findings suggest a causal protective effect of genetic predisposition to OSA on the risk of unspecified dementia. Intriguingly, our analyses also hinted at potential causal links from genetic predisposition to general dementia and dementia in AD towards a reduced risk of both OSA and generalized sleep apnea, though these latter findings warrant cautious interpretation. These results offer novel genetic perspectives on the complex interplay between sleep apnea and dementia, holding potential implications for disease prevention and management.

The observed causal association suggesting that OSA may reduce the risk of unspecified dementia (IVW OR=0.830) is a noteworthy phenomenon that warrants careful consideration. This finding appears to contradict a substantial body of observational research reporting OSA as an independent risk factor for dementia [31-33]. This discrepancy may be attributable to the inherent strength of MR in mitigating confounding, a common limitation in observational studies. Alternatively, it could stem from the heterogeneity within the "unspecified dementia" outcome. This category is broadly defined to include various conditions such as 'dementia NOS,' 'psychosis NOS,' and 'primary degenerative dementia NOS' (both presenile and senile), while excluding specific diagnoses like 'senile dementia with delirium' (F05.1) or 'senility NOS' (R54). Consequently, "unspecified dementia" might encompass early-stage, atypical, or etiologically unclear cognitive impairments whose underlying pathophysiology could

differ significantly from that of well-defined dementia subtypes.

One potential explanation is that the genetic predisposition to OSA identified by MR analysis might differ from the clinically diagnosed OSA phenotype with significant pathophysiological consequences. Individuals carrying certain OSA genetic risk alleles might experience mild, subclinical intermittent hypoxia exposure throughout their early life, particularly as many risk factors for OSA are genetically determined [34]. While chronic, severe intermittent hypoxia undoubtedly promotes neurodegeneration through oxidative stress, neuroinflammation, impaired cerebral autoregulation, and accelerated A β deposition [35], the existence of a "hypoxic preconditioning" or "mild stress adaptation" window is debatable [36]. Some studies suggest that mild intermittent hypoxia may induce neuroprotective preconditioning effects, mediated by hypoxia-inducible factor (HIF) and its downstream targets like vascular endothelial growth factor (VEGF) and erythropoietin [37], thereby enhancing neuronal tolerance to subsequent, more severe insults and promoting neurovascular unit remodeling. If genetically predisposed individuals to OSA experience a lifelong pattern of hypoxic exposure that includes such preconditioning effects, possibly mediated by *HIF* and *VEGF*, and if this adaptive response outweighs the direct damage from mild hypoxia, it might confer a degree of "relative protection" against certain "vulnerable" or slowly progressing types of unspecified dementia. However, this remains a highly speculative explanation, requiring more refined mechanistic studies and MR analyses stratified by OSA severity. Furthermore, undetected horizontal pleiotropy of the IVs cannot be entirely excluded, meaning these OSA-related genetic variants might influence the risk of unspecified dementia through pathways independent of OSA.

Regarding the unexpected MR finding that genetic liability to general dementia and

dementia in AD might be causally linked to a reduced risk of sleep apnea, careful consideration is warranted. This genetically-inferred directionality is counterintuitive, contrasting with clinical observations that often associate manifest dementia with increased sleep apnea prevalence due to factors like pharyngeal muscle weakness or later-stage medication effects [38-40]. While speculative, one potential biological avenue for such a genetically-driven protective effect could involve early, subtle dementia-related alterations in neurotransmitter systems (such as cholinergic pathways [41]) or sleep-wake regulation that paradoxically stabilize respiration before widespread neurodegeneration. However, this counterintuitive observation could also stem from other factors. Complex genetic pleiotropy, where dementia-associated variants independently influence pathways reducing sleep apnea risk, is a possibility. Methodological considerations, such as potential survival or selection biases in GWAS for late-onset diseases or residual confounding from uncharacterized horizontal pleiotropy, are also pertinent [29]. Furthermore, the lifelong effect of genetic liability captured by MR may not fully align with the multifaceted clinical picture of advanced dementia.

The careful selection of datasets for dementia in AD and PD in this study, emphasizing ICD codes distinct from isolated AD or PD, helps to more accurately capture the causal effect of the "dementia state" itself on sleep apnea. GWAS data for AD (ICD-10, G30) or PD (ICD-10, G20) might predominantly represent earlier disease stages, often without overt dementia or associated severe neuromuscular or central respiratory dysregulation. In contrast, analyses focusing on "dementia in AD (F5_ALZHDEMENT)" or "dementia in PD (PD_DEMENTIA_EX)" are more reflective of the impact when neurodegenerative processes have advanced sufficiently to cause a

dementia syndrome.

The strengths of this study lie in its use of the MR design, effectively controlling for confounding inherent in observational studies and allowing for the assessment of potential causal directions. We utilized summary data from large-scale GWAS, enhancing statistical power. Furthermore, analyzing multiple dementia subtypes and differentiating between OSA and generalized sleep apnea provided more nuanced insights. The application of various sensitivity analyses also bolstered the robustness of our findings. However, this study also has limitations. First, MR analysis relies on three core assumptions; while we conducted several sensitivity analyses to assess these (e.g., for horizontal pleiotropy), the potential influence of unknown pleiotropy cannot be entirely ruled out. Second, the GWAS data used were primarily from European and Finnish populations, which may limit the generalizability of our findings to other ethnic groups. Third, despite distinguishing between dementia subtypes, GWAS data may not fully capture the heterogeneity within these conditions, such as dementia severity or disease duration. Fourth, for some dementia subtypes (e.g., frontotemporal dementia), the limited number of cases might have resulted in insufficient statistical power and weaker IVs, leading to uncertain results. Fifth, given that MR analyses are grounded in genetic susceptibility and diverge from clinical observational studies, it is plausible that sleep apnea treatment could modulate dementia risk in clinical settings, though this requires further validation in future clinical studies. Finally, for several of the dementia-to-sleep-apnea associations, the observed odds ratios, while reaching statistical significance in some analyses, were modest. Such effect sizes warrant cautious interpretation as they might represent smaller true effects or be more influenced by subtle biases, underscoring the need for validation in significantly larger

samples.

Future research must prioritize replication of these findings, especially the unexpected association between dementia predisposition and reduced sleep apnea risk, in independent, large-scale, and diverse cohorts. Investigating potential methodological reasons for this counterintuitive finding should be a priority. Further studies should also aim to use more refined phenotypes and investigate potential mediating biological pathways if these causal links are substantiated. In addition, clarifying dementia subtypes may facilitate earlier risk prediction and may promote the development of subtype-specific therapeutic or precision prevention strategies.

In conclusion, this MR study provides suggestive genetic evidence supporting a causal protective effect of OSA predisposition on unspecified dementia risk. It also presents exploratory, and more tentative, genetic evidence suggesting that predisposition to general dementia and AD may be causally linked to a reduced risk of sleep apnea. While the former aligns with hypotheses like hypoxic preconditioning, the latter is unexpected and requires substantial further investigation to distinguish potential novel biological insights from methodological or complex genetic phenomena. These findings underscore the intricate, and potentially counterintuitive, causal dynamics between sleep apnea and dementia.

Acknowledgements

None

Authors' contributions

Lirong Zhao and Yamin Xu carried out the studies, participated in collecting data, and drafted the manuscript. Lu Xia and Yan Zhang performed the statistical analysis and participated in its design. Lirong Zhao and Lu Xia participated in acquisition, analysis, or interpretation of data and draft the manuscript. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not Applicable

Consent for publication

Not Applicable

Competing interests

None

Funding

None

Availability of data and materials

All data generated or analysed during this study are included in this published article

References

1. Varghese J, Goothy SS, Karunakaran B. Dementia: A silent killer which kills millions in its wake: An overview. *International journal of health sciences* 2022; 6831-6836.
2. Alexopoulos P, Soldatos R, Kontogianni E, et al. COVID-19 Crisis Effects on Caregiver Distress in Neurocognitive Disorder. *J Alzheimers Dis* 2021; 79: 459-466.
3. Sun Y. Alzheimer's Disease and Dementia, Under-Recognized Public Health Crisis in China. *Journal of Gerontology & Geriatric Research* 2014; 03.
4. WHO. Dementia World Health Organization, 2018 [Available from: <https://www.who.int/news-room/fact-sheets/detail/dementia> (accessed Sep 14, 2025).
5. Yiannopoulou KG, Papageorgiou SG. Current and Future Treatments in Alzheimer Disease: An Update. *J Cent Nerv Syst Dis* 2020; 12: 1179573520907397.
6. Ayas NT, Hirsch AA, Laher I, et al. New frontiers in obstructive sleep apnoea. *Clin Sci (Lond)* 2014; 127: 209-216.
7. Garvey JF, Pengo MF, Drakatos P, Kent BD. Epidemiological aspects of obstructive sleep apnea. *J Thorac Dis* 2015; 7: 920-929.
8. Lam JC, Sharma SK, Lam B. Obstructive sleep apnoea: definitions, epidemiology & natural history. *Indian J Med Res* 2010; 131: 165-170.
9. Zou Y, Pan M, guo h. The associations between leisure sedentary behaviors, physical activity and chronic respiratory diseases: a Mendelian randomization study. *Archives of Medical Science* 2025.
10. Gonzaga C, Bertolami A, Bertolami M, Amodeo C, Calhoun D. Obstructive sleep apnea, hypertension and cardiovascular diseases. *J Hum Hypertens* 2015; 29: 705-712.
11. Vijayan VK. Morbidities associated with obstructive sleep apnea. *Expert Rev*

Respir Med 2012; 6: 557-566.

12. Assallum H, Song TY, Aronow WS, Chandy D. Obstructive sleep apnoea and cardiovascular disease: a literature review. Archives of Medical Science 2021; 17: 1200-1212.
13. Shastri A, Bangar S, Holmes J. Obstructive sleep apnoea and dementia: is there a link? Int J Geriatr Psychiatry 2016; 31: 400-405.
14. Weihs A, Frenzel S, Grabe H. The Link Between Obstructive Sleep Apnoea and Neurodegeneration and Cognition. Current Sleep Medicine Reports 2021; 7.
15. Buratti L, Luzzi S, Petrelli C, et al. Obstructive Sleep Apnea Syndrome: An Emerging Risk Factor for Dementia. CNS Neurol Disord Drug Targets 2016; 15: 678-682.
16. Liguori C, Maestri M, Spanetta M, et al. Sleep-disordered breathing and the risk of Alzheimer's disease. Sleep Med Rev 2021; 55: 101375.
17. Emdin CA, Khera AV, Kathiresan S. Mendelian Randomization. Jama 2017; 318: 1925-1926.
18. Sleiman PM, Grant SF. Mendelian randomization in the era of genomewide association studies. Clin Chem 2010; 56: 723-728.
19. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted Median Estimator. Genet Epidemiol 2016; 40: 304-314.
20. Skrivankova VW, Richmond RC, Woolf BAR, et al. Strengthening the Reporting of Observational Studies in Epidemiology Using Mendelian Randomization: The STROBE-MR Statement. Jama 2021; 326: 1614-1621.
21. Verma A, Huffman JE, Rodriguez A, et al. Diversity and scale: Genetic architecture

- of 2068 traits in the VA Million Veteran Program. *Science* 2024; 385: eadj1182.
22. Huang SJ, Li R, Xu S, Liu Y, Li SH, Duan SZ. Assessment of bidirectional relationships between circulating cytokines and periodontitis: Insights from a mendelian randomization analysis. *Front Genet* 2023; 14: 1124638.
23. Chen H, Sun X, Zhuo C, et al. Cardiac Troponin I and Risk of Stroke: A Mendelian Randomization Study. *Int J Gen Med* 2022; 15: 1575-1582.
24. Palmer TM, Lawlor DA, Harbord RM, et al. Using multiple genetic variants as instrumental variables for modifiable risk factors. *Stat Methods Med Res* 2012; 21: 223-242.
25. Burgess S, Small DS, Thompson SG. A review of instrumental variable estimators for Mendelian randomization. *Stat Methods Med Res* 2017; 26: 2333-2355.
26. Wang X, Chen Z, Tian W, et al. Plasma homocysteine levels and risk of congestive heart failure or cardiomyopathy: A Mendelian randomization study. *Front Cardiovasc Med* 2023; 10: 1030257.
27. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol* 2015; 44: 512-525.
28. Hartwig FP, Davey Smith G, Bowden J. Robust inference in summary data Mendelian randomization via the zero modal pleiotropy assumption. *Int J Epidemiol* 2017; 46: 1985-1998.
29. Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet* 2018; 50: 693-698.
30. Papadimitriou N, Dimou N, Tsilidis KK, et al. Physical activity and risks of breast

and colorectal cancer: a Mendelian randomisation analysis. *Nat Commun* 2020; 11: 597.

31. Shieu MM, Dunietz GL, Paulson HL, Chervin RD, Braley TJ. The association between obstructive sleep apnea risk and cognitive disorders: a population-based study. *J Clin Sleep Med* 2022; 18: 1177-1185.

32. Legault J, Thompson C, Martineau-Dussault M, et al. Obstructive Sleep Apnea and Cognitive Decline: A Review of Potential Vulnerability and Protective Factors. *Brain Sci* 2021; 11.

33. Ercolano E, Bencivenga L, Palaia ME, et al. Intricate relationship between obstructive sleep apnea and dementia in older adults. *Geroscience* 2024; 46: 99-111.

34. Mukherjee S, Saxena R, Palmer LJ. The genetics of obstructive sleep apnoea. *Respirology* 2018; 23: 18-27.

35. Snyder B, Shell B, Cunningham JT, Cunningham RL. Chronic intermittent hypoxia induces oxidative stress and inflammation in brain regions associated with early-stage neurodegeneration. *Physiol Rep* 2017; 5.

36. Baillieul S, Chacaroun S, Doutreleau S, Detante O, Pépin JL, Verges S. Hypoxic conditioning and the central nervous system: A new therapeutic opportunity for brain and spinal cord injuries? *Exp Biol Med (Maywood)* 2017; 242: 1198-1206.

37. Brzecka A. Brain preconditioning and obstructive sleep apnea syndrome. *Acta Neurobiol Exp (Wars)* 2005; 65: 213-220.

38. Lee JH, Ryan J, Andreescu C, Aizenstein H, Lim HK. Brainstem morphological changes in Alzheimer's disease. *Neuroreport* 2015; 26: 411-415.

39. Seidel K, Mahlke J, Siswanto S, et al. The brainstem pathologies of Parkinson's disease and dementia with Lewy bodies. *Brain Pathol* 2015; 25: 121-135.

40. Cori JM, O'Donoghue FJ, Jordan AS. Sleeping tongue: current perspectives of

genioglossus control in healthy individuals and patients with obstructive sleep apnea.

Nat Sci Sleep 2018; 10: 169-179.

41. Chen ZR, Huang JB, Yang SL, Hong FF. Role of Cholinergic Signaling in Alzheimer's Disease. Molecules 2022; 27.

Preprint

Figure Legends

Figure 1. The flow diagram of the process in this Mendelian randomization analysis.

Supplementary Figure Legends

Figure S1. The association between Obstructive sleep apnea and Unspecified dementia is presented in (A) a forest plot, (B) a leave-one-out sensitivity analysis, (C) a scatter plot, and (D) a funnel plot.

Figure S2. The association between Obstructive sleep apnea and Unspecified dementia/after exclusion is presented in (A) a forest plot, (B) a leave-one-out sensitivity analysis, (C) a scatter plot, and (D) a funnel plot.

Figure S3. The association between General dementia and Obstructive sleep apnea is presented in (A) a forest plot, (B) a leave-one-out sensitivity analysis, (C) a scatter plot, and (D) a funnel plot.

Figure S4. The association between General dementia and generalized Sleep apnea is presented in (A) a forest plot, (B) a leave-one-out sensitivity analysis, (C) a scatter plot, and (D) a funnel plot.

Figure S5. The association between General dementia and Obstructive sleep apnea /after exclusion is presented in (A) a forest plot, (B) a leave-one-out sensitivity analysis, (C) a scatter plot, and (D) a funnel plot.

Figure S6. The association between B General dementia and generalized Sleep apnea /after exclusion is presented in (A) a forest plot, (B) a leave-one-out sensitivity analysis, (C) a scatter plot, and (D) a funnel plot.

Figure S7. The association between Unspecified dementia and Obstructive sleep

apnea is presented in (A) a forest plot, (B) a leave-one-out sensitivity analysis, (C) a scatter plot, and (D) a funnel plot.

Figure S8. The association between Dementia due to Parkinson's disease and Obstructive sleep apnea is presented in (A) a forest plot, (B) a leave-one-out sensitivity analysis, (C) a scatter plot, and (D) a funnel plot.

Figure S9. The association between Dementia in Alzheimer's disease and Obstructive sleep apnea is presented in (A) a forest plot, (B) a leave-one-out sensitivity analysis, (C) a scatter plot, and (D) a funnel plot.

Figure S10. The association between Dementia in Alzheimer's disease and Generalized Sleep apnea is presented in (A) a forest plot, (B) a leave-one-out sensitivity analysis, (C) a scatter plot, and (D) a funnel plot.

Supplementary Table Legends

Table S1. Detailed Information on GWAS Data

Table S2. Table S2. Summary of Mendelian Randomization Analyses and Instrument Variable Characteristics

Table S3. Comprehensive Instrument Variable (IV) Information for Forward Analysis

Table S4. Details of SNPs After Matching and Their Effects on Exposure and Outcomes for Forward Analysis

Table S5. Comprehensive Instrument Variable (IV) Information for Reverse Analysis

Table S6. Details of SNPs After Matching and Their Effects on Exposure and Outcomes for Reverse Analysis

Table S7. MR Analysis Results for Forward Analysis

Table S8. MR Analysis Results for Forward Analysis After Outlier Removal

Table S9. Heterogeneity and Pleiotropy Tests for Forward Analysis

Table S10. Heterogeneity and Pleiotropy Tests for Forward Analysis After Outlier Removal

Table S11. MR-PRESSO Results for Forward Analysis

Table S12. MR-PRESSO Results for Forward Analysis After Outlier Removal

Table S13. MR Analysis Results for Reverse Analysis

Table S14. MR Analysis Results for Reverse Analysis After Outlier Removal

Table S15. Heterogeneity and Pleiotropy Tests for Reverse Analysis

Table S16. Heterogeneity and Pleiotropy Tests for Reverse Analysis After Outlier Removal

Table S17. MR-PRESSO Results for Reverse Analysis

Table S18. MR-PRESSO Results for Reverse Analysis After Outlier Removal

Preprint




Table 1. Causal Associations.

Analysis direction	Outcome	Exposure	N.SN P	Method	OR (95% CI)	P-value	FDR
Forward Analysis (Before Outlier Removal)							
	Unspecified dementia	Obstructive sleep apnea	73	Inverse variance weighted	0.600 (0.310 - 1.160)	0.1275	0.4368
	Unspecified dementia	Obstructive sleep apnea	73	MR Egger	0.450 (0.060 - 3.250)	0.4275	0.7973
	Unspecified dementia	Obstructive sleep apnea	73	Weighted median	0.750 (0.590 - 0.950)	0.0178	0.2498
	Unspecified dementia	Obstructive sleep apnea	73	Weighted mode	0.690 (0.470 - 1.020)	0.0665	0.6311
Forward Analysis (After Outlier Removal)							
	Unspecified dementia	Obstructive sleep apnea	70	Inverse variance weighted	0.830 (0.700 - 0.970)	0.021	0.021
	Unspecified dementia	Obstructive sleep apnea	70	MR Egger	0.880 (0.550 - 1.420)	0.6132	0.6132
	Unspecified dementia	Obstructive sleep apnea	70	Weighted median	0.740 (0.580 - 0.940)	0.015	0.015
	Unspecified dementia	Obstructive sleep apnea	70	Weighted mode	0.680 (0.460 - 0.990)	0.0503	0.0503
Reverse Analysis (Before Outlier Removal)							

Sleep apnea	Vascular dementia	15	Inverse variance weighted	0.9671 (0.9419 - 0.9931)	0.0133	0.031
Sleep apnea	Vascular dementia	15	MR Egger	0.9271 (0.8920 - 0.9635)	0.002	0.0121
Sleep apnea	Vascular dementia	15	Weighted median	0.9539 (0.9346 - 0.9735)	<0.001	<0.001
Sleep apnea	Vascular dementia	15	Weighted mode	0.9495 (0.9322 - 0.9671)	1.00E-04	3.00E-04
Obstructive sleep apnea	Vascular dementia	15	Inverse variance weighted	0.9596 (0.9354 - 0.9845)	0.0016	4.40E-03
Obstructive sleep apnea	Vascular dementia	15	MR Egger	0.9183 (0.8855 - 0.9523)	5.00E-04	0.006
Obstructive sleep apnea	Vascular dementia	15	Weighted median	0.9479 (0.9291 - 0.9670)	<0.001	<0.001
Obstructive sleep apnea	Vascular dementia	15	Weighted mode	0.9440 (0.9251 - 0.9633)	1.00E-04	3.00E-04
Obstructive sleep apnea	Unspecified dementia	7	Inverse variance weighted	0.9644 (0.9334 - 0.9964)	0.0293	0.0552
Obstructive sleep apnea	Unspecified dementia	7	MR Egger	0.9441 (0.8939 - 0.9971)	0.0938	0.2078
Obstructive sleep apnea	Unspecified dementia	7	Weighted median	0.9553 (0.9405 - 0.9704)	<0.001	<0.001
Obstructive sleep apnea	Unspecified dementia	7	Weighted mode	0.9554 (0.9401 - 0.9709)	0.0014	0.0021
Sleep apnea	Dementia in Alzheimer disease	15	Inverse variance weighted	0.9680 (0.9552 - 0.9809)	<0.001	<0.001

Sleep apnea	Dementia in Alzheimer disease	15	MR Egger	0.9771 (0.9609 - 0.9937)	0.0181	0.0723
Sleep apnea	Dementia in Alzheimer disease	15	Weighted median	0.9700 (0.9586 - 0.9816)	<0.001	<0.001
Sleep apnea	Dementia in Alzheimer disease	15	Weighted mode	0.9700 (0.9594 - 0.9807)	0.0001	0.0003
Obstructive sleep apnea	Dementia in Alzheimer disease	16	Inverse variance weighted	0.9668 (0.9514 - 0.9824)	<0.001	0.0003
Obstructive sleep apnea	Dementia in Alzheimer disease	16	MR Egger	0.9737 (0.9527 - 0.9952)	0.0316	0.0947
Obstructive sleep apnea	Dementia in Alzheimer disease	16	Weighted median	0.9670 (0.9550 - 0.9792)	<0.001	<0.001
Obstructive sleep apnea	Dementia in Alzheimer disease	16	Weighted mode	0.9669 (0.9541 - 0.9799)	0.0002	0.0005
Obstructive sleep apnea	Dementia due to Parkinsons disease	13	Inverse variance weighted	0.9760 (0.9546 - 0.9979)	0.0315	0.0552
Obstructive sleep apnea	Dementia due to Parkinsons disease	13	MR Egger	0.9861 (0.9059 - 1.0734)	0.753	0.8214
Obstructive sleep apnea	Dementia due to Parkinsons disease	13	Weighted median	0.9869 (0.9678 - 1.0064)	0.1867	0.2489
Obstructive sleep apnea	Dementia due to Parkinsons disease	13	Weighted mode	0.9890 (0.9661 - 1.0125)	0.3755	0.4506
Sleep apnea	Dementia	27	Inverse variance weighted	0.9412 (0.9122 - 0.9711)	0.0001	0.0007
Sleep apnea	Dementia	27	MR Egger	0.9651 (0.9233 - 1.0089)	0.1293	0.2078

	Sleep apnea	Dementia	27	Weighted median	0.9470 (0.9237 - 0.9710)	<0.001	<0.001
	Sleep apnea	Dementia	27	Weighted mode	0.9510 (0.9284 - 0.9741)	0.0004	0.0007
	Obstructive sleep apnea	Dementia	28	Inverse variance weighted	0.9418 (0.9107 - 0.9740)	0.0005	0.0017
	Obstructive sleep apnea	Dementia	28	MR Egger	0.9622 (0.9157 - 1.0110)	0.1385	0.2078
	Obstructive sleep apnea	Dementia	28	Weighted median	0.9537 (0.9283 - 0.9798)	0.0006	0.0009
	Obstructive sleep apnea	Dementia	28	Weighted mode	0.9497 (0.9264 - 0.9735)	0.0004	0.0007
Reverse Analysis (After Outlier Removal)							
	Sleep apnea	Vascular dementia	14	Inverse variance weighted	1.0079 (0.9714 - 1.0457)	0.6773	0.9016
	Sleep apnea	Vascular dementia	14	MR Egger	0.9695 (0.8903 - 1.0557)	0.4894	0.4894
	Sleep apnea	Vascular dementia	14	Weighted median	1.0046 (0.9646 - 1.0464)	0.8239	0.8239
	Sleep apnea	Vascular dementia	14	Weighted mode	1.0059 (0.9456 - 1.0701)	0.8542	0.8542
	Obstructive sleep apnea	Vascular dementia	14	Inverse variance weighted	0.9977 (0.9620 - 1.0347)	0.9016	0.9016
	Obstructive sleep apnea	Vascular dementia	14	MR Egger	0.9479 (0.8708 - 1.0318)	0.2401	0.3202

Obstructive sleep apnea	Vascular dementia	14	Weighted median	0.9928 (0.9525 - 1.0349)	0.7348	0.8239
Obstructive sleep apnea	Vascular dementia	14	Weighted mode	0.9913 (0.9341 - 1.0521)	0.779	0.8542
Obstructive sleep apnea	Dementia	25	Inverse variance weighted	0.9399 (0.9187 - 0.9615)	<0.001	<0.001
Obstructive sleep apnea	Dementia	25	MR Egger	0.9512 (0.9200 - 0.9835)	0.0074	0.0257
Obstructive sleep apnea	Dementia	25	Weighted median	0.9536 (0.9288 - 0.9791)	4.00E-04	8.00E-04
Obstructive sleep apnea	Dementia	25	Weighted mode	0.9500 (0.9263 - 0.9744)	6.00E-04	0.0013
Sleep apnea	Dementia	23	Inverse variance weighted	0.9141 (0.8863 - 0.9427)	<0.001	<0.001
Sleep apnea	Dementia	23	MR Egger	0.9229 (0.8711 - 0.9779)	0.0129	0.0257
Sleep apnea	Dementia	23	Weighted median	0.9190 (0.8801 - 0.9596)	0.0001	0.0005
Sleep apnea	Dementia	23	Weighted mode	0.9180 (0.8800 - 0.9577)	0.0007	0.0013

Table 2. Heterogeneity Tests and Pleiotropy Tests for Instrumental Variables.

Analysis direction	Outcome	Exposure	Heterogeneity		Pleiotropy	
			Q statistic (IVW)	P value	MR-Egger Intercept	P value
Forward Analysis						
Before Outlier Removal	Unspecified dementia	Obstructive sleep apnea	1257.118	<0.001	0.011	0.762
After Outlier Removal	Unspecified dementia	Obstructive sleep apnea	67.326	0.535	-0.003	0.767
Reverse Analysis (Before Outlier Removal)						
	Sleep apnea	Vascular dementia	41.6064	1.00E-04	0.0166	0.0209
	Obstructive sleep apnea	Vascular dementia	34.5027	0.0017	0.0174	0.0127
	Obstructive sleep apnea	Unspecified dementia	30.9729	<0.001	0.013	0.3833
	Sleep apnea	Dementia in Alzheimer disease	20.1879	0.1243	-0.0042	0.123
	Obstructive sleep apnea	Dementia in Alzheimer disease	28.9605	0.0163	-0.0031	0.358
	Obstructive sleep apnea	Dementia due to Parkinsons disease	31.7179	0.0015	-0.0042	0.8088
	Sleep apnea	Dementia	73.5277	<0.001	-0.0053	0.1383
	Obstructive sleep apnea	Dementia	79.7211	<0.001	-0.0042	0.2625
Reverse Analysis (After Outlier Removal)						
	Obstructive sleep apnea	Vascular dementia	22.735	0.045	0.012	0.217
	Sleep apnea	Vascular dementia	26.473	0.015	0.009	0.341
	Obstructive sleep apnea	Dementia	30.979	0.154	-0.003	0.343
	Sleep apnea	Dementia	26.813	0.218	-0.001	0.701

Table 3. MR-PRESSO results

Analysis direction	Outcome	Exposure	Raw		Outlier corrected		Global <i>p</i>	outliers	Distortion <i>p</i>
			OR_CI	<i>P</i>	OR_CI	<i>P</i>			
Forward Analysis									
Before Outlier Removal	Unspecified dementia	Obstructive sleep apnea	0.5954 (0.3057 - 1.1598)	0.1318	0.8257 (0.703 - 0.9697)	0.0224	<0.001	3	<0.001
After Outlier Removal	Unspecified dementia	Obstructive sleep apnea	0.8257 (0.703 - 0.9697)	0.0224	NA (NA - NA)	NA	0.527	NA	NA
Reverse Analysis (Before Outlier Removal)									
	Sleep apnea	Vascular dementia	0.9671 (0.9419 - 0.9931)	2.67E-02	1.0079 (0.9714 - 1.0457)	0.6841	0.04	1	<0.001
	Obstructive sleep apnea	Vascular dementia	0.9596 (0.9354 - 0.9845)	0.0069	NA (NA - NA)	NA	0.058	NA	NA
	Obstructive sleep apnea	Unspecified dementia	0.9644 (0.9334 - 0.9964)	0.0722	NA (NA - NA)	NA	0.308	NA	NA
	Sleep apnea	Dementia in Alzheimer disease	0.968 (0.9552 - 0.9809)	0.0003	NA (NA - NA)	NA	0.295	NA	NA
	Obstructive sleep apnea	Dementia in Alzheimer disease	0.9668 (0.9514 - 0.9824)	0.0009	NA (NA - NA)	NA	0.109	NA	NA
	Obstructive sleep apnea	Dementia due to Parkinsons disease	0.976 (0.9546 - 0.9979)	0.0526	0.9924 (0.9808 - 1.004)	0.2238	0.003	1	<0.001
	Sleep apnea	Dementia	0.9412 (0.9122 - 0.9711)	0.0008	0.9141 (0.8863 - 0.9427)	0	0.003	4	0.072

Reverse Analysis (After Outlier Removal)	Obstructive sleep apnea	Dementia	0.9418 (0.9107 - 0.974)	0.0017	0.9399 (0.9187 - 0.9615)	0	<0.001	3	0.889
	Obstructive sleep apnea	Vascular dementia	0.9977 (0.962 - 1.0347)	0.9035	NA (NA - NA)	NA	0.041	NA	NA
	Sleep apnea	Vascular dementia	1.0079 (0.9714 - 1.0457)	0.6841	1.0251 (0.9897 - 1.0617)	0.1918	0.008	1	0.545
	Obstructive sleep apnea	Dementia	0.9399 (0.9187 - 0.9615)	0	NA (NA - NA)	NA	0.197	NA	NA
	Sleep apnea	Dementia	0.9141 (0.8863 - 0.9427)	0	NA (NA - NA)	NA	0.294	NA	NA

