

Association between childbirth history and multiple health outcomes

Lizhong Yin¹, Huiwen Hu¹, Xuting Shi², Jiali Xiong^{1*}

¹Changzhou Maternal and Child Health Care Hospital, Changzhou Medical Center of Nanjing Medical University, China

²The Second People's Hospital of Changzhou, the Third Affiliated Hospital of Nanjing Medical University, China

Submitted: 28 January 2026; **Accepted:** 4 June 2026

Online publication: 25 July 2026

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms/224184>

Copyright © 2026 Termedia & Banach

***Corresponding author:**

Jiali Xiong
Changzhou Maternal and
Child Health Care Hospital
Changzhou Medical Center
of Nanjing Medical
University, China
E-mail: 13861083577@163.
com

Abstract

Introduction: The aim of the study was to investigate the association between childbirth experience and multiple health outcomes, and to discuss the role of blood lipids.

Material and methods: The cross-sectional study used data from the National Health and Nutrition Examination Survey (NHANES) from 2007 to 2020. Childbirth history (e.g., parity or number of live births) was obtained from questionnaire data. Blood lipid parameters included triglycerides, total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), and high-density lipoprotein cholesterol (HDL-C). Remnant cholesterol (RC) was calculated based on standard lipid equations..

Results: A total of 3889 participants were included in this study. Compared with women without childbirth history, the prevalence of several conditions was significantly higher among women with childbirth history, including high blood pressure (40.77% vs. 19.57%, $p < 0.001$), diabetes (13.15% vs. 5.86%, $p < 0.001$), heart disease (7.75% vs. 1.69%, $p < 0.001$) and arthritis (35.22% vs. 17.57%, $p < 0.001$). Individuals with two diseases were the most common (29.1%), and overweight–high blood pressure–arthritis was the most important comorbidity pattern. After multivariable adjustment, childbirth history was associated with higher odds of seven medical conditions (heart disease, cancer, kidney stones, arthritis, thyroid problem, diabetes, and high blood pressure). Odds ratios (OR) ranged from 1.57 to 3.54 and 95% confidence intervals (CIs) from 1.15 to 6.97 ($p < 0.0001$), depending on the outcome. In addition, the level of RC was significantly higher among women with these conditions ($p < 0.001$). A nonlinear (L-shaped) association between RC and several outcomes, including heart disease, arthritis, cancer, diabetes, and high blood pressure, was observed. Mediation analysis suggested that blood RC explained 8.2%, 10.6%, 6.84%, 17.7%, and 12.5% of the association between childbirth history and these outcomes.

Conclusions: Childbirth history was associated with multiple health outcomes in women, and RC may serve as a potential biomarker and partial mediator in these associations.

Key words: childbirth, pregnancy, health outcome, lipid, cholesterol.

Introduction

Pregnancy and childbirth are key biological and psychosocial events in a woman's life that may influence short- and long-term health outcomes [1], including physiology, metabolism, and psychology [2]. Beyond

their reproductive function, these processes are increasingly recognized as being associated with long-term health patterns. Evidence from previous studies suggested that metabolic, hormonal, and vascular changes in pregnancy may be associated with long-term health risks in women. The most frequently reported conditions include hypertension, cardiovascular disease, and diabetes. Hypertensive disorders of pregnancy (HDP) were associated with an increased risk of cardiovascular disease, potentially reflecting shared underlying vascular mechanisms related to blood pressure regulation. Timely recognition and modification of the optimal BP range in peripartum and postpartum may contribute to improvement of their outcomes [3]. One systematic review showed that gestational diabetes mellitus (GDM) and HDP were associated with higher risk of type 2 diabetes mellitus (T2DM) and hypertension after pregnancy [4]. Nakimuli *et al.* also reported that over one-third of women with pre-eclampsia would develop hypertension at one year postpartum [5].

Previous studies have predominantly focused on the adverse effects of pregnancy complications on women's health, while the influence of normal pregnancy and childbirth experience on women's long-term health remains less well understood. Conversely, reproductive experiences have been

associated with physiological adaptations that confer certain health advantages. For instance, pregnancy might lower the risks of hormone-sensitive malignancies, such as breast⁶ and ovarian cancers [6, 7]. Juang's group also found that pregnancy had an impact on primary dysmenorrhea [8]. Nevertheless, research in this field remains limited. Generally, pregnant studies to date have predominantly employed single-disease paradigms, which fail to reflect the clinical reality of the women's short- and long-term health.

Evidence suggests that pregnancy is associated with significant changes in lipid metabolism during the peripartum and postpartum periods. These changes often affect women's health [9, 10]. Recently, remnant cholesterol (RC), defined as cholesterol other than low-density lipoprotein cholesterol (LDL-C) and high-density lipoprotein cholesterol (HDL-C), has been identified as a potential marker of perinatal diseases, especially GDM [11, 12]. Gao *et al.* reported RC, but not other cholesterol parameters, may serve as an early indicator of GDM at 12–14 weeks of gestation. Elevated RC levels (> 24.30 mg/dl) were associated with a higher risk of developing GDM [13]. In addition, remnant cholesterol was found to have a positive association with blood pressure and sustained hypertension in pregnancy and 6 and 9 years after pregnancy, and the effect was more significant than that of total cholesterol (TC), LDL-C and HDL-C [14]. However, the longitudinal trajectory of RC in women with typical reproductive histories and its influence on women's health are still unclear.

In order to explore the association between childbirth experience and multiple health outcomes, we investigated seven cycles of the National Health and Nutrition Examination Survey (NHANES, 2007–2020) [15, 16]. We first examined these relations. The selected health outcomes included 16 chronic diseases and 26 disease subtypes, such as diabetes, heart disease, cancer, liver condition, failing kidney, kidney stones, thyroid problem, high blood pressure, sleep disorders, and arthritis. We further evaluated the association between blood lipid parameters, particularly cholesterol-related indices, and these outcomes. This study provides evidence on potential associations between childbirth experience and multiple health outcomes, which may have implications for future research on women's health.

Material and methods

Data source and study population

The cross-sectional study was from seven cycles of NHANES (2007–2020). Selection of participants is illustrated in Figure 1. Of 66,148 initial subjects, 32,793 were excluded due to male gender. 16,627




Figure 1. Flowchart of participant selection from the NHANES dataset in the present study

were excluded due to incomplete data of age, education level, race, poverty income ratio, and body mass index (BMI). 5468 were excluded due to missing the data of blood lipid indices. 7371 were excluded due to missing the information of pregnancy, childbirth, and multiple health outcomes. Finally, 3889 women were included. Among excluded female participants, 11,847 reported no pregnancy history but were excluded due to incomplete data on blood lipid indices or health outcomes. To assess potential selection bias in this specific subgroup, we compared baseline characteristics between these excluded nulligravid women and included nulligravid women ($n = 649$). No significant differences were observed in age (36.8 ± 18.2 vs. 37.2 ± 17.7 years, $p = 0.62$), racial composition, or education level (college graduate: 40.2% vs. 41.5%), suggesting limited evidence of selection bias in the nulligravid subgroup. Additionally, no substantial differences were observed in the measured baseline characteristics between the overall excluded female participants and the included participants, suggesting that selection bias due to these measured variables may be limited.

Definition of pregnancy and childbirth history

Pregnancy history was measured by the following question: “Ever been pregnant? (RHQ131)”. Pregnancy history included current pregnancy, live births, miscarriages, stillbirths, tubal pregnancies, and abortions. Participants who answered “No” were considered as having no pregnancy history (G1). Similarly, childbirth history and birth number were defined according to the question: “How many live births have you had? (RHQ171)”. Women who reported one or more live births were classified as having a history of childbirth (G2). Age at first birth (AFB) and age at last birth (ALB) were defined using the questionnaire items “Age at first live birth (RHQ180)” and “Age at last live birth (RHD190)”, respectively. The interval between the first and last live birth was calculated as ALB minus AFB.

Definition of multiple health outcomes

As in a previous study [17], multiple health outcomes were assessed using self-reported NHANES questionnaire data, primarily from the Medical Conditions Questionnaire (MCQ), Diabetes Questionnaire (DIQ), Kidney Conditions Questionnaire (KIQ), Sleep Disorders Questionnaire (SLQ), and Blood Pressure Questionnaire (BPQ). Women were considered as having the condition when they answered “Yes” to the question “Has a doctor or other health professional ever told you that you have?”

In this study, 16 conditions and 26 subtypes were included: heart disease (heart attack, angio-

na pectoris, coronary heart disease, congestive heart failure), stroke, emphysema, asthma, chronic bronchitis, liver condition, failing kidney, kidney stones, cancer (breast, cervix, ovary, uterus), overweight, arthritis (osteoarthritis, rheumatoid arthritis), gout, thyroid problem, diabetes, high blood pressure, and sleep disorder.

Measurements of blood lipid indices

Women’s blood lipid indices in questionnaire were obtained from laboratory data, including triglyceride (mmol/l), total cholesterol (TC, mmol/l), LDL-cholesterol (LDL-C, mmol/l) and HDL-cholesterol (HDL-C, mmol/l). According to the study [18], remnant cholesterol (RC) was calculated as deducting the levels of LDL-C and HDL-C from TC. According to the literature [19], we divided RC into four quartile levels: Q1 (< 0.43 mmol/l), Q2 ($0.41\text{--}0.61$ mmol/l), Q3 ($0.61\text{--}0.88$ mmol/l) and Q4 (≥ 0.88 mmol/l). Meanwhile, Q1 was defined as the reference in the follow-up study.

Other covariates

Similar to some other studies [20–22], age, race, education level, and poverty income ratio were obtained from demographics data. Gender was not included as a covariate since only female participants were enrolled.

Statistical analysis

DecisionLinnc1.0 software [23], a platform that integrates multiple programming language environments, was employed for data analysis. A logistic regression analysis model was employed to analyze two distinct models to examine the relationship between childbirth experience and multiple health outcomes. Next, restricted cubic splines (RCS) was used to assess potential nonlinear relationships between blood lipid indices and the risk of multiple health outcomes associated with childbirth experience. Parallel mediation analysis was carried out to clarify whether blood RC plays a potential intermediary role between childbirth and multiple health outcomes. $P < 0.05$ was considered statistically significant.

Results

Baseline participant characteristics

A total of 3889 participants with complete data from NHANES (2007–2020) were included in this study. Table I presents the baseline characteristics of the participants according to childbirth history. Compared to women who had never been pregnant (G1), those with a history of childbirth showed higher age and BMI, but lower education level and poverty income ratio ($p < 0.001$). At the

Table I. Baseline participant characteristics according to childbirth history

Variable	Overall	Childbirth history		P-value
		G1 (no)	G2 (yes)	
N	3889	649	3240	
Age [years]	49.879 ±17.817	37.228 ±17.699	52.413 ±16.731	< 0.001
Race (%)				
Mexican American	558 (14.35)	66 (10.17)	492 (15.19)	< 0.001
Other Hispanic	392 (10.08)	60 (9.24)	332 (10.25)	
Non-Hispanic White	1835 (47.18)	312 (48.07)	1523 (47.01)	
Non-Hispanic Black	762 (19.59)	115 (17.72)	647 (19.97)	
Other race	342 (8.79)	96 (14.79)	246 (7.59)	
Poverty income ratio	2.416 ±1.618	2.664 ±1.669	2.366 ±1.603	< 0.001
< 5.0	3272 (84.13)	516 (79.51)	2756 (85.06)	
≥ 5.0	617 (15.87)	133 (20.49)	484 (14.94)	
Education level (%)				< 0.001
Less than 9 th grade	347 (8.92)	21 (3.24)	326 (10.06)	
9–11 th grade	586 (15.07)	36 (5.55)	550 (16.98)	
High school graduate	822 (21.14)	92 (14.18)	730 (22.53)	
Some college	1199 (30.83)	231 (35.59)	968 (29.88)	
College graduate or above	935 (24.04)	269 (41.45)	666 (20.56)	
BMI [kg/m ²]	29.251 ±7.461	27.826 ±8.688	29.536 ±7.158	< 0.001
Normal weight (%)	1229 (31.60)	307 (47.30)	922 (28.46)	
Overweight (%)	1529 (39.32)	191 (29.43)	1338 (41.30)	
Obesity (%)	1131 (29.08)	151 (23.27)	980 (30.25)	
Triglycerides	1.289 ±0.689	1.091 ±0.583	1.328 ±0.702	< 0.001
Total cholesterol	5.067 ±1.044	4.889 ±0.979	5.103 ±1.053	< 0.001
LDL cholesterol	2.962 ±0.905	2.831 ±0.848	2.988 ±0.914	< 0.001
HDL cholesterol	1.515 ±0.407	1.558 ±0.397	1.506 ±0.409	0.003
Remnant cholesterol	0.59 ±0.316	0.5 ±0.267	0.609 ±0.322	< 0.001
Q1	1402 (36.05)	306 (47.15)	1096 (33.83)	
Q2	1022 (26.28)	175 (26.96)	847 (26.14)	
Q3	854 (21.96)	110 (16.95)	744 (22.96)	
Q4	611 (15.71)	58 (8.94)	553 (17.07)	

same time, they also showed abnormal levels of several blood lipid indices, which were characterized by significantly lower HDL-C and significantly higher values of other indices (triglyceride, TC, LDL-C and RC).

The occurrence of multiple health outcomes

Table II compares the occurrence of multiple health outcomes according to childbirth history (Supplementary Figure S1). The incidence of most chronic diseases among women who had given birth was significantly higher ($p < 0.05$), except asthma, chronic bronchitis, and liver condition.

It was particularly noteworthy that the incidence of high blood pressure (40.77% vs. 19.57%, $p < 0.001$), diabetes (13.15% vs. 5.86%, $p < 0.001$), heart disease (7.75% vs. 1.69%, $p < 0.001$), and arthritis (35.22% vs. 17.57%, $p < 0.001$) was significantly higher. The incidence of four common subtypes of cardiovascular diseases (congestive heart failure, coronary heart disease, angina pectoris, heart attack) was significantly higher. Interestingly, the incidence of uterus, cervix, and breast cancer in women with childbirth experience was significantly higher ($p < 0.001$), but the incidence of ovarian cancer was not significantly different.

Table II. Multiple health outcomes according to childbirth history [n (%)]

Variable	Overall	Childbirth history		P-value
		G1 (no)	G2 (yes)	
Heart disease				< 0.001
No	3627 (93.26)	638 (98.31)	2989 (92.25)	
Yes	262 (6.74)	11 (1.69)	251 (7.75)	
Congestive heart failure				< 0.001
No	3771 (96.97)	645 (99.38)	3126 (96.48)	
Yes	118 (3.03)	4 (0.62)	114 (3.52)	
Coronary heart disease				
No	3786 (97.35)	645 (99.38)	3141 (96.94)	
Yes	101 (2.60)	4 (0.62)	97 (2.99)	
Angina pectoris				0.022
No	3811 (97.99)	644 (99.23)	3167 (97.75)	
Yes	75 (1.93)	4 (0.62)	71 (2.19)	
Heart attack				< 0.001
No	3773 (97.02)	644 (99.23)	3129 (96.57)	
Yes	116 (2.98)	5 (0.77)	111 (3.43)	
Stroke				0.002
No	3736 (96.07)	638 (98.31)	3098 (95.62)	
Yes	153 (3.93)	11 (1.69)	142 (4.38)	
Asthma				0.629
No	3262 (83.88)	549 (84.59)	2713 (83.73)	
Yes	627 (16.12)	100 (15.41)	527 (16.27)	
Emphysema				0.011
No	3822 (98.28)	646 (99.54)	3176 (98.02)	
Yes	67 (1.72)	3 (0.46)	64 (1.98)	
Chronic bronchitis				0.135
No	3604 (92.67)	611 (94.14)	2993 (92.38)	
Yes	285 (7.33)	38 (5.86)	247 (7.62)	
Liver condition				0.097
No	3768 (96.89)	636 (98.00)	3132 (96.67)	
Yes	121 (3.11)	13 (2.00)	108 (3.33)	
Failing kidney				0.038
No	3778 (97.15)	639 (98.46)	3139 (96.88)	
Yes	111 (2.85)	10 (1.54)	101 (3.12)	
Kidney stones				< 0.001
No	3582 (92.11)	621 (95.69)	2961 (91.39)	
Yes	307 (7.89)	28 (4.31)	279 (8.61)	
Cancer				< 0.001
No	3493 (89.82)	622 (95.84)	2871 (88.61)	
Yes	396 (10.18)	27 (4.16)	369 (11.39)	
Uterus				< 0.001
No	3493 (89.82)	622 (95.84)	2871 (88.61)	
Yes	28 (0.72)	2 (0.31)	26 (0.80)	
Ovary				0.280
No	3493 (89.82)	622 (95.84)	2871 (88.61)	
Yes	18 (0.46)	3 (0.46)	15 (0.46)	

Table II. Cont.

Variable	Overall	Childbirth history		P-value
		G1 (no)	G2 (yes)	
Cervix				< 0.001
No	3493 (89.82)	622 (95.84)	2871 (88.61)	
Yes	60 (1.54)	2 (0.31)	58 (1.79)	
Breast				< 0.001
No	3493 (89.82)	622 (95.84)	2871 (88.61)	
Yes	111 (2.85)	7 (1.08)	104 (3.21)	
Overweight				< 0.001
No	2406 (61.87)	454 (69.95)	1952 (60.25)	
Yes	1483 (38.13)	195 (30.05)	1288 (39.75)	
Arthritis				< 0.001
No	2634 (67.73)	535 (82.43)	2099 (64.78)	
Yes	1255 (32.27)	114 (17.57)	1141 (35.22)	
Osteoarthritis				< 0.001
No	2634 (95.30)	535 (98.71)	2099 (94.46)	
Yes	130 (4.70)	7 (1.29)	123 (5.54)	
Rheumatoid arthritis				0.033
No	2634 (97.77)	535 (99.07)	2099 (97.45)	
Yes	60 (2.23)	5 (0.93)	55 (2.55)	
Gout				0.036
No	3785 (97.33)	640 (98.61)	3145 (97.07)	
Yes	104 (2.67)	9 (1.39)	95 (2.93)	
Thyroid problem				0.001
No	3284 (84.44)	576 (88.75)	2708 (83.58)	
Yes	605 (15.56)	73 (11.25)	532 (16.42)	
Diabetes				< 0.001
No	3425 (88.07)	611 (94.14)	2814 (86.85)	
Yes	464 (11.93)	38 (5.86)	426 (13.15)	
High blood pressure				< 0.001
No	2441 (62.77)	522 (80.43)	1919 (59.23)	
Yes	1448 (37.23)	127 (19.57)	1321 (40.77)	
Sleep disorder				0.006
No	3593 (92.39)	617 (95.07)	2976 (91.85)	
Yes	296 (7.61)	32 (4.93)	264 (8.15)	

For the 16 diseases of interest, individuals with no disease accounted for 21.5% ($n = 697$), those with one disease made up 22.9% ($n = 741$), with two diseases 19.8% ($n = 641$), with three diseases 11.9% ($n = 385$), and with four diseases 10.3% ($n = 333$). Among all comorbid patients, individuals with two diseases were the most common (29.1%), followed by those with three diseases (25.2%). In total, there were 74 combinations of two diseases. The top three combinations were overweight–high blood pressure (15.2%), arthritis–high blood pressure (12.3%), and overweight–

arthritis (8.0%). There were 113 combinations of three diseases. The top three combinations were overweight–arthritis–high blood pressure (14.5%), overweight–high blood pressure–thyroid problems (4.7%), and overweight–high blood pressure–diabetes (4.4%) (Figure 2 A). In addition, a network graph depicted the comorbidity relationships between diseases. The three largest nodes in the patient network diagram were overweight, hypertension, and arthritis. The strongest association between nodes was observed for overweight–blood pressure–arthritis (Figure 2 B).




Figure 2. Comorbidity patterns among diseases. **A** – UpSet plot showing comorbidity patterns among diseases. **B** – Network graph depicting comorbidity relationships between diseases




Figure 3. Association between childbirth experience and multiple health outcomes. Model 1: unadjusted; Model 2: adjusted for age, race, education, poverty income ratio and BMI

Associations between childbirth experience and multiple health outcomes

We established two models before and after adjusting for confounding factors. As shown in Figure 3, two models established a statistically significant association between childbirth experience and multiple health outcomes. After additionally adjusting for general data confounding factors such as race, education, poverty income ratio, and BMI (Model 2), the risks of seven kinds of medical conditions (heart disease, cancer, kidney stones, arthritis, thyroid problem, diabetes, and high blood pressure) in women who had given birth were significantly higher. The odds ratios (OR) were 1.57–3.54 and 95% confidence intervals (CIs) were 1.15–6.97 ($p < 0.0001$). The three conditions with the highest OR were heart

disease (OR = 3.54, $p < 0.0001$), cancer (OR = 3.12, $p < 0.0001$), and high blood pressure (OR = 2.25, $p < 0.0001$). Next, we conducted the logistic regression analysis of disease subtypes (Supplementary Figure S2). Childbirth experience was associated with higher odds of breast cancer (OR = 3.52, $p = 0.0016$) and cervical cancer (OR = 5.27, $p = 0.0223$), but not uterine cancer (OR = 2.39, $p = 0.2448$) or ovarian cancer (OR = 0.64, $p = 0.4961$). Similarly, reproductive experience was associated with higher odds of osteoarthritis (OR = 4.07, $p = 0.0004$), but not rheumatoid arthritis. Childbirth experience was also associated with higher odds of several cardiovascular conditions, including congestive heart failure, coronary heart disease, and heart attack, but not angina pectoris.

We conducted further analysis of seven conditions that showed significant associations with childbirth experience, including heart disease, cancer, kidney stones, arthritis, thyroid problem, diabetes, and high blood pressure. We examined the association between reproductive indicators (number of live births, age at first birth, age at last birth, and interval between first and last live birth) and these health outcomes. Higher number of live births was associated with increased odds of heart disease, cancer, arthritis, diabetes, and high blood pressure (Figure 4). In contrast, age at first birth, age at last birth, and birth interval were not significantly associated with these outcomes.




Figure 4. Relationship between reproductive indicators and health outcomes

Effects of blood lipid indices

C levels were significantly higher in women with the seven medical conditions ($p < 0.001$, Table III). In contrast, TC, LDL-C, and HDL-C showed hetero-

Table III. Comparison of cholesterol levels

Parameter	N	TC [mmol/l]	LDL-C [mmol/l]	HDL-C [mmol/l]	Remnant cholesterol [mmol/l]
Heart disease					
No	3627	5.081 ±1.038	2.979 ±0.900	1.522 ±0.404	0.581 ±0.311
Yes	262	4.870 ±1.107	2.726 ±0.951	1.421 ±0.439	0.723 ±0.354
P-value		0.002	< 0.001	< 0.001	< 0.001
Kidney stones					
No	3582	5.064 ±1.048	2.961 ±0.908	1.520 ±0.406	0.583 ±0.313
Yes	307	5.098 ±0.999	2.974 ±0.872	1.453 ±0.423	0.672 ±0.343
P-value		0.587	0.806	0.005	< 0.001
Cancer					
No	3493	5.047 ±1.043	2.954 ±0.905	1.512 ±0.400	0.582 ±0.315
Yes	396	5.242 ±1.042	3.035 ±0.905	1.542 ±0.465	0.666 ±0.313
P-value		< 0.001	0.092	0.163	< 0.001
Arthritis					
No	2634	5.021 ±1.041	2.950 ±0.900	1.514 ±0.398	0.556 ±0.304
Yes	1255	5.164 ±1.046	2.986 ±0.915	1.516 ±0.426	0.662 ±0.329
P-value		< 0.001	0.258	0.906	< 0.001
Thyroid problem					
No	3284	5.051 ±1.046	2.956 ±0.906	1.516 ±0.405	0.579 ±0.308
Yes	605	5.153 ±1.032	2.992 ±0.903	1.509 ±0.419	0.652 ±0.349
P-value		0.027	0.370	0.715	< 0.001
Diabetes					
No	3425	5.098 ±1.038	2.994 ±0.897	1.534 ±0.408	0.569 ±0.302
Yes	464	4.839 ±1.060	2.721 ±0.931	1.372 ±0.374	0.746 ±0.367
P-value		< 0.001	< 0.001	< 0.001	< 0.001
High blood pressure					
No	2441	5.032 ±1.035	2.955 ±0.898	1.537 ±0.406	0.541 ±0.289
Yes	1448	5.126 ±1.058	2.974 ±0.917	1.477 ±0.407	0.674 ±0.341
P-value		0.007	0.524	< 0.001	< 0.001

geneous patterns across different conditions; for example, lower levels were observed among women with heart disease and diabetes, whereas higher levels were observed among women with cancer and arthritis. RC levels, however, remained higher in participants with these conditions due to its mathematical derivation from TC, LDL-C, and HDL-C. Restricted cubic spline analyses showed nonlinear associations between RC and several health outcomes, including heart disease, arthritis, cancer, diabetes, and hypertension (Figure 5). No clear nonlinear associations were observed for TC, LDL-C, or HDL-C. In mediation analyses, RC accounted for 6.84–17.7% of the association between childbirth experience and selected health outcomes; however, these results should be interpreted as statistical mediation within a cross-sectional design rather than causal pathways. ROC analysis showed modest discriminatory ability of RC for diabetes (AUC = 0.658) (Figure 6, Supplementary Figure S3).

Discussion

This large cross-sectional study suggests that childbirth history is associated with increased risk of multiple health outcomes in women, and that remnant cholesterol (RC) may mediate part of these associations. These findings contribute to the literature suggesting that reproductive milestones may be associated with long-term cardiovascular and metabolic health in women. Using nationally representative NHANES data, we observed that women with a history of childbirth had higher odds of hypertension, diabetes, cardiovascular conditions, arthritis, and certain cancers (breast and cervical cancer), after adjustment for socioeconomic and anthropometric covariates. RC partially statistically mediated 8.2–17.7% of the observed associations between childbirth history and selected health outcomes. These findings suggest that RC may play a role in explaining part

of the observed associations in this cross-sectional dataset.

Our observation of higher odds of hypertension, diabetes, and cardiovascular diseases aligns with prior studies describing pregnancy as a period of substantial metabolic and hemodynamic change. These physiological adaptations during gestation have been hypothesized in prior studies to be associated with long-term vascular and metabolic alterations. Previous studies have hypothesized that pregnancy is associated with substantial hemodynamic and metabolic adaptations that may have long-term implications for cardiovascu-

lar and metabolic health. In our analysis, a higher number of live births was associated with higher odds of diabetes, and a monotonic association was observed in the unadjusted models. Similar findings have been reported in previous studies, including Moon *et al.* They found that women with higher parity (more than three deliveries) had higher levels of obesity, and their plasma glucose concentrations were elevated at 2 months postpartum. The authors suggested that these findings may be related to pregnancy-associated biological changes, although the underlying mechanisms remain to be clarified [24, 25]. Childbirth history was




Figure 5. Blood remnant cholesterol in the association between childbirth and health outcomes. Restricted cubic spline (RCS) analysis; adjusted for confounders; reference = RC Q1 (< 0.43 mmol/l); L-shaped nonlinear relationship, *p* for nonlinearity < 0.05




Figure 6. Mediation analysis of remnant cholesterol in the association between childbirth and health outcomes. Percentage indicates the proportion of the statistical association mediated by RC; all indirect effects were significant ($p < 0.001$); models were adjusted for confounders

associated with higher odds of breast and cervical cancers, but no significant association with ovarian and uterine cancers was observed. These findings contradicted previous studies [7, 26, 27]. This discrepancy may be related to differences in population characteristics, such as age at first birth and body weight, which have changed over time and are known to be associated with these outcomes.

RC was associated with part of the relationship between childbirth history and multiple health outcomes. Previous studies have suggested that pregnancy is associated with changes in lipid metabolism, although the underlying mechanisms were not examined in this study. In this cross-sectional analysis, RC was more strongly associated with the studied outcomes than LDL-C in the observed dataset. While RC was evaluated as a potential statistical mediator, these findings should be interpreted in the context of the cross-sectional design. Age may partially confound the observed associations given its relationship with both reproductive history and metabolic outcomes. Given that age is

non-modifiable whereas RC represents a potential therapeutic target, exploring its association with reproductive history may be clinically relevant. The cross-sectional design precludes establishing temporal sequence; thus, our mediation analysis should be interpreted as exploratory rather than confirmatory. The observed associations may reflect broader metabolic correlates of lipid profiles, which have been linked in previous studies to inflammatory and metabolic processes [28]. A nonlinear (L-shaped) association between RC and disease risks was observed, with higher odds of several conditions observed at relatively low to moderate RC levels. These findings may have potential implications for future research on lipid monitoring.

Our findings suggest that reproductive history may be considered in future studies of cardiovascular risk stratification. The observed association between childbirth history and higher odds of heart disease (OR = 3.54) indicates a potential relationship that warrants further investigation in prospective studies. RC showed modest discrim-

inatory ability for diabetes (AUC = 0.658). These findings suggest a potential association between RC and diabetes that warrants further evaluation in longitudinal studies.

This study has limitations. First, the cross-sectional design precludes causal inference, and self-reported outcomes may introduce recall bias. While NHANES provides large, nationally representative samples, its continuous cross-sectional structure – independent samples per 2-year cycle without repeated individual measurements – precludes within-person trajectory analysis. Future longitudinal studies should employ prospective cohort designs with repeated measurements of the same women across reproductive milestones to establish the temporal sequence and validate our findings. Second, residual confounding by unmeasured factors (e.g., gestational complications, lifestyle changes postpartum) cannot be excluded. The substantial age difference between nulligravid and parous women (approximately 15 years) represents a significant confounding factor. While our multivariable models adjusted for chronological age, residual confounding from age-related biological aging may persist. Future studies with age-matched controls are needed to disentangle these contributions. Third, several outcomes, including uterine cancer (0.72%), ovarian cancer (0.46%), and emphysema (1.72%), exhibited very low prevalence. While we retained these outcomes given their biological relevance to reproductive history, estimates for rare outcomes carry wide confidence intervals and limited statistical power. These findings should be considered exploratory. Fourth, we recognize that categorizing women into three distinct groups – (1) term live births, (2) pregnancy loss without live births, and (3) nulligravid – would provide finer mechanistic insights. However, the subgroup with pregnancy complications but no live births was too small ($n = 118$, 3.0%) for reliable inference. We therefore retained the binary classification. Fifth, we acknowledge that obesity ($\text{BMI} \geq 30 \text{ kg/m}^2$) represents a more severe clinical endpoint. However, we deliberately retained overweight ($\text{BMI} \geq 25 \text{ kg/m}^2$) as the outcome based on current standards: (1) overweight and obesity remain independent, standardized categories in contemporary national health surveys [29]; (2) recent reproductive epidemiology studies employ $\text{BMI} \geq 25$ as a sensitive endpoint to capture postpartum metabolic deviation [30]; (3) metabolically unhealthy phenotypes are significantly prevalent in the $\text{BMI} 25\text{--}29.9$ range [31]. Future studies should employ standard BMI stratification to clarify dose-response relationships. Finally, we used complete case analysis for missing data, but this may have led to a slight loss of statistical power. Future studies could employ multiple imputation to verify robustness. To overcome the cross-sectional limita-

tion, future research should employ prospective cohort designs with repeated measurements of the same women across reproductive milestones. Such designs would enable: (1) establishment of the temporal sequence; (2) assessment of within-person metabolic changes; (3) evaluation of cumulative pregnancy effects using time-varying covariate analyses; (4) stratification by standard BMI categories to offer additional granularity. Potential data sources include established pregnancy cohorts or electronic health record linkages with longitudinal lipid monitoring.

In conclusion, childbirth history was associated with multiple health outcomes in this cross-sectional study, and RC was statistically associated with part of these relationships. These findings highlight the need for further longitudinal research to clarify the relationships between reproductive history and metabolic health.

Acknowledgments

We thank all of the project participants for their contributions.

Availability of data and materials

The questionnaire and datasets used are available from the corresponding author on request.

Funding

This study was funded by Changzhou Medical Center Project of Nanjing Medical University (CMCC202410), Applied Basic Research Project of Changzhou Science and Technology Bureau (CJ20252043), Changzhou Maternal and Child Health Care Hospital Research project (YJ202602).

Ethical approval and consent to participate

This study is derived from the mining of the NHANES database and is deemed exempt from ethical review and informed consent. The National Center for Health Statistics (NCHS) Ethics Review Board (ERB) ensures that research involving human participants protects the rights and welfare of study participants and conforms to U.S. federal regulations. The NCHS ERB, and the formal review bodies that preceded it, have approved each NHANES study protocol since the survey began running continuously in 1999.

Conflict of interest

The authors declare no conflict of interest

References

1. Hivert MF, Backman H, Benhalima K, et al. Pathophysiology from preconception, during pregnancy, and beyond. *Lancet* 2024; 404: 158-74.

2. Nahae J, Rezaie M, Abdoli E, et al. Association of childbirth experience with long-term psychological outcomes: a prospective cohort study. *Reprod Health* 2024; 21: 71.
3. Countouris M, Mahmoud Z, Cohen JB, et al. Hypertension in pregnancy and postpartum: current standards and opportunities to improve care. *Circulation* 2025; 151: 490-507.
4. Wambua S, Singh M, Okoth K, et al. Association between pregnancy-related complications and development of type 2 diabetes and hypertension in women: an umbrella review. *BMC Med* 2024; 22: 66.
5. Nakimuli A, Okello E, Kesiiga A, et al. Postpartum cardiovascular health in African women following pre-eclampsia: a prospective cohort study. *BJOG* 2025; 132: 1319-28.
6. Feigman MJ, Moss MA, Chen C, et al. Pregnancy reprograms the epigenome of mammary epithelial cells and blocks the development of premalignant lesions. *Nat Commun* 2020; 11: 2649.
7. Husby A, Wohlfahrt J, Melbye M. Pregnancy duration and ovarian cancer risk: a 50-year nationwide cohort study. *Int J Cancer* 2022; 151: 1717-25.
8. Juang CM, Yen MS, Twu NF, et al. Impact of pregnancy on primary dysmenorrhea. *Int J Gynaecol Obstet* 2006; 92: 221-7.
9. Benshop L, Bergen NE, Schalekamp-Timmermans S, et al. Maternal lipid profile 6 years after a gestational hypertensive disorder. *J Clin Lipidol* 2018; 12: 428-436.e4.
10. Adank MC, Benshop L, van Streun SP, et al. Gestational lipid profile as an early marker of metabolic syndrome in later life: a population-based prospective cohort study. *BMC Med* 2020; 18: 394.
11. Su S, Zhang E, Gao S, et al. Associations of remnant cholesterol in early pregnancy with gestational diabetes mellitus risk: a prospective birth cohort study. *Lipids Health Dis* 2024; 23: 243.
12. Wang W, Li N, Wang X, et al. Remnant cholesterol is associated with gestational diabetes mellitus: a cohort study. *J Clin Endocrinol Metab* 2023; 108: 2924-30.
13. Gao Y, Hu Y, Xiang L. Remnant cholesterol, but not other cholesterol parameters, is associated with gestational diabetes mellitus in pregnant women: a prospective cohort study. *J Transl Med* 2023; 21: 531.
14. Adank MC, Bensshop L, Peterbroers KR, et al. Is maternal lipid profile in early pregnancy associated with pregnancy complications and blood pressure in pregnancy and long term postpartum? *Am J Obstet Gynecol* 2019; 221: 150.e1-13.
15. Fang J, Alderman MH. Serum uric acid and cardiovascular mortality the NHANES I epidemiologic follow-up study, 1971-1992. *National Health and Nutrition Examination Survey. JAMA* 2000; 283: 2404-10.
16. Alderman MH, Cohen H, Madhavan S. Dietary sodium intake and mortality: the National Health and Nutrition Examination Survey (NHANES I). *Lancet* 1998; 351: 781-5.
17. Wang Y, Sun Y, Jie T, et al. Association between serum copper-zinc-selenium mixture and multiple health outcomes. *Bioactive Materials* 2025; 50: 432-42.
18. Chen Z, Li R, Guo J, et al. Association between remnant cholesterol (RC) and endometriosis: a cross-sectional study based on NHANES data. *Lipids Health Dis* 2025; 24: 2.
19. Qian S, You S, Sun Y, et al. Remnant cholesterol and common carotid artery intima-media thickness in patients with ischemic stroke. *Circ Cardiovasc Imaging* 2021; 14: e010953.
20. Dang K, Wang X, Hu J, et al. The association between triglyceride-glucose index and its combination with obesity indicators and cardiovascular disease: NHANES 2003-2018. *Cardiovasc Diabetol* 2024; 23: 8.
21. Guo HJ, Ye YL, Gao YF, Liu ZH. Age at first birth is associated with the likelihood of frailty in middle-aged and older women: a population-based analysis from NHANES 1999-2018. *Maturitas* 2024; 181: 107904.
22. Wang L, Wu X, Guo Z, et al. Prolonged sitting time and all-cause mortality: the mediating and predictive role of kidney function markers. *Renal Fail* 2025; 47: 2486568.
23. Chen HL, Wu C, Cao L, et al. The association between the neutrophil-to-lymphocyte ratio and type 2 diabetes mellitus: a cross-sectional study. *BMC Endocr Disord* 2024; 24: 107.
24. Ryczkowska K, Adach W, Janikowski K, et al. Menopause and women's cardiovascular health: is it really an obvious relationship? *Arch Med Sci* 2023; 19: 458-66.
25. Moon JH, Lee J, Kim KH, et al. Multiparity increases the risk of diabetes by impairing the proliferative capacity of pancreatic β cells. *Exp Mol Med* 2023; 55: 2269-80.
26. Husby A, Wohlfahrt J, Øyen N, Melbye M. Pregnancy duration and breast cancer risk. *Nature Commun* 2018; 9: 4255.
27. Husby A, Wohlfahrt J, Melbye M. Pregnancy duration and endometrial cancer risk: nationwide cohort study. *BMJ* 2019; 366: l4693.
28. Ugovšek S, Zupan J, Likožar AR, Sebestjen M. Influence of lipid-lowering drugs on inflammation: what is yet to be done? *Arch Med Sci* 2022; 18: 855-69.
29. Hales CM, Carroll MD, Fryar CD, Ogden CL. Prevalence of obesity and severe obesity among adults: United States, 2017-2018. *NCHS Data Brief* 2020; 360: 18.
30. Guo HJ, Ye YL, Gao YF, Liu ZH. Age at first birth is associated with the likelihood of frailty in middle-aged and older women: a population-based analysis from NHANES 1999-2018. *Maturitas* 2024; 181: 107904.
31. Stefan N. Causes, consequences, and treatment of metabolically unhealthy fat distribution. *Lancet Diabetes Endocrinol* 2020; 8: 616-27.