

Serum triglycerides, remnant cholesterol, and small dense low-density lipoprotein cholesterol levels: lessons from the NATPOL 2011 survey

Bogdan Solnica^{1*}, Adam Wyszomirski², Maciej Banach^{3,4}, Krzysztof Chlebus⁵, Marcin Rutkowski⁶, Piotr Bandosz⁶, Tomasz Zdrojewski⁶

¹Department of Clinical Biochemistry, Jagiellonian University Medical College, Krakow, Poland

²Department of Adult Neurology, Medical University of Gdansk, Poland

³Department of Preventive Cardiology and Lipidology, Medical University of Lodz, Poland

⁴Ciccarone Center for the Prevention of Cardiovascular Disease, Johns Hopkins University School of Medicine, Baltimore, MD, USA

⁵1st Department of Cardiology, Medical University of Gdansk, Poland

⁶Department of Prevention and Didactics, Medical University of Gdansk, Poland

*Corresponding author:

Bogdan Solnica
Department of Clinical Biochemistry
Jagiellonian University Medical College
Skawińska 8
31-066 Krakow, Poland
E-mail: bogdan.solnica@uj.edu.pl

Submitted: 19 May 2026; Accepted: 23 June 2026

Online publication: 23 June 2026

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms/224964>

Copyright © 2026 Termedia & Banach

Abstract

Introduction: The role of triglyceride-rich lipoproteins in atherogenesis is insufficiently recognized and often underestimated. We aimed to evaluate the distribution of serum triglyceride (TG), remnant cholesterol (RC), and small dense LDL cholesterol (sdLDL-C) levels and their interrelationships in participants of the NATPOL 2011 survey.

Material and methods: In approximately 2,000 NATPOL participants, serum TG was measured, and serum RC and sdLDL-C were calculated and compared between women and men, across age groups, and in relation to overweight/obesity and plasma glucose. RC/sdLDL-C levels were also compared across serum TG intervals (< 100, 100–149, and 150–499 mg/dl), and their percentile distributions were analyzed.

Results: Serum TG, RC and sdLDL-C were higher in overweight/obese participants and in those with prediabetes. In subjects with serum TG levels < 100, 100–149, and 150–499 mg/dl, RC and sdLDL-C levels were successively higher, with values exceeding the cutoffs used to indicate high atherosclerotic cardiovascular disease (ASCVD) risk observed in both TG intervals > 100 mg/dl. Participants with serum RC and sdLDL-C above these cutoffs had higher TG. Serum TG levels of 100 mg/dl and 150 mg/dl corresponded to the 55.9th and 88.2nd percentiles, respectively; serum RC of 24 mg/dl corresponded to the 73.8th percentile; and serum sdLDL-C of 30 mg/dl (adopted cutoff) corresponded to the 38th percentile of the respective distributions.

Conclusions: We found an association between serum RC and sdLDL-C, and TG levels > 100 mg/dl indicating that serum TG levels of 100–150 mg/dl may already be associated with TRL remnants and sdLDL accumulation. Calculated serum RC and sdLDL-C may be considered useful supplements to the lipid profile.

Key words: atherosclerotic cardiovascular disease, atherosclerosis, lipid profile, triglyceride-rich lipoproteins.

Introduction

Triglyceride (TG)-rich lipoproteins (TRL) include chylomicrons (CM), very-low-density lipoproteins (VLDL), and the end products of their metabolism, remnants. The association of TRL with atherogenesis and the risk of atherosclerotic cardiovascular disease (ASCVD) is well established [1, 2]. Direct atherogenic effects are exerted by TRL remnants and small dense LDL (sdLDL) produced in increased amounts in hypertriglyceridemia [3–9].

CM, secreted by enterocytes, in the blood, are deprived of TG by lipoprotein lipase (LPL) and enriched in cholesteryl esters transferred from HDL and LDL by cholesteryl ester transfer protein (CETP), and they acquire apolipoprotein E (apoE). The resulting CM remnants are cleared by the liver [3, 4]. VLDL are secreted by hepatocytes in two molecular forms: a larger, TG-rich VLDL1 and a smaller VLDL2, present in roughly equal proportions when plasma TG is below 100 mg/dl, whereas in hypertriglyceridemia, the VLDL1 fraction predominates. VLDL remnants formed after TG removal, unlike CM remnants, are not necessarily cleared by the liver but are mainly converted to LDL, with VLDL1 being considered the main precursor of sdLDL [3, 4].

The structural and functional properties of TRL remnant particles, including their relatively small size and enrichment in cholesteryl esters and apoE, predispose them to participate in atherogenesis. They can enter the arterial intima through transcytosis and are trapped by proteoglycans in the subendothelial space. Remnant particles are engulfed by vascular macrophages through interactions involving the LDL receptors and LDL receptor related protein 1 (LRP-1) mediated by apoE to form foam cells. Additionally, macrophage LPL mediates remnant TG hydrolysis, releasing free fatty acids that may contribute to the inflammatory response [3–7].

Small dense LDL particles, contrary to large buoyant LDL (ldLDL), are characterized by a reduced affinity for the LDL receptor, longer blood residence time, greater susceptibility to oxidation, penetration into the arterial intima, and binding to subendothelial proteoglycans. Oxidized sdLDL particles are taken up by macrophages via scavenger receptors [9, 10].

Many studies, including the PESA (Progression of Early Subclinical Atherosclerosis) study, which showed an association between vascular inflammation and plasma TG ≥ 150 mg/dl [11], have demonstrated that hypertriglyceridemia is a risk factor for ASCVD. In 2021, the European Atherosclerosis Society (EAS) proposed a consensus classification of plasma TG concentrations, considering TG < 100 mg/dl as optimal, and higher values as borderline, moderate, severe, and very severe (ex-

treme) hypertriglyceridemia, with TG levels of 100–150 mg/dl, 150–500 mg/dl, 500–880 mg/dl, and > 880 mg/dl, respectively. This classification is based on the relationship between plasma TG levels and TRL accumulation and the associated ASCVD risk [3]. This recommendation is part of the 2024 Guidelines of the Polish Society of Laboratory Diagnostics and the Polish Lipid Association on laboratory diagnostics of lipid metabolism disorders [12].

However, plasma TG levels reflect TRL metabolism indirectly and approximately. Measurements of plasma remnant or remnant-like particles (RLPs) using ultracentrifugation, electrophoresis, or immunoseparation techniques are available [4] but are not recommended for clinical practice. Plasma remnant cholesterol (RC) can be measured using a direct enzymatic assay or calculated as the difference between total cholesterol and the sum of HDL and LDL cholesterol, or between non-HDL and LDL cholesterol concentrations [2, 4]. However, these measurements and calculations have not been routinely recommended for clinical practice. Similarly, sdLDL-C content in plasma is widely measured for scientific purposes using ultracentrifugation and other techniques but is not quantified in diagnostic practice. In 2021, Sampson *et al.* developed an equation enabling the calculation of plasma sdLDL cholesterol based on the results of a standard lipid profile [13]. In the current study, we evaluated the distribution of serum TG, RC, and sdLDL-C levels and their relationships in the NATPOL 2011 survey participants.

Material and methods

Study population

The NATPOL 2011 (Arterial Hypertension and Other CVD Risk Factors in Poland) survey was a cross-sectional observational study conducted between January and August 2011 to assess the prevalence and control of CVD risk factors. The study cohort was a sample of Polish men and women aged 18–79 [14]. Subjects who declared the use of lipid-lowering treatment were excluded from this analysis. The NATPOL 2011 survey design conformed to the ethical guidelines of the 1975 Declaration of Helsinki and was approved by the Bioethics Committee of the Medical University of Gdansk, Poland. Prior to enrollment, each participant gave informed consent.

Measurements

Blood samples were collected at the participants' homes after 10–12 h of fasting. Frozen serum and plasma samples were transported to the central laboratory, where tests were performed. Serum total cholesterol (TC) was measured by an enzymatic method based on cholesterol esterase

and cholesterol oxidase reactions; HDL-C was measured using Accelerator Selective Detergent (Abbott Laboratories). TG was measured by an enzymatic method with glycerol kinase and glycerol-3-phosphate oxidase. Plasma glucose was measured by the hexokinase/glucose-6-phosphate dehydrogenase method. All tests were performed on the Architect c8000 clinical chemistry analyzer (Abbott Laboratories). Serum LDL-C was calculated using the Friedewald formula: LDL-C [mg/dl] = TC – HDL-C – TG/5 [mg/dl]. When serum TG exceeded 400 mg/dl, the direct LDL-C assay was applied instead of calculation.

Serum RC was calculated as the difference: TC – HDL-C – LDL-C. Small dense LDL cholesterol (sdLDL-C) was calculated as the difference between serum LDL-C and lbLDL-C [13]: lbLDL-C = $1.43 \times \text{LDL-C} - 0.14 \times (\ln \text{TG} \times \text{LDL-C}) - 8.9$ [mg/dl], sdLDL-C = LDL-C – lbLDL-C [mg/dl].

Serum TG was measured in 2,099 survey participants (998 men and 1,101 women); LDL-C, sdLDL-C, lbLDL-C, and RC were determined in 2,073 participants (975 men and 1,098 women); and plasma glucose was measured in 2,088 participants (993 men and 1,095 women).

The serum TG, RC, and sdLDL-C levels were compared between women and men, across age groups, and in relation to body mass index (BMI), visceral obesity, and fasting plasma glucose (FPG). The distribution of serum RC, LDL-C, lbLDL-C, and sdLDL-C levels was compared among subgroups of survey participants with optimal serum TG, borderline, and moderate hypertriglyceridemia, in line with the 2021 EAS consensus statement [3]. No cases of severe (500–880 mg/dl) or very severe (> 880 mg/dl) hypertriglyceridemia were found in the survey cohort. Based on clinical trial reports, serum RC of 24 mg/dl and sdLDL-C of 30 mg/dl were adopted as cutoff values for increased ASCVD risk [4, 15–20]. Serum TG distribution was analyzed in participants with RC and sdLDL-C levels below and above the adopted cutoff values. The percentile distribution of serum TG, RC, LDL-C, and sdLDL-C was also analyzed.

Statistical analysis

Statistical analyses were conducted using R software, version 4.4.2 (R Core Team). Continuous

variables were reported as means accompanied by 95% confidence intervals (CIs), calculated based on the *t*-distribution. All subgroups analyzed were defined a priori. To evaluate the statistical significance of differences between subgroup means, the criterion of non-overlapping 95% confidence intervals was employed. It is important to note that this approach is more conservative compared to a formal hypothesis test at a significance level of $\alpha = 0.05$. Missing data were addressed through complete-case analysis, whereby observations with missing values for the variables of interest were excluded. The analysis did not adjust for multiple comparisons, and the complex survey design was not taken into account.

Results

In the survey cohort, we observed a similar distribution of serum TG, RC, and sdLDL-C, with minor differences mainly between age groups (Table I). In the entire cohort, and independently in men and women, serum TG, RC, and sdLDL-C were higher in those with overweight/obesity compared with those with BMI < 25 kg/m², and in those with rather than without visceral obesity. Serum TG and RC in all BMI categories, and serum sdLDL-C in subjects with BMI > 30 kg/m², were higher in men than in women. In the entire cohort, and independently in men and women, serum TG was higher in subjects with impaired fasting glucose (IFG) and plasma glucose indicative of diabetes than in those with those with normal fasting glucose (NFG), while serum RC and sdLDL-C were higher in subjects with IFG than in those with NFG (Table I).

In the subgroups with serum TG < 100 mg/dl, 100–149 mg/dl, 150–499 mg/dl, serum RC was consecutively and significantly higher (Table II). In subsequent serum TG groups, the concentrations of LDL-C and sdLDL-C were higher, while serum lbLDL-C did not differ significantly (Table II). Additionally, across these three serum TG categories, the mean sdLDL-C levels were 24.7%, 31%, and 37.2% of the mean serum LDL-C, respectively. Of note, in both analyzed TG intervals above 100 mg/dl, the study participants had serum RC and sdLDL-C levels of ≥ 24 mg/dl and ≥ 30 mg/dl, respectively (Table II, Figure 1). On the other hand,

Table I. Serum triglyceride, remnant cholesterol, and small dense LDL cholesterol levels in the survey participants

Groups	Triglycerides:			Remnant cholesterol:			Small dense LDL cholesterol:		
	N, mean (95% CI) [mg/dl]			N, mean (95% CI) [mg/dl]			N, mean (95% CI) [mg/dl]		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
The entire cohort	998	1101	2099	975	1098	2073	975	1098	2073
	120.70	96.95	108.24	22.30	19.19	20.65	36.90	34.06	35.40
	(115.43,	(94.32,	(105.34,	(21.66,	(18.72,	(20.26,	(36.12,	(33.40,	(34.89,
	125.97)	99.58)	111.14)	22.94)	19.65)	21.04)	37.69)	34.72)	35.91)

Table I. Cont.

Groups	Triglycerides: N, mean (95% CI) [mg/dl]			Remnant cholesterol: N, mean (95% CI) [mg/dl]			Small dense LDL cholesterol: N, mean (95% CI) [mg/dl]		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Age groups									
18–39	411	471	882	400	471	871	400	471	871
	116.58	85.17	99.80	21.45	17.04	19.07	33.82	29.46	31.46
	(108.83, 124.33)	(81.94, 88.39)	(95.68, 103.93)	(20.48, 22.42)	(16.40, 17.69)	(18.48, 19.65)	(32.71, 34.93)	(28.66, 30.27)	(30.78, 32.15)
40–59	365	350	715	355	348	703	355	348	703
	130.27	100.67	115.78	23.56	19.64	21.62	40.75	36.97	38.88
	(119.99, 140.56)	(95.30, 106.03)	(109.82, 121.74)	(22.45, 24.67)	(18.83, 20.46)	(20.91, 22.33)	(39.38, 42.12)	(35.79, 38.15)	(37.96, 39.79)
60–79	222	280	502	220	279	499	220	279	499
	112.60	112.12	112.33	21.81	22.23	22.05	36.30	38.21	37.37
	(104.35, 120.85)	(106.98, 117.26)	(107.71, 116.96)	(20.55, 23.08)	(21.27, 23.20)	(21.28, 22.82)	(34.69, 37.92)	(36.82, 39.59)	(36.32, 38.42)
Body mass index (BMI)									
< 25 kg/m ²	345	569	914	342	569	911	342	569	911
	99.39	86.76	91.53	19.27	17.35	18.07	32.82	31.42	31.95
	(92.69, 106.09)	(83.71, 89.80)	(88.35, 94.70)	(18.32, 20.22)	(16.74, 17.96)	(17.55, 18.59)	(31.68, 33.96)	(30.57, 32.27)	(31.26, 32.63)
25–29 kg/m ²	425	324	749	412	321	733	412	321	733
	125.76	106.45	117.41	23.04	20.61	21.97	38.44	36.78	37.71
	(118.20, 133.31)	(100.64, 112.26)	(112.40, 122.41)	(22.07, 24.00)	(19.77, 21.44)	(21.31, 22.63)	(37.20, 39.67)	(35.52, 38.04)	(36.82, 38.60)
≥ 30 kg/m ²	222	200	422	215	200	415	215	200	415
	144.86	110.60	128.63	25.80	22.13	24.03	40.60	37.11	38.92
	(129.90, 159.83)	(104.75, 116.45)	(120.15, 137.10)	(24.33, 27.28)	(20.96, 23.30)	(23.07, 25.00)	(38.84, 42.36)	(35.57, 38.65)	(37.73, 40.10)
Visceral obesity									
Yes	590	653	1243	573	650	1223	573	650	1223
	135.88	105.22	119.77	24.81	20.71	22.63	39.90	36.27	37.97
	(128.31, 143.45)	(101.58, 108.86)	(115.62, 123.93)	(23.93, 25.69)	(20.10, 21.31)	(22.09, 23.17)	(38.85, 40.95)	(35.41, 37.12)	(37.30, 38.65)
No	408	444	852	402	444	846	402	444	846
	98.75	84.39	91.26	18.72	16.88	17.75	32.63	30.69	31.61
	(92.51, 104.99)	(81.00, 87.77)	(87.77, 94.76)	(17.94, 19.50)	(16.20, 17.55)	(17.24, 18.27)	(31.57, 33.69)	(29.73, 31.65)	(30.90, 32.33)
Cardiovascular disease									
Yes	96	98	194	93	97	190	93	97	190
	130.11	105.83	117.85	23.39	20.62	21.97	34.10	35.32	34.72
	(110.58, 149.65)	(97.14, 114.51)	(107.19, 128.50)	(21.10, 25.68)	(19.26, 21.97)	(20.65, 23.29)	(31.68, 36.52)	(33.31, 37.33)	(33.17, 36.28)
No	902	1003	1905	882	1001	1883	882	1001	1883
	119.70	96.08	107.26	22.18	19.05	20.52	37.20	33.94	35.47
	(114.24, 125.15)	(93.32, 98.84)	(104.26, 110.27)	(21.52, 22.85)	(18.55, 19.54)	(20.10, 20.93)	(36.37, 38.03)	(33.24, 34.64)	(34.92, 36.01)
Fasting plasma glucose									
NFG (< 100 mg/dl)	721	927	1648	708	925	1633	708	925	1633
	112.57	93.71	101.96	21.28	18.56	19.74	35.90	33.36	34.47
	(107.55, 117.59)	(90.92, 96.51)	(99.23, 104.70)	(20.60, 21.97)	(18.07, 19.05)	(19.33, 20.15)	(35.01, 36.80)	(32.65, 34.08)	(33.90, 35.03)
IFG (100–125 mg/dl)	217	146	363	213	145	358	213	145	358
	138.06	114.23	128.48	25.88	22.48	24.50	40.73	38.45	39.81
	(126.12, 150.00)	(106.76, 121.71)	(120.67, 136.28)	(24.22, 27.54)	(21.14, 23.81)	(23.37, 25.63)	(38.95, 42.52)	(36.62, 40.27)	(38.52, 41.10)
Indicative of diabetes (≥ 126 mg/dl)	55	22	77	49	22	71	49	22	71
	162.73	121.32	150.90	22.16	24.32	22.83	35.62	35.33	35.53
	(112.87, 212.59)	(97.75, 144.89)	(114.79, 187.00)	(19.96, 24.37)	(19.58, 29.06)	(20.77, 24.89)	(32.24, 39.00)	(30.06, 40.59)	(32.75, 38.30)

Table II. Distribution of serum remnant cholesterol (RC), low-density lipoprotein cholesterol (LDL-C), small dense LDL cholesterol (sdLDL-C), and large buoyant LDL cholesterol (lbLDL-C) among subjects with optimal triglyceridemia, borderline, and moderate hypertriglyceridemia (EAS 2021)


Fasting serum triglycerides	Remnant cholesterol: <i>N</i> , mean (95% CI) [mg/dl]			LDL cholesterol: <i>N</i> , mean (95% CI) [mg/dl]			sdLDL cholesterol: <i>N</i> , mean (95% CI) [mg/dl]			lbLDL cholesterol: <i>N</i> , mean (95% CI) [mg/dl]		
	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total
< 100 mg/dl optimal	484	669	1153	484	669	1153	484	669	1153	484	669	1153
	14.95 (14.67, 15.24)	14.32 (14.07, 14.56)	14.58 (14.40, 14.77)	116.43 (113.58, 119.28)	118.33 (115.99, 120.67)	117.53 (115.73, 119.34)	29.14 (28.47, 29.82)	28.65 (28.11, 29.20)	28.86 (28.43, 29.28)	87.29 (85.02, 89.56)	89.68 (87.79, 91.57)	88.67 (87.22, 90.13)
100–149 mg/dl borderline hypertriglyceridemia	338	353	691	338	353	691	338	353	691	338	353	691
	24.51 (24.21, 24.81)	24.20 (23.90, 24.50)	24.35 (24.14, 24.57)	131.00 (127.24, 134.77)	130.94 (127.09, 134.80)	130.97 (128.28, 133.66)	40.79 (39.81, 41.76)	40.53 (39.54, 41.51)	40.65 (39.96, 41.35)	90.22 (87.39, 93.05)	90.41 (87.51, 93.32)	90.32 (88.29, 92.34)
150–499 mg/dl moderate hypertriglyceridemia	153	76	229	153	76	229	153	76	229	153	76	229
	40.65 (39.18, 42.13)	38.75 (37.01, 40.49)	40.02 (38.88, 41.16)	141.42 (135.3, 147.48)	139.58 (130.45, 148.71)	140.81 (135.79, 145.83)	52.88 (50.87, 54.90)	51.66 (48.66, 54.66)	52.48 (50.82, 54.14)	88.54 (84.32, 92.76)	87.92 (81.62, 94.22)	88.33 (84.85, 91.82)

survey participants with serum RC ≥ 24 mg/dl and sdLDL-C ≥ 30 mg/dl had higher TG levels compared with participants with levels below these cutoff values (Table III).

Percentile distribution analysis showed that the median values of serum TG, RC, LDL-C, and sdLDL-C were 95 mg/dl, 19 mg/dl, 122 mg/dl, and 33.5 mg/dl, respectively. Serum TG values of 100 mg/dl and 150 mg/dl corresponded to the 55.9th and 88.2nd percentiles, respectively; serum RC of 24 mg/dl was at the 73.8th percentile, serum LDL-C of 100 mg/dl was at the 26.1st percentile, and serum sdLDL-C of 30 mg/dl was at the 38th percentile (Table IV).

Discussion

TRL remnants, together with LDL and lipoprotein (a) [Lp(a)], belong to the atherogenic lipoproteins directly involved in all stages of atherogenesis. Therefore, their elevated plasma levels should

**Figure 1.** Mean (95% CI) serum remnant cholesterol (RC) and small dense LDL cholesterol (sdLDL-C) across triglyceride level intervals according to the EAS [3]

be considered separate risk factors, also because the treatment of these forms of dyslipidemia differs [21, 22]. For many years, plasma/serum TG

Table III. Serum TG in participants with remnant cholesterol (RC) and small dense LDL cholesterol (sdLDL-C) levels below and above 24 mg/dl and 30 mg/dl, respectively

Serum triglycerides	Male	Female	Total
Serum RC 24–69 mg/dl			
<i>N</i>	355	265	620
Mean (95% CI)	162.52 (157.61, 167.44)	149.91 (145.63, 154.19)	157.13 (153.75, 160.51)
RC < 24 mg/dl			
<i>N</i>	620	833	1453
Mean (95% CI)	82.13 (80.56, 83.71)	82.13 (80.56, 83.71)	78.76 (77.36, 80.15)
Serum sdLDL-C 30–111 mg/dl			
<i>N</i>	648	636	1284
Mean (95% CI)	129.67 (125.81, 133.52)	114.51 (111.51, 117.51)	122.16 (119.68, 124.64)
Serum sdLDL-C < 30 mg/dl			
<i>N</i>	326	462	788
Mean (95% CI)	74.62 (71.94, 77.29)	74.62 (71.94, 77.29)	70.35 (68.29, 72.41)

Table IV. Percentile distribution of serum triglyceride (TG), remnant cholesterol (RC), low-density lipoprotein cholesterol (LDL-C), and small dense LDL cholesterol (sdLDL-C) levels (mg/dl) in the entire survey cohort

Percentile	Serum TG	Serum RC	Serum LDL-C	Serum sdLDL-C
5 th	49	10	73	19.69
10 th	57	11	83	21.97
25 th	72	14	99	26.69
25.1 st			100	
38 th				30
50 th (median)	95	19	122	33.50
55.9 th	100			
73.8 th		24		
75 th	125	25	146	42.13
88.2 nd	150			
90 th	160	31	169	51.23
95 th	206	38	187	56.76




Figure 2. Relationship between serum small dense LDL cholesterol (sdLDL-C) and triglyceride levels. Serum sdLDL-C above 30 mg/dl was observed in borderline and moderate hypertriglyceridemia according to the EAS [3]

was considered a measure of TRL and their remnants' content in the blood. In principle, this rule is correct. Recently, this relationship has been described more accurately – the EAS proposed shifting optimal TG levels to < 100 mg/dl and defined higher levels as progressive degrees of hypertriglyceridemia, linking serum TG levels with TRL accumulation and the associated ASCVD risk [3, 12].

However, serum TG levels are an indirect and not always accurate measure of TRL remnant formation and content in plasma. TRL metabolism, mediated mainly by LPL and hepatic lipase (HL), is affected by many factors, including the regulation of the synthesis and activity of LPL, leading to inter-individual variability [23]. Therefore, serum TG alone may be insufficient in this context. An extreme example is familial chylomicronemia – the accumulation of CM and very high plasma TG due to a lack of LPL functionality without increased ASCVD risk in affected patients [24].

Therefore, plasma/serum RC and sdLDL-C determined together with TG levels may provide additional information for assessing TRL-related

ASCVD risk and may help inform decisions about TG-lowering treatment. Some TRL remnants or remnant-like particles (RLP) determination methods are available to clinical laboratories [4, 6]. Calculation of plasma/serum RC and sdLDL-C is considered a convenient tool because dedicated formulas use the results of a standard lipid profile [4, 13]. We used these formulas in our study.

We analyzed the relationship between serum RC and sdLDL-C and TG levels based on their distribution in approximately 2,000 NATPOL 2011 participants, representative of the Polish population aged 18–79. We found that serum TG, RC, and sdLDL-C were higher in older age groups, in subjects with overweight/obesity, including visceral obesity, and in subjects with IFG (prediabetes) compared to survey participants in the 18–39 age group, with BMI < 25 kg/m², without visceral obesity, and with NFG, respectively. Furthermore, serum TG, RC, and sdLDL-C were higher in men than in women (Table I). The relationship between TRL metabolism and serum TG levels with age is well established and explained [25, 26]. Also, in many studies involving subjects with metabolic syndrome, hypertriglyceridemia with a parallel increase in serum RC and sdLDL-C has been observed, which is consistent with our findings [27–31].

In the categories of serum TG proposed by EAS, RC levels increased gradually and differed significantly between subjects with optimal serum TG, borderline, and moderate hypertriglyceridemia. Of note, subjects with borderline hypertriglyceridemia had serum RC close to the adopted cutoff value for increased ASCVD risk (24 mg/dl) (Figure 1, Table II). The equation used in our study to calculate serum RC as the transformed Fredrickson formula closely relates RC and TG levels, with a regression equation close to $RC = 0.2 \times TG$ [mg/dl]

and a correlation coefficient of 0.999 found. To keep this calculation accurate, it is recommended to use LDL-C levels measured directly (at least in subjects with hypertriglyceridemia, as in our study) or obtained using the Martin-Hopkins or Sampson NIH equation in this formula [4, 5, 32–35]. The relationship, together with the distribution of RC levels in participants with borderline hypertriglyceridemia, indicates that serum RC close to or higher than 24 mg/dl can be expected within this TG level interval. Furthermore, there are data showing that at concentrations > 20 mg/dl, directly measured serum RC may be higher than the calculated one [32].

Serum sdLDL-C also increased and differed significantly between successive TG level categories, exceeding, in borderline hypertriglyceridemia, the adopted cutoff value, with a mean of 40.65 mg/dl (Table II, Figures 1 and 2). Increases in serum sdLDL-C across successive TG concentration intervals were the main contributor to the increase in serum LDL-C because serum lLDL-C did not differ between these intervals. Across the serum TG intervals, the mean sdLDL-C levels increased both in absolute values and relatively, as a percentage of mean serum LDL-C (Table II). These findings support the hypothesis that sdLDL is a second link, beyond TRL remnants, between increased serum TG with TRL accumulation and ASCVD risk.

On the other hand, in survey participants, serum RC $\geq$ 24 mg/dl and sdLDL-C $\geq$ 30 mg/dl were associated with higher TG levels compared to participants whose serum RC and sdLDL-C levels were below these cutoffs. TG levels in subjects with serum RC $\geq$ 24 mg/dl and sdLDL-C $\geq$ 30 mg/dl were within moderate and borderline/moderate hypertriglyceridemia, respectively (Table III). Our two-way analysis showed associations between serum RC and sdLDL-C and TG levels, consistent with the triglyceridemia classification proposed by the EAS [3]. Such data shed new light on atherogenic dyslipidemia, previously defined by high serum TG and low HDL-C, and currently by increased non-HDL-C reflecting TRL remnants accumulation [36].

Dyslipidemia is the most common modifiable risk factor for ASCVD. The NATPOL 2011 survey revealed hypercholesterolemia (LDL-C > 115 mg/dl) in 57.8% of the Polish population aged 18–79 [37]. A similar incidence of hypercholesterolemia has also been observed in other countries [38]. In our present study, percentile-distribution analysis showed that serum TG levels > 100 mg/dl and > 150 mg/dl were found in 44.1% and 11.8% of participants, respectively (in Poland this corresponds to approx. 14 million and 4 million people, respectively). Serum RC > 24 mg/dl was found in 26.2% (in Poland, approx. 9 million people); and serum sdLDL-C > 30 mg/dl was found in 62% of

participants. Of note, this frequency was close to the frequency of LDL-C > 115 mg/dl – in Poland, approximately 20 million people. These numbers highlight the potential population-level relevance of TRL-related ASCVD risk.

To the best of our knowledge, the present study is the first analysis of the distribution of RC and sdLDL-C in a cohort intended to be representative of the Polish population. Standardized preanalytical procedures and centralized laboratory testing ensured high-quality measurements.

The time elapsed since the survey was completed may be considered a limitation of our study. The motivation for this study was the 2021 EAS consensus statement [3]. Taking into account the increasing prevalence of CVD risk factors such as obesity, diabetes, and metabolic syndrome, we may assume that the epidemiological situation, including the distribution of RC and sdLDL-C in the population, may have worsened since the survey [39, 40].

In conclusion, our results show an association between serum RC and sdLDL-C levels and TG, including at TG levels exceeding 100 mg/dl. These findings should draw attention to borderline hypertriglyceridemia (100–150 mg/dl), until recently considered normal, as a possible state of increased RC and/or sdLDL-C levels. Calculation of serum RC and sdLDL-C, although still under discussion, is a convenient tool to characterize lipid profiles that may reflect increased plasma concentrations of these atherogenic lipoprotein fractions. Taking this further, these measures could potentially contribute to personalized assessment of serum TG in the context of ASCVD risk. Supplementing the standard lipid profile with these calculations could allow for a more comprehensive assessment of dyslipidemia-related residual ASCVD risk. However, further research is required to evaluate the diagnostic characteristics of RC and sdLDL-C.

Funding

The NATPOL 2011 Survey was partially funded by the Polish Ministry of Health as part of the National CVD Prevention and Treatment Program and by statutory grants from the Medical University of Gdansk and the Medical University of Warsaw. It was also partly funded by the following industry sponsors: Sanofi-Aventis, Abbott Laboratories Poland Ltd, Siemens Ltd., and Polpharma.

Ethical approval

Approval number: NKEBN/251/99.

Conflict of interest

The authors declare no conflict of interest.

References

- Nordestgaard BG. Triglyceride-rich lipoproteins and atherosclerotic cardiovascular disease new insights from epidemiology, genetics, and biology. *Circ Res* 2016; 118: 547-63.
- Xu D, Xie L, Cheng C, et al. Triglyceride-rich lipoproteins and cardiovascular diseases. *Front Endocrinol* 2024; 15: 1409653.
- Ginsberg HN, Packard CJ, Chapman MJ, et al. Triglyceride-rich lipoproteins and their remnants: metabolic insights, role in atherosclerotic cardiovascular disease, and emerging therapeutic strategies – a consensus statement from the European Atherosclerosis Society. *Eur Heart J* 2021; 42: 4791-806.
- Durana EK, Pradhanb AD. Triglyceride-rich lipoprotein remnants and cardiovascular disease. *Clin Chem* 2021; 67: 183-97.
- Balling M, Afzal S, Varbo A, et al. Remnant cholesterol: quantification, concentrations by sex and age, and risk of ischemic heart disease. *Clin Chem* 2025; 71: 451-62.
- Zhao Y, Zhuang Z, Li Y, et al. Elevated blood remnant cholesterol and triglycerides are causally related to the risks of cardiometabolic multimorbidity. *Nature Com* 2024; 15: 2451.
- Baratta F, Cocomello N, Coronati M, et al. Cholesterol remnants, triglyceride-rich lipoproteins and cardiovascular risk. *Int J Mol Sci* 2023; 24: 4268.
- Duran EK, Aday AW, Cook NR, et al. Triglyceride-rich lipoprotein cholesterol, small dense LDL cholesterol, and incident cardiovascular disease. *JACC* 2020; 75: 122-35.
- Krauss RM. Small dense LDL particles: clinically relevant? *Cur Opin Lipidol* 2022; 33: 160-6.
- Vekic J, Zeljkovic A, Cicero AFG, et al. Atherosclerosis development and progression: the role of atherogenic small, dense LDL. *Medicina* 2022; 58: 299.
- Raposeiras-Roubin S, Rossello X, Oliva B, et al. Triglycerides and residual atherosclerotic risk. *J Am Coll Cardiol* 2021; 77: 3031-41.
- Solnica B, Sygitowicz G, Sitkiewicz D, et al. 2024 Guidelines of the Polish Society of Laboratory Diagnostics and the Polish Lipid Association on laboratory diagnostics of lipid metabolism disorders. *Arch Med Sci* 2024; 20: 357-74.
- Sampson M, Wolska A, Warnick R, et al. New equation based on the standard lipid panel for calculating small dense low-density lipoprotein-cholesterol and its use as a risk-enhancer test. *Clin Chem* 2021; 67: 987-97.
- Zdrojewski T, Rutkowski M, Bandosz P, et al. Prevalence and control of cardiovascular risk factors in Poland. Assumptions and objectives of the NATPOL 2011 Survey. *Kardiol Pol* 2013; 71: 381-92.
- Quispe R, Martin SS, Michos ED, et al. Remnant cholesterol predicts cardiovascular disease beyond LDL and ApoB: a primary prevention study. *Eur Heart J* 2021; 42: 4324-32.
- Varbo A, Freiberg JJ, Nordestgaard BG. Remnant cholesterol and myocardial infarction in normal weight, overweight, and obese individuals from the Copenhagen General Population Study. *Clin Chem* 2018; 64: 1219-30.
- Doi T, Langsted A, Nordestgaard BG. Elevated remnant cholesterol reclassifies risk of ischemic heart disease and myocardial infarction. *J Am Coll Cardiol* 2022; 79: 2383-97.
- Ikezaki H, Lim E, Cupples LA, et al. Small dense low-density lipoprotein cholesterol is the most atherogenic lipoprotein parameter in the prospective Framingham Offspring Study. *J Am Heart Assoc* 2021; 10: e019140.
- Inyaku M, Tanaka M, Sato T, et al. Calculated small dense low-density lipoprotein cholesterol level is a pre-dominant predictor for new onset of ischemic heart disease. *J Atheroscler Thromb* 2024; 31: 232-48.
- Koba S, Satoh N, Ito Y, et al. Impact of direct measurement of small dense low-density lipoprotein cholesterol for long-term secondary prevention in patients with stable coronary artery disease. *Clin Chem* 2024; 70: 957-66.
- Toth PP, Banach M. It is time to address the contribution of cholesterol in all apoB-containing lipoproteins to atherosclerotic cardiovascular disease. *Eur Heart J Open* 2024; 4: oeae057.
- Makover ME, Surma S, Banach M, et al. Eliminating atherosclerotic cardiovascular disease residual risk. *Eur Heart J* 2023; 44: 4731-3.
- Wu SA, Kersten S, Qi L. Lipoprotein lipase and its regulators: an unfolding story. *Trends Endocrinol Metab* 2021; 32: 48-61.
- Moulin P, Dufour R, Avernac M, et al. Identification and diagnosis of patients with familial chylomicronaemia syndrome (FCS): expert panel recommendations and proposal of an "FCS score". *Atherosclerosis* 2018; 275: 265-72.
- Spitler KM, Davies BSJ. Aging and plasma triglyceride metabolism. *J Lipid Res* 2020; 61: 1161-7.
- Feng L, Nian S, Tong Z, et al. Age-related trends in lipid levels: a large-scale cross-sectional study of the general Chinese population. *BMJ Open* 2020; 10: e034226.
- ZhangID F, Zhang Y, Zhang J, et al. Small dense low density lipoprotein predominance in patients with type 2 diabetes mellitus using Mendelian randomization. *PLoS One* 2024; 19: e0298070.
- Li B, Liu Y, Zhou X, Chen L, et al. Remnant cholesterol is more positively related to diabetes, prediabetes, and insulin resistance than conventional lipid parameters and lipid ratios: a multicenter, large sample survey. *J Diabetes* 2024; 16: e13592.
- Hirano T. Hypertriglyceridemia contributes significantly to high prevalence of small dense LDL-cholesterol in patients with type 2 diabetes, even when LDL-C targets are achieved. *Diabetol Int* 2023; 14: 288-93.
- Srisawasdi P, Vanavanan S, Rochanawutanon M, et al. Small-dense LDL/large-buoyant LDL ratio associates with the metabolic syndrome. *Clin Biochem* 2015; 48: 495-502.
- Hirano T, Satoh N, Ito Y. Specific increase in small dense low-density lipoprotein-cholesterol levels beyond triglycerides in patients with diabetes: implications for cardiovascular risk of MAFLD. *J Atheroscler Thromb* 2024; 31: 36.
- Wadstrom BN, Pedersen KM, Wulf AB, et al. Elevated remnant cholesterol in atherosclerotic cardiovascular disease in diabetes: a population-based prospective cohort study. *Eur Heart J* 2023; 44 (Suppl 2).
- Langlois MR, Nordestgaard BG, Langsted A, et al.; European Atherosclerosis Society (EAS) and the European Federation of Clinical Chemistry and Laboratory Medicine (EFLM). Joint Consensus Initiative Quantifying atherogenic lipoproteins for lipid-lowering strategies: consensus-based recommendations from EAS and EFLM. *Clin Chem Lab Med* 2020; 58: 496-517.
- Grant JK, Kaufman HW, Martin SS. Extensive evidence supports the Martin–Hopkins equation as the LDL-C calculation of choice. *Clin Chem* 2024; 70: 392-8.
- Sampson M, Wolska A, Meeusen JW, et al. The Sampson-NIH equation is the preferred calculation method for LDL-C. *Clin Chem* 2024; 70: 399-402.

36. Raja V, Aguiar C, Alsayed N, et al. Non-HDL-cholesterol in dyslipidemia: review of the state-of-the-art literature and outlook. *Atherosclerosis* 2023; 383: 117312.
37. Zdrojewski T, Solnica B, Cybulska B, et al. Prevalence of lipid abnormalities in land. The NATPOL 2011 survey. *Kardiologia Polska* 2016; 74: 213-23.
38. NCD Risk Factor Collaboration (NCD-RisC). Repositioning of the global epicenter of non-optimal cholesterol. *Nature* 2020; 582 (7810): 73-7.
39. Chen Y, Gue Y, Banach M, et al.; LIPIDOGRAM Investigators. Phenotypes of Polish primary care patients using hierarchical clustering: Exploring the risk of mortality in the LIPIDOGEM2015 study cohort. *Eur J Clin Invest* 2024; 54: e14261.
40. Osadnik T, Nowak D, Osadnik K, et al.; LIPIDOGRAM Investigators. Association of body mass index and long-term mortality in patients from nationwide LIPIDOGRAM 2004-2015 cohort studies: no obesity paradox? *Cardiovasc Diabetol* 2023; 22: 323.